

## A SIMPLE ANALYTIC APPROXIMATION FOR DUSTY STRÖMGREN SPHERES

VAHÉ PETROSIAN

Institute for Plasma Research, Stanford University, Stanford, California

AND

JOSEPH SILK AND GEORGE B. FIELD

Astronomy Department, University of California, Berkeley

*Received 1972 July 12; revised 1972 July 31*

### ABSTRACT

We interpret recent far-infrared observations of H II regions in terms of true absorption by internal dust of a significant fraction of the Lyman-continuum photons. We present approximate analytic expressions describing the effects of internal dust on the ionization structure of H II regions, and outline a procedure for deducing the properties of this dust from optical and infrared observations.

The discovery of large far-infrared emissions from H II regions (Harper and Low 1971), significantly in excess of the apparent Lyman-continuum output of the exciting stars inferred from measurements of the thermal radio continuum, on the assumption that the Lyman-continuum photons are absorbed by the gas has stimulated us to reexamine the effects of internal dust on Strömgren's classical solution to the problem of H II regions. It is the purpose of this *Letter* to indicate a simple analytic approximation, which leads to results in reasonable agreement with earlier numerical work (Mathis 1971). Our result moreover both clarifies and adds considerable insight into the problem, and is useful in indicating how one may extend Mathis's results over a wide range of optical depths for dust absorption, with no additional numerical computation. Moreover, we apply our results to interpret the observed far-infrared emission from H II regions, and demonstrate the observations to be consistent with a moderate amount of absorption of Lyman-continuum photons by internal dust.

The standard equation of ionization equilibrium for a spherically symmetric H II region of uniform density  $n$ , containing a central exciting star emitting  $S$  ionizing quanta per second can be written in the form

$$\frac{1-x}{x^2} = \frac{3y^2 e^\tau}{\tau_s}, \quad (1)$$

where  $x$  is the fractional degree of ionization,  $y \equiv r/r_s$  is the dimensionless radial coordinate in units of the Strömgren radius, which is  $r_s$ , defined by

$$S = \frac{4\pi}{3} r_s^3 n^2 a, \quad (2)$$

and

$$\tau_s = n\sigma r_s. \quad (3)$$

In these equations  $a$  is the hydrogen recombination coefficient summed over all levels excluding the ground level, and  $\sigma$  is an appropriately averaged value of the photoionization cross-section for hydrogen. The optical depth  $\tau$  is defined by

$$\frac{d\tau}{dy} = (1-x)\tau_s + \tau_d, \quad (4)$$

where

$$\tau_d = n\sigma_g x_g r_s, \quad (5)$$

$\sigma_g$  is the effective absorption cross-section of the grains, and  $x_g$  is the number of dust grains per H atom. The dust and gas are assumed to be uniformly distributed in the ionized region.

We make the approximation  $1-x \ll 1$ , which should be well satisfied throughout the ionized region. It is then a straightforward matter to eliminate  $\tau$  from equations (1) and (4), and obtain the following solution for  $x$  as a function of radial coordinate:

$$\frac{1}{1-x} = \frac{\tau_s}{3y^2} \left\{ \exp(-\tau_d y) + \frac{6}{\tau_d^3} [\exp(-\tau_d y) - 1 + \tau_d y - \frac{1}{2} \tau_d^2 y^2] \right\}. \quad (6)$$

For  $\tau_d \ll 1$ , equation (6) reduces to the anticipated limit

$$\frac{1}{1-x} \sim \frac{\tau_s}{3y^2} [1 - y^3 - \tau_d y(1 - y^{3/4}) + O(\tau_d^2)]. \quad (7)$$

The neutral fraction  $1-x$  is plotted in figure 1 as a function of  $y$  for various values of  $\tau_d$ . The parameter  $\tau_s$  is set equal to 5000. For comparison, the numerical results of Mathis (1971) are also shown (*dashed line*) for one of the cases he computed, namely,  $\tau_d = 0.4$ . The agreement for the ionization structure is remarkably good, despite the approximations of our analysis. The small disagreement at low values of  $y$  is due to scattering of photons by dust which we have ignored completely.

Another useful quantity that we can compare with Mathis's numerical work is the fraction of Lyman-continuum photons absorbed by the gas. The analytic ex-




FIG. 1.— $\text{Log}_{10}$  (neutral fraction) plotted against radial distance in units of  $r_g$ . Each curve is labeled by the corresponding value of  $\tau_d$ . For comparison, the numerical computation by Mathis (1971) is shown for the case  $\tau_d = 0.4$  (*dashed line*).




FIG. 2.—Fraction of ionizing photons absorbed by gas plotted against optical depth for true absorption by dust within the ionized region. *Crosses*, the numerical results of Mathis (1971).

pression for this fraction as a function of  $y$  is found to be  $y^3/(1 - e^{-\tau})$ . At the boundary of the H II region ( $y \equiv y_0$ )  $(1 - x)\tau_s \gg 1$ , and  $\tau \rightarrow \infty$ . Consequently, we obtain the extent of the ionized region by setting the left-hand side of equation (7) equal to zero. The fraction  $f$  of stellar ionizing photons absorbed by gas in the entire ionized region is therefore given by

$$f = y_0^3 = \frac{1}{8} \tau_0^3 \exp(-\tau_0) / \{\tau_0^2 - 2\tau_0 + 2[1 - \exp(-\tau_0)]\}, \quad (8)$$

where  $\tau_0 = \tau_d y_0$  is the optical depth for absorption by dust within the ionized region.

In figure 2 we plot the fraction  $f$  versus  $\tau_0$ . For comparison with our analytic curves, we show results computed numerically by Mathis. As is evident, our solution is in perfect agreement with the computations of Mathis for  $\tau_0 \leq 1.2$ . For larger amounts of dust absorption, our solution tends to overestimate the fraction of ionizing radiation absorbed by the gas.

Part of this discrepancy is due to our neglect of helium. Mathis (1971) shows that the He<sup>+</sup> and H<sup>+</sup> ionization zones tend to coincide in the presence of dust, because the H<sup>+</sup> ionization zone moves inward with increasing  $\tau_d$  as H-ionizing photons are absorbed by dust. One of us has also given an analytic expression for this effect (Petrosian 1972). The principal effect of He on the hydrogen ionization structure is to act as a converter of relatively hard He-ionizing photons to softer photons (each He recombination liberating  $\sim 0.9$  H-ionizing photons), for which the cross-section for ionizing H is much higher (Hummer and Seaton 1964). Consequently, the ionization of H is increased, which implies that at  $\tau_d \gtrsim 1$  our neglect of He leads us to overestimate the neutral fraction, and therefore the fraction of ionizing photons absorbed by the gas.

Part of the difference between the analytic and numerical solutions appears to be due also to our neglect of scattering by the dust, the effect of which is to provide a higher probability for absorption of photons by either gas or dust grains. We could account for the effect of scattering by redefining all our optical depths as

$$\tau(r) = \int_0^r \sigma n \frac{ds}{dr} dr,$$

where  $ds$  is an element of length measured along the actual scattering path of the photon. With this new definition of optical depth our results remain unchanged, but the points of figure 2 from Mathis's numerical work move to the left (toward higher optical depths) approximately by a factor

$$\int_0^{\tau_0} \frac{ds}{dr} \frac{dr}{r_0},$$

where  $r_0 = r_s y_0$  is the Strömgen radius of the dusty nebula. This factor depends on the scattering optical depth  $\tau_{\text{scat}} = \tau_d \tilde{\omega} / (1 - \tilde{\omega})$ , and increases with increasing values of the dust albedo,  $\tilde{\omega}$ . The exact dependence of this factor on the albedo is complicated and beyond the scope of the present paper.

In order to apply figure 2 to observations of far-infrared emission from H II regions, we note that the expected infrared luminosity is

$$L(\text{IR}) = L(\text{La}) + (1 - f) \langle h\nu \rangle_{\text{Ly}\alpha} S + L_{\nu < \text{Ly}\alpha} [1 - \exp(-\tau'_0)], \quad (9)$$

where  $\langle h\nu \rangle_{\text{Ly}\alpha}$  is the average energy of stellar Lyman-continuum photons,  $L_{\nu < \text{Ly}\alpha}$  is the stellar luminosity below the Lyman continuum, and  $\tau'_0$  is the effective absorption optical depth of dust in the H II region and in the surrounding H I region for photons with  $h\nu < 13.6$  eV. For a nebula which is optically thick to La photons (case B), the value of the La luminosity of the nebula (assuming that the La can eventually escape from the nebula), is given by  $L(\text{La}) = \frac{2}{3} h\nu_{\text{La}} S f$ , and can be inferred either from the observations of Balmer photons suitably corrected for reddening or from the radio-continuum luminosity. If the spectrum of the ionizing stars and the observed infrared flux of H II regions are known, we can use equation (9) to determine the fraction  $f$ , and from figure (2) the effective absorption optical depth of the dust. This procedure may allow one to empirically determine the far-ultraviolet absorption properties of the dust. For example, in the case of the Orion Nebula, the total extinction by dust near H $\beta$  has been found to be 1.8 mag (Mathis 1970); this value is consistent with other independent determinations at optical wavelengths (cf. Münch and Persson 1971). On the other hand, we estimate that, in our earlier notation  $L(\text{IR})/L(\text{La}) = 7.5$ . Hence if we neglect the last term in equation (8), as might be the case for a hot central star or if  $\tau'_0 \ll \tau_0$ , and set  $\langle h\nu \rangle_{\text{Ly}\alpha} = 15$  eV, we find that  $f \approx 0.26$ . We then deduce from figure 2 that the true absorption by dust of Lyman-continuum photons is about 1.7 mag. It follows from these numbers that, for a ratio of total absorption cross-section in the Lyman continuum to that at H $\beta$  of between 2:1 and 3:1, the albedo at H $\beta$  must be about  $0.6 \pm 0.1$ . In principle, one could now infer further physical characteristics of the dust from this result; however, the paucity of available data on optical constants at ultraviolet wavelengths does not appear to allow any definitive conclusion to be drawn at present. In the event that the last term in equation (8) is not negligible, similar calculations can be carried out if the spectrum of the central star is known.

In summary, our analytic approximation (6) to the ionization structure of H II regions with a uniform distribution of dust should be useful in aiding future studies of line and continuum emission from H II regions. In particular, we have obtained a simple expression for the fraction of ionizing photons absorbed by the dust (eq. [8]), a quantity which is consistent with the infrared observations provided that the dust absorption optical depths are typically between 1.0 and 2.0 mag. This approach offers an independent way of obtaining the total absorption optical depth to internal dust in H II regions, which can then be compared with the optical depth at H $\beta$ . If the stellar spectrum is known, this approach can be used for determination of the albedo of the dust.

We are grateful to Dr. W. Mathews for interesting discussions. The work of V. P. has been supported by the National Aeronautics and Space Administration grant NGR 05-020-510; the work of J. S. by National Aeronautics and Space Administration grant NGR 05-003-453; and the work of G. F. by National Science Foundation grant GP 18476.

## REFERENCES

- Harper, D. A., and Low, F. J. 1971, *Ap. J. (Letters)*, **165**, L9.  
Hummer, D. G., and Seaton, M. J. 1964, *M.N.R.A.S.*, **127**, 217.  
Mathis, J. S. 1970, *Ap. J.*, **159**, 263.  
———. 1971. *ibid.*, **167**, 261.  
Münch, G., and Persson, S. E. 1971, *Ap. J.*, **165**, 241.  
Petrosian, Vahé. 1972, paper presented at IAU Symposium No. 52, Albany, New York.