

THE SOFT GAMMA-RAY SPECTRUM OF A0535+26: DETECTION OF AN ABSORPTION FEATURE AT 110 keV BY OSSE

J. E. GROVE,¹ M. S. STRICKMAN, W. N. JOHNSON, J. D. KURFESS, AND R. L. KINZER
 E. O. Hulburt Center for Space Research, Code 7650, Naval Research Laboratory, Washington, DC 20375

C. H. STARR AND G. V. JUNG
 Universities Space Research Association, Washington, DC 20024

AND

E. KENDZIORRA, P. KRETSCHMAR, M. MAISACK, AND R. STAUBERT
 Astronomisches Institut der Universität Tübingen, D-72076 Tübingen, Germany

Received 1994 August 11; accepted 1994 October 11

ABSTRACT

We present soft γ -ray observations by the Oriented Scintillation Spectrometer Experiment (OSSE) on the *Compton Gamma Ray Observatory* (GRO) of the transient X-ray binary pulsar A0535+26. The observations were made 1994 February 8–17, immediately prior to the peak of a giant outburst. The phase averaged spectrum is complex and cannot be described by a single-component model. We find that structure in the spectrum above 100 keV can best be modeled by an absorption feature near 110 keV, which we interpret as the signature of cyclotron resonant scattering. Because of OSSE's 45 keV threshold, we are unable to make a definitive statement on the presence of a 55 keV absorption line; however, we can conclude that if this line does exist, it must have a smaller optical depth than the line at 110 keV. A first harmonic (=fundamental) cyclotron resonance at 110 keV corresponds to a magnetic field strength at the surface of the neutron star of $\sim 1 \times 10^{13}$ G ($\sim 5 \times 10^{12}$ G if the first harmonic is at 55 keV).

Subject headings: gamma rays: observation — line identification — stars: individual (A0535+26) — stars: magnetic fields — stars: neutron — X-rays: stars

1. INTRODUCTION

The massive X-ray binary A0535+26 is a recurrent transient pulsar, with pulse period ≈ 104 s (Rosenberg et al. 1975) and recurrence period ≈ 111 days (Motch et al. 1991). The optical companion is the Be star HDE 245770 (Li et al. 1979) at an estimated distance of 1.8 ± 0.3 kpc (Giangrande et al. 1980). For a review, see Giovannelli & Graziati (1992). The X-ray outbursts are thought to be powered by enhanced mass transfer from the companion star near periastron. The intensity of the X-ray emission can vary dramatically from one periastron passage to the next: No outburst may be observed; a “typical” outburst may occur, with peak flux of a few hundred mCrab units (1–10 keV); or a “giant” outburst may occur, with peak X-ray flux of several Crab units. Giant outbursts may appear away from periastron (Motch et al. 1991).

Previous hard X-ray spectra of A0535+26 have been modeled variously as exponential ($kT = 17$ keV; Hameury et al. 1983), blackbody ($kT = 9$ keV; Frontera et al. 1985), and Wien ($kT = 8$ keV; Dal Fiume et al. 1988). Kendziorra et al. (1992, 1994) reported the detection of two absorption features, near 50 and 100 keV, in the spectrum of A0535+26 in outburst. Previous estimates of the magnetic field strength have ranged over $4\text{--}20 \times 10^{12}$ G, based on the suggestion of cyclotron line features (Kendziorra et al. 1992, 1994), the energy dependence of the pulsed fraction (Dal Fiume et al. 1988), or the multiplicity of pulse shapes at low energies (Kanno 1980).

We report here on spectral observations of A0535+26 by the Oriented Scintillation Spectrometer Experiment (OSSE) on

the *Compton GRO* over the interval 1994 February 8–17, during the rise immediately preceding the peak of a giant outburst. These observations were scheduled as a “target of opportunity” triggered by the detection of the outburst by the BATSE instrument on *GRO*. The rise to maximum of this giant outburst spanned a binary phase interval of ~ 0.15 , with the peak intensity occurring on 1994 February 18 at binary phase 0.95 ± 0.03 , according to the ephemeris of Motch et al. (1991). At the maximum, the pulsed emission reached 8 Crab (nebula + pulsar) flux units in the 20–40 keV band (Finger, Wilson, & Hagedorn 1994; Wilson et al. 1994).

2. OBSERVATIONS

The OSSE instrument consists of four identical large-area NaI(Tl)–CsI(Na) phoswich detector systems, as described in detail in Johnson et al. (1993). Full-resolution spectra were accumulated every 4.096 s in a sequence of 2 minute measurements of the source field alternated with 2 minute, offset-pointed measurements of background. The observation was made in twice-normal gain to improve the low-energy response. A total of 1.8×10^5 s of highest-quality data, equally divided between source and background, were collected for each of the four detectors.

For this observation, the source and background fields were placed to ensure that the Crab Nebula, a potentially contaminating source, did not cause a significant contribution to the flux measurement of A0535+26. Below 100 keV, the net (source – background) Crab contribution is less than 0.1% of the measured flux from A0535+26, and in the critical range from 100–200 keV where we see spectral structure, the contribution is less than 10^{-4} photons $\text{cm}^{-2} \text{s}^{-1} \text{MeV}^{-1}$, or $\lesssim 1\%$ of the measured A0535+26 flux.

¹ E-mail: grove@osse.nrl.navy.mil

3. RESULTS

We detected emission from A0535+26 with high statistical significance from 45 keV to ~ 200 keV. The average 45–200 keV luminosity during 1994 February 8–17 was $4 \times 10^{36}(d/1.8 \text{ kpc})^2 \text{ ergs s}^{-1}$; the luminosity doubled over the nine days of the observation, from 2.5×10^{36} to $5.1 \times 10^{36}(d/1.8 \text{ kpc})^2 \text{ ergs s}^{-1}$.

We fitted the phase-averaged spectrum between 45 and 300 keV from the sum of the four detectors for the entire observation using a forward-folding method (Johnson et al. 1993). We derived error estimates in the parameters of the various spectral models by χ^2 -mapping, with confidence intervals specified according to Lampton, Margon, & Bowyer (1976). With such a strong source, it is important to address possible sources of systematic error. We have identified a number of likely contributors, and in all the analysis described herein we have added our best estimate of the systematic error to the statistical error. The systematic error becomes the dominant source of error below ≈ 70 keV, rising to $\sim 3\%$ of the flux at 45 keV. It does not significantly alter the values of the best-fit parameters.

Figure 1a shows the phase-averaged spectrum summed over the entire observation. The spectrum is quite soft (Wien temperature ≈ 10 keV), although no single-component model adequately describes the spectrum. Simple continuum models show very significant residuals and have unacceptably large χ^2 values (Table 1). The structure apparent above 100 keV could indicate the emergence of weak, underlying, high-energy continuum emission, or it could indicate a broad emission or absorption line. To determine which of these is most likely, we tested a number of models in two classes: a class of two-component continuum spectra, and a class of single-component continuum spectra together with line profiles for emission or absorption (Table 1). In both classes, the continuum model that accounts for the bulk of the emission is a power law times an exponential (hereafter, “PLExp”), chosen because of its general adequacy in describing the standard and

TABLE 1
MINIMUM χ^2_{dof} VALUES FOR CANDIDATE SPECTRAL MODELS

Description	χ^2_{dof} (dof)
Single-Continuum Models	
OTTB ^a	59.4 (94)
Wien	11.7 (94)
PLExp ^b	5.44 (93)
Two-Continuum Models	
PLExp + Power Law	5.42 (91)
PLExp + PLExp	5.30 (90)
Continuum + Emission-Line Models	
Wien + Gaussian	11.9 (91)
PLExp + Gaussian	4.72 (90)
Continuum with Absorption Models	
Wien \times One Line ^c	1.86 (91)
Wien \times Two Lines	1.13 (90)
PLExp \times One Line ^c	1.11 (90)
PLExp \times Two Lines	1.06 (89)

^a Optically thin thermal bremsstrahlung.

^b Power-law times exponential.

^c $E_L \approx 110$ keV.




FIG. 1.—(a) Phase-averaged spectrum of A0535+26 from 1994 February 8–17. The outburst peaked 1994 February 18 (Wilson et al. 1994). Solid line is the model fit, consisting of a power law times an exponential continuum (eq. [2]) with transmission function of a single absorption line near 110 keV (eq. [3]). Inset shows joint 68% ($\chi^2_{\text{min}} + 5.9$) and 95% ($\chi^2_{\text{min}} + 11.1$) confidence intervals for the line centroid and optical depth from the single-line model (b) Residuals from best fit. (c) Transmission function for the best-fit absorption line near 110 keV. Data are divided by the continuum portion (eq. [2]) of the best-fit model.

hard X-ray spectra of X-ray binaries (White, Swank, & Holt 1983).

The models from the first class, the two-component continuum models, do not give acceptable fits (Table 1). The weak, hard continuum necessary to describe the emission above 100 keV—either a simple power law or PLExp—has a substantial effect on the spectrum below 100 keV and generates significant residuals. Models from the second class employing a Gaussian emission feature (centroid energy ≈ 150 keV) are similarly poor, with χ^2 probabilities of less than 10^{-7} .

The remaining models from the second class, those with broad absorption lines, provide substantially better fits. In these models, the continuum is modified by a multiplicative transmission function from one or more absorption lines, which we presume arise from cyclotron resonant scattering. First proposed by Tanaka (1986), the transmission function assumes a Lorentzian times E^2 form, which resembles the cyclotron scattering cross section (Herold 1979; Daugherty & Harding 1986; Harding & Daugherty 1991) and has been successfully used to model cyclotron absorption features in *Ginga* data (e.g., Nagase et al. 1991; Tamura et al. 1992). The model

function has the form

$$f(E) = AC(E) \exp[-L(E)], \quad (1)$$

where

$$C(E) = E^{-\alpha} \exp(-E/kT), \quad (2)$$

$$L(E) = \sum_{n=1}^2 \frac{\tau_n(nW)^2(E/nE_L)^2}{(E - nE_L)^2 + (nW)^2}. \quad (3)$$

The continuum term, $C(E)$, is the PLEXP model with folding energy kT . In the absorption profile $L(E)$, parameters τ_n , nE_L , and nW are the optical depth, energy, and width (the half-width at half-maximum of the Lorentzian, HWHM) of absorption line $n = 1$ or 2. We have assumed a harmonic relationship of 1:2 between the line energies and widths. While in the relativistic treatment the resonant energies are not strict multiples of the cyclotron energy (Herold 1979), the nonlinearities are small (of order several percent for the first few levels) and have been ignored. Trial fits with independent line energies and widths were consistent with the harmonic assumption and did not substantially improve the fit.

Table 1 shows that the best fit is achieved with the general PLEXP continuum model with two absorption lines, although the first harmonic (=fundamental) near 55 keV has a small optical depth ($\tau_1 = 0.1_{-0.1}^{+0.5}$; 68% confidence, $\chi^2_{\min} + 7.0$), which serves mostly to flatten the model spectrum at low energies. Indeed this absorption line is not strictly required with the PLEXP continuum model: χ^2 probabilities for models with free or zero optical depth at 55 keV are both acceptable (33% and 22%, respectively). The 95% confidence upper limit is $\tau < 1.1$ ($\chi^2_{\min} + 12.6$). We note also that the Wien continuum with two absorption lines is acceptable; however, the lines are extremely broad (42% $^{+4}_{-3}$ % HWHM), and the 55 keV line is difficult to distinguish from a smooth change in the continuum.

Parameters of the best-fit model are given in Table 2. The errors given correspond to 68% confidence intervals for six interesting parameters (we judge the intensity A to be irrelevant); thus we use $\chi^2_{\min} + 7.0$. Care should be taken in interpreting the error estimates for this nonlinear model; for example, the 95% (i.e., "2 σ ") confidence interval is not given by twice the 68% (i.e., "1 σ ") confidence interval. The best-fit

optical depth of the line at 110 $_{-4}^{+2}$ keV is 1.8 $_{-0.5}^{+1.1}$, implying $\sim 15\%$ transmission at the centroid. The width of the line (28 $_{-9}^{+16}$ keV HWHM) is typical of the absorption features detected by *Ginga* ($\sim 25\%$; e.g., Tamura et al. 1992). The width and depth of the line combine to affect the overall spectrum significantly throughout the 45–200 keV band (Fig. 1c). Because of the large uncertainties in the data > 150 keV, the existence of possible higher harmonics near 165 and 220 keV is not well constrained; e.g., the 95% confidence upper limit on the optical depth at 165 keV is $\tau < 4$ ($\chi^2_{\min} + 14.1$). Figure 1a shows a PLEXP continuum model with a single absorption feature (near 110 keV); the optical depth of the potential first harmonic near 55 keV has been set to zero. With only a single absorption line, significant correlation among the parameters is removed, and the confidence intervals (inset Fig. 1a) are significantly smaller than those in Table 2. The residuals (Fig. 1b) demonstrate that a feature near 55 keV is not required, reinforcing the conclusion drawn from the χ^2 probabilities. The total excess emission above the model in the 200–600 keV band is less than 2 σ .

We have also employed a more general line profile than that of equations (1) and (3), a multiplicative Gaussian. Fits with this symmetric profile are essentially identical in quality to those given above (e.g., for the two-line model $\chi^2_{\min} = 1.09$ with 89 degrees of freedom) and lead to identical conclusions, in particular that only the line near 110 keV is strongly required. The line centroid is shifted to a slightly higher energy 115 $_{-4}^{+3}$ keV, presumably because the Gaussian lacks the blue tail of the exponentiated Lorentzian times E^2 . The line width is unchanged 26 $_{-7}^{+69}$ keV HWHM, although it is not well constrained. The line width is governed by thermal broadening, as well as the averaging over magnetic field conditions and viewing angle present in the phase-averaged spectrum.

Fits to the spectrum for each day show that the line centroid and width vary by less than 2 σ , which suggests that we have not introduced a significant bias by summing the data over the entire observation. However, as the luminosity increases during the outburst, the spectrum does soften. In order to characterize the spectral change in a simple manner, we fit the daily spectra over 45–80 keV with a Wien law. Although formally these fits are only marginally acceptable (χ^2 probabilities typically $\sim 1\%$), they are sufficient to convey that the magnitude of the softening is modest. The Wien temperature decreases by 7% as the luminosity doubles.

4. DISCUSSION

We have shown that an absorption feature near 110 keV is present in the phase-averaged spectrum of A0535+26 during the giant outburst of 1994 February. This is the first time that both the low-energy and high-energy wings of an absorption line have been clearly detected at such high energies in any binary pulsar. We interpret this absorption as a signature of cyclotron resonance scattering. Two-continuum and continuum + emission-line models are strongly rejected by the χ^2 test (Table 1).

We are unable to make a definitive statement on the presence of a 55 keV absorption line in the phase-averaged spectrum. Furthermore, there have been no reports to date on the existence of such a feature in the BATSE data. Adequate fits are obtained with and without this feature, with a slight preference for a weak line in combination with the general power-law times exponential continuum (Table 1). However, we caution that it is always difficult to distinguish between the

TABLE 2

BEST-FIT PARAMETERS FOR POWER-LAW TIMES EXPONENTIAL CONTINUUM AND TWO ABSORPTION LINES

Parameter	Value
Continuum Parameters	
Intensity ($\text{cm}^{-2} \text{s}^{-1} \text{MeV}^{-1}$) ^a	2.35 $_{-0.35}^{+4.32}$ ^b
Power-law index, α ^c	-0.13 $_{-1.67}^{+0.65}$
Folding energy, kT (keV)	17.8 $_{-3.8}^{+5.4}$
Line Parameters	
Line energy, first harmonic, E_L (keV)	55.0 $_{-1.9}^{+1.1}$
Line HWHM, first harmonic, W (keV)	14.2 $_{-4.5}^{+8.1}$
Optical depth, first harmonic, τ_1	0.10 $_{-0.10}^{+0.48}$
Optical depth, second harmonic, τ_2	1.8 $_{-0.5}^{+1.1}$

^a At 70 keV.

^b Errors correspond to 68% confidence, assuming six interesting parameters ($\chi^2_{\min} + 7.0$; Lampton, Margon, & Bowyer 1976).

^c I.e., photon spectral index.

appearance of a weak, broad line feature and an inadequate description of the underlying continuum: There is significant coupling between the line and continuum model parameters. Such is the case here, and it is compounded by uncertainties in the instrument response and the absence of adequate spectral coverage below the nominal line energy. We can conclude, however, that if a 55 keV feature is present, its optical depth must be significantly smaller than that of the 110 keV feature. It is surprising that the ratio of depths would be small ($\sim 1:20$ in the best-fit model with both depths free; $< 1:3.5$ at 95% confidence). Radiative transfer calculations based on a neutron-star paradigm for the hard-spectrum γ -ray bursts do show significant filling-in of the first harmonic by two- γ decay of the second excited state (e.g., Wang et al. 1989; Alexander & Mészáros 1991). However, Bulik et al. (1992) point out that this "photon spawning" process should be much less important in accreting X-ray pulsars, where the typically soft, thermal spectrum would not populate the higher harmonics significantly, and therefore would not lead to any appreciable down-scattering into the first harmonic.

Kendziorra et al. (1992, 1994) reported the detection of two absorption features (≈ 50 and ≈ 100 keV) in the spectrum of A0535+26 at periastron. An exponentially truncated power-law model for the underlying continuum was required by their combined TTM and HEXE spectrum; a Wien law was unacceptable. The 50 keV absorption was detected at low signifi-

cance ($\lesssim 3\sigma$) in the phase-averaged spectrum, and HEXE lacked the sensitivity to detect with certainty the high-energy wing of the 100 keV line; however, most importantly, the amplitudes of the absorption features depended on rotation phase. We have begun analysis of the phase-resolved spectroscopy data from OSSE, which will be sensitive to a phase-dependent absorption at 55 keV, if it exists. We note that the 95% confidence limit we set here on the optical depth at 55 keV ($\tau < 1.1$) in the phase-averaged spectrum does not exclude the HEXE value.

If the feature at 110 keV does indeed result from cyclotron resonant scattering in a strongly magnetized plasma, it can be used to estimate the magnetic field strength. A first harmonic resonance at 110 keV corresponds to a field strength at the surface of the neutron star of $9.5 \times 10^{12}(1+z)$ G, where z is the gravitational redshift. If instead the first harmonic is at 55 keV, the field is $4.7 \times 10^{12}(1+z)$ G. In either case, this is the highest field strength for an accreting neutron star ever measured through direct detection of cyclotron absorption, although the value is within reason and is consistent with the strongest magnetic fields inferred from the spin-down of radio pulsars.

This work was supported by NASA contract S-10987-C and by DARA grant 50 OR 92054.

REFERENCES

- Alexander, S. G., & Mészáros, P. 1991, *ApJ*, 344, L1
 Bulik, T., et al. 1992, *ApJ*, 395, 564
 Dal Fiume, D., et al. 1988, *ApJ*, 331, 313
 Daugherty, J. K., & Harding, A. K. 1986, *ApJ*, 309, 362
 Finger, M. H., Wilson, R. B., & Hagedorn, K. S. 1994, *IAU Circ.*, No. 5931
 Frontera, F., et al. 1985, *ApJ*, 298, 585
 Giangrande, A., et al. 1980, *A&AS*, 40, 289
 Giovanelli, F., & Graziati, L. S. 1992, *Space Sci. Rev.*, 59, 1
 Hameury, J. M., et al. 1983, *ApJ*, 270, 144
 Harding, A. K., & Daugherty, J. K. 1991, *ApJ*, 374, 687
 Herold, H. 1979, *Phys. Rev. D*, 19, 2868
 Johnson, W. N., et al. 1993, *ApJS*, 86, 693
 Kanno, S. 1980, *PASJ*, 32, 105
 Kendziorra, E., et al. 1992, in *Proc. 28th Yamada Conf., Frontiers of X-Ray Astronomy*, ed. Y. Tanaka & K. Koyama (Tokyo: Univ. Acad.), 51
 Kendziorra, E., et al. 1994, *A&A*, in press
 Lampton, M., Margon, B., & Bowyer, S. 1976, *ApJ*, 208, 177
 Li, F., et al. 1979, *ApJ*, 228, 893
 Motch, C., et al. 1991, *ApJ*, 369, 490
 Nagase, F., et al. 1991, *ApJ*, 375, L49
 Rosenberg, F. D., et al. 1975, *Nature*, 256, 628
 Tamura, K., et al. 1992, *ApJ*, 389, 676
 Tanaka, Y. 1986, in *IAU Colloq. 89, Radiation Hydrodynamics in Stars and Compact Objects*, ed. D. Mihalas & K.-H. A. Winkler (Berlin: Springer), 198
 Wang, J. C. L., et al. 1989, *Phys. Rev. Lett.*, 63, 1550
 White, N. E., Swank, J. H., & Holt, S. S. 1983, *ApJ*, 270, 711
 Wilson, R. B., et al. 1994, *IAU Circ.*, No. 5945