

H₂ EMISSION FROM THE INNER 400 PARSECS OF THE GALAXY

SOOJONG PAK, D. T. JAFFE, AND L. D. KELLER

Department of Astronomy, University of Texas at Austin, Austin, TX 78712; soojong, dtj, keller@astro.as.utexas.edu

Received 1995 July 11; accepted 1995 November 8

ABSTRACT

We have mapped the H₂ $v = 1 \rightarrow 0$ $S(1)$ ($\lambda = 2.1215 \mu\text{m}$) emission line along a 400 pc-long strip and in a 50 pc region in the Galactic center. There is H₂ emission throughout the surveyed region. The typical dereddened ($A_K = 2.5$ mag) H₂ $v = 1 \rightarrow 0$ $S(1)$ surface brightness, $\sim 3 \times 10^{-5}$ ergs s⁻¹ cm⁻² sr⁻¹, is similar to the surface brightness in large-scale photon-dominated regions in the Galactic disk. We investigate two possible excitation mechanisms for the H₂ emission, UV excitation by photons from OB stars and shock waves, and conclude that UV excitation is more likely. The total H₂ $v = 1 \rightarrow 0$ $S(1)$ luminosity in the inner 400 pc region of the Galaxy is 8000 L_\odot . The ratio of the H₂ to far-IR luminosity in the inner 400 pc of the Galaxy agrees with that in starburst galaxies and ultraluminous IR-bright galaxies.

Subject headings: galaxies: ISM — infrared: ISM: lines and bands — ISM: molecules

1. INTRODUCTION

Physical conditions in the interstellar medium of the Galactic center¹ are significantly different from those in the solar neighborhood. The thin disk of interstellar material in the Galactic center (size: 450 pc \times 40 pc) contains $\sim 10^8 M_\odot$ of dense molecular gas, $\sim 10\%$ of the Galaxy's molecular mass (Güsten 1989). The molecular clouds in the Galactic center have higher density, metallicity, and internal velocity dispersion than the clouds in the solar neighborhood (Blitz et al. 1993). Strong radio continuum radiation from giant H II regions and extended low-density (ELD) H II (Sofue 1985), as well as far-IR radiation from dust (Odenwald & Fazio 1984), indicate that the UV radiation field is intense. The energetic conditions in the Galactic center can provide a unique view of the interaction between stellar UV radiation and molecular clouds and a nearby example for the nuclei of other galaxies.

Ro-vibrational lines of H₂ trace photon-dominated regions (PDRs), where far-UV photons excite the H₂, and shocked regions, where the H₂ is thermally excited. As a result, the central regions (~ 1 kpc) in starburst galaxies are powerful emitters of near-IR H₂ emission (Puxley, Hawarden, & Mountain 1989; Joseph 1989; Lester et al. 1990; Moorwood & Oliva 1990). Vigorous star formation in these galaxies produces large numbers of UV photons, which can excite H₂, while subsequent supernovas can shock-excite the H₂.

Gatley et al. (1984, 1986) and Gatley & Merrill (1993) have observed H₂ emission from the inner 5 pc diameter (2') in the Galactic center, a much smaller region than those observed in starburst galaxies. With the University of Texas Near-Infrared Fabry-Perot Spectrometer (Luhman et al. 1995), it is now possible to observe H₂ emission over much larger angular scales. We describe here a program to map the Galactic center in H₂ emission on a scale of several degrees (several hundred parsecs) and discuss the likely H₂ excitation mechanism. We can then compare the central region of our Galaxy with those in other galaxies.

¹ We use here the term "Galactic center" to denote the inner several-hundred-parsec region of our Galaxy. We adopt a distance of 8.5 kpc, at which 1° corresponds to 148 pc.

2. OBSERVATIONS

We observed the H₂ $v = 1 \rightarrow 0$ $S(1)$ ($\lambda = 2.1215 \mu\text{m}$) line at the McDonald Observatory 0.91 m telescope in 1994 May and June, using the University of Texas Near-Infrared Fabry-Perot Spectrometer (Luhman et al. 1995). To select a single order from the Fabry-Perot interferometer, we used a 1% interference filter cooled to 77 K. The telescope (f-ratio 13.5), a collimator (effective focal length 343 mm), and a field lens (effective focal length 20 mm) combined to produce a beam diameter of 3'3 (equivalent disk).

The Fabry-Perot interferometer operated in 92d order with an effective finesse of 17.7, yielding a spectral resolution (FWHM) of 184 km s⁻¹. The scanning spectral range was ± 335 km s⁻¹ centered at $V_{\text{LSR}} = 0$, with 20 sequentially exposed channels. We nodded the telescope between the object and the sky every ~ 60 s to subtract background and the telluric OH line emission. The sky positions were offset by $\pm 1^\circ$ in declination ($\Delta l = \pm 0:85$, $\Delta b = \pm 0:53$) from the object positions. The telescope pointing error was $\pm 15''$. We made a strip map along the Galactic plane running across Sgr A*, at $b = -0:05$, from $l = -1:2$ (178 pc) to $+1:6$ (237 pc), taking spectra at 0:1 or 0:2 intervals (Fig. 1). We also mapped the central 50 pc region including Sgr A and the radio "arc" (the arched filaments and the vertical filaments, Yusef-Zadeh, Morris, & Chance 1984) on a 0:05 grid (see Fig. 2). The relative flux calibration is accurate to $\pm 15\%$.

3. RESULTS

There is H₂ emission throughout the 400 pc diameter region around the Galactic center. About 70% of the observed positions along the strip at $l = -0:05$ have detections of the H₂ $v = 1 \rightarrow 0$ $S(1)$ line with a significance of 3 σ or more. Figure 1 shows the intensity distribution of H₂ $v = 1 \rightarrow 0$ $S(1)$ along the strip. The H₂ intensity peaks strongly at Sgr A and decreases continuously from $l = -0:1$ to $l = -0:7$. The "dust ridge" seen in 800 μm continuum emission (Lis & Carlstrom 1994) may cause the dip between $l = +0:1$ and $l = +0:3$. Away from the nucleus, the intensity distribution is fairly flat.

Figure 2 shows the H₂ $v = 1 \rightarrow 0$ $S(1)$ intensity distribution in the inner 50 pc of the Galaxy overlaid on a simplified radio continuum map (based on Yusef-Zadeh et al. 1984). We have




FIG. 1.—Observed intensity distribution of $\text{H}_2 \nu = 1 \rightarrow 0 S(1)$ ($\lambda = 2.1215 \mu\text{m}$) along the Galactic plane at $b = -0^\circ 05$. The intensity values have not been corrected for interstellar extinction. The error bars represent 1σ measurement uncertainties. The strongest emission, at $l = -0^\circ 05$, is from Sgr A. The vertical lines show the positions of the prominent radio continuum sources (Altenhoff et al. 1978).

detected H_2 emission in the arched filaments and in the Sickle at levels comparable to those of the H_2 emission elsewhere along the plane. The inset in Figure 2 shows the $\nu = 1 \rightarrow 0 S(1)$ distribution in the $0^\circ 022 \times 0^\circ 038$ region around Sgr A West mapped by Gatley et al. (1986). In Gatley's work, the H_2 appears to be brightest along the inner edge of the circumnuclear gas ring at radius of 1.0–2.5 pc. Our measured flux at ($l = -0^\circ 05$, $b = -0^\circ 05$), 1.5×10^{-11} ergs $\text{s}^{-1} \text{cm}^{-2}$, agrees within the errors with the total flux from the map of Gatley et al. (1986; $F = 2.0 \times 10^{-11}$ ergs $\text{s}^{-1} \text{cm}^{-2}$). The H_2 emission observed adjacent to ($l = -0^\circ 05$, $b = -0^\circ 05$) most likely




FIG. 2.—Positions (circles with a diameter of the telescope beam size, $3'.3$) and measured H_2 intensities (numbers inside the circles in units of 10^{-6} ergs $\text{s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$) observed in $\text{H}_2 \nu = 1 \rightarrow 0 S(1)$ in the inner 50 pc of the Galaxy. Typical measurement uncertainties are 0.7×10^{-6} ergs $\text{s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$. The heavy lines show a schematic version of the radio continuum distribution (Yusef-Zadeh et al. 1984). The small box and inset show the $\text{H}_2 \nu = 1 \rightarrow 0 S(1)$ distribution in the circumnuclear gas ring (Gatley et al. 1986). The plus sign marks the position of Sgr A* ($l = -0^\circ 0558$, $b = -0^\circ 0462$).

arises from portions of the circumnuclear ring beyond their map.

4. DISCUSSION

4.1. Extinction Correction

At $2.2 \mu\text{m}$, the emission from the Galactic center is attenuated by interstellar material in the foreground (“foreground extinction,” mostly 4–8 kpc from the Galactic center) and by material in the Galactic center itself (“Galactic center extinction”). Catchpole, Whitelock, & Glass (1990) mapped the extinction toward the Galactic center by observing the near-IR reddening of giant stars in the central few hundred parsecs. Along our H_2 strip at $b = -0^\circ 05$ (for $-0^\circ 6 < l < +0^\circ 6$), the extinction is fairly uniform, with a value $A_K \sim 2.5$ mag. Although most of this extinction is in the foreground, some of it could occur within the Galactic center since Catchpole et al. (1990) were able to identify patches in their maps with $A_K > 2.5$ mag with known molecular clouds in the Galactic center (see Plate 4 in Glass, Catchpole, & Whitelock 1987). Based on this work, we adopt $A_K = 2.5$ mag for the foreground extinction.

The Galactic center extinction greatly exceeds the foreground extinction. Typical $^{12}\text{CO } J = 1 \rightarrow 0$ line strengths along the strip we have surveyed in H_2 are 1500 K km s^{-1} (Oort 1977). This line strength implies an A_K of 10–40 mag, depending on the CO/H_2 and A_K/H_2 ratios in the Galactic center (Sodroski et al. 1994). The extinction through individual clouds may also be substantial ($A_K \sim 10$ –30 for a 10 pc-long cloud with $n_{\text{H}_2} = 10^4 \text{ cm}^{-3}$). The relevance of the Galactic center extinction depends on the source of the H_2 emission. Any H_2 emission that originates within the clouds will be highly extinguished. If the H_2 emission arises on the cloud surfaces, however, we only miss the H_2 flux from the back side of each cloud.

Clouds that lie in front of other clouds will further reduce the flux that reaches us from the front surfaces. If, in the Galactic center, the velocity-integrated area-filling factor of clouds, f , is substantially greater than unity, extinction by overlapping clouds will reduce the H_2 flux observed from the front surfaces by a factor of $\sim 1/f$ in addition to the foreground extinction and to the loss of emission on the opposite sides of the clouds. Typical clouds in the Galactic center disk have kinetic temperatures of $\sim 70 \text{ K}$ and line widths of $\sim 20 \text{ km s}^{-1}$ (Güsten 1989). An ensemble of such clouds could produce the observed $^{12}\text{CO } J = 1 \rightarrow 0$ lines in the Galactic center with $f \sim 1$. We therefore conclude that the extinction of any H_2 emission from cloud surfaces facing the Sun beyond the foreground extinction of $A_K = 2.5$ mag discussed above is not substantial. Since the extinction of emission from within the clouds or from the sides facing away from us is difficult to estimate and since no correction is usually made for such effects in giant molecular clouds and galactic nuclei, we make no additional extinction corrections here.

4.2. UV Excitation of H_2

If one ignores the region immediately around Sgr A*, the dereddened ($A_K = 2.5$ mag; see § 4.1) $\text{H}_2 \nu = 1 \rightarrow 0 S(1)$ surface brightness along the Galactic plane has a roughly constant value of $\sim 3 \times 10^{-5}$ ergs $\text{s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$. Any excitation mechanism for the H_2 must be able to explain both the absolute intensity and the uniformity and extent of the emission. The excitation of the $\text{H}_2 \nu = 1$, $J = 3$ state can result

either from radiative decay from UV-excited electronic states or from energetic collisions. H₂ can absorb 91–123 nm photons in the $B^1\Sigma_u^+ - X^1\Sigma_g^+$ Lyman and $C^1\Pi_u - X^1\Sigma_g^+$ Werner bands. About 90% of the time, the excited H₂ decays to some ro-vibrational level of the ground electronic state. The relative line intensities that arise in UV-excited H₂ are insensitive to density or to UV field strength if $n_{\text{H}_2} < 10^4 \text{ cm}^{-3}$ (Black & van Dishoeck 1987). At densities $\geq 10^5 \text{ cm}^{-3}$, UV-excited gas can become hot enough that collisions populate states with $v = 1$ (Sternberg & Dalgarno 1989). Collisional excitation can also result from shocks that abruptly heat the gas to greater than 10^3 K (see, e.g., Hollenbach, Chernoff, & McKee 1989). Several observational results lead us to believe that UV excitation can explain the $v = 1 \rightarrow 0 S(1)$ emission in the Galactic center.

The denser parts of clouds like Orion and NGC 2024 produce H₂ emission with an intensity close to that observed in the Galactic center. In Orion and NGC 2024, the degree-scale H₂ emission has a typical surface brightness of $\sim 6 \times 10^{-6} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$ (Luhman et al. 1994). Along the molecular ridges in these clouds, the H₂ surface brightness is 3–5 times higher. Observations of H₂ transitions arising from higher lying states indicates that, in these sources, the $v = 1 \rightarrow 0 S(1)$ emission is a result of UV fluorescence.

If large-scale H₂ emission arises in the surface layers of the clouds where UV photons can excite the molecules, the dust, which absorbs the bulk of the incident flux, ought to radiate in the far-IR continuum as well. Luhman & Jaffe (1996) have compared the H₂ $v = 1 \rightarrow 0 S(1)$ observations of clouds in the Galactic disk with *IRAS* far-IR continuum results and derived a relation between the H₂ $v = 1 \rightarrow 0 S(1)$ line and far-IR continuum intensities. We can use this relationship and the measured far-IR intensities in the Galactic center to see if UV excitation is plausible for our observed $v = 1 \rightarrow 0 S(1)$ emission. In most of the region along our Galactic center H₂ cut, the *IRAS* 100 μm -band detectors were saturated. In order to compare the Galactic center H₂ data to far-IR continuum measurements with comparable angular resolution, we have combined the unsaturated *IRAS* measurements from the outer ends of our H₂ strip with the 40–250 μm continuum measurements of Odenwald & Fazio (1984). To make the two data sets comparable, we first converted the *IRAS* 60 μm and 100 μm fluxes into a total far-IR flux (the FIR parameter of Fullmer & Lonsdale 1989). The *IRAS* total far-IR flux agrees well with the far-IR flux derived by Odenwald & Fazio in the regions where their data and the unsaturated *IRAS* data overlap. We then converted the combined data sets into integrated far-IR intensity for comparison with our H₂ strip. We used the Luhman & Jaffe Galactic disk H₂ data set to rederive their H₂/far-IR relation in intensity units. We obtain

$$\log I_{\text{H}_2, v=1-0 S(1)} = -4.65 + 0.39 \log I_{\text{FIR}},$$

where both intensities are in $\text{ergs s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$. The dispersion of the Galactic disk cloud H₂ intensities about this relation is $\log \sigma = 0.23$. If we deredden the Galactic center H₂ observations by $A_K = 2.5 \text{ mag}$ (*but otherwise do nothing to fit the data to the Galactic disk relation*), the Galactic center H₂ intensities have a dispersion $\log \sigma = 0.26$ about this relation. The Galactic center results are therefore completely consistent with the empirical far-IR versus H₂ relationship derived for the UV-excited surfaces of clouds in the Galactic disk.

We can also compare the H₂ line intensities predicted by

models of PDRs to the observed intensities. The models use indirect observations of the far-UV field in the Galactic center (radio and far-IR continuum fluxes) as inputs. For the radio, we predict the far-UV field using emission from ELD H II regions because the molecular cloud column densities, and therefore the extinction at the wavelength of H₂, tend to be high (and uncertain) toward the discrete H II regions. Away from discrete H II regions, the typical 10.5 GHz flux density is 2.2 Jy in a 3'3 beam (Sofue 1985). Assuming $T_e = 10^4 \text{ K}$, this flux density corresponds to 2.3×10^{49} Lyman continuum photons per second (Mezger, Smith, & Churchwell 1974) in the corresponding region (8.2 pc). For an ionizing source with an effective stellar temperature $T_{\text{eff}} = 3.5 \times 10^4 \text{ K}$ as the UV source, the 2.3×10^{49} Lyman continuum photons imply $\sim 2.3 \times 10^{49}$ photons in the range that can excite the H₂ (91–123 nm), or a luminosity of $1.2 \times 10^5 L_\odot$ (Black & van Dishoeck 1987). From our observations, the average H₂ flux in a 3'3 beam is $2.4 \times 10^{-12} \text{ ergs s}^{-1} \text{ cm}^{-2}$. The corresponding total H₂ luminosity in the 8.2 pc (3'3) region is $3.3 \times 10^3 L_\odot$, if we correct for an extinction of $A_K = 2.5 \text{ mag}$ and use the PDR model of Black & van Dishoeck (1987) to extrapolate to the H₂ cooling in all lines [$I_{\text{H}_2, v=1-0 S(1)}/I_{\text{H}_2} = 0.016$]. The ratio of the near-IR H₂ luminosity to the luminosity in the far-UV band that is effective in exciting H₂ is 0.028, which is close to the value of 0.034 from an appropriate PDR model for the Galactic center (model 19 in Black & van Dishoeck 1987, which has $n_{\text{H}} = 10^4$ and a UV field $I_{\text{UV}} = 10^3$).

The far-IR continuum intensities along our Galactic center strip are typically $0.8 \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$ (Odenwald & Fazio 1984). If all of this emission arises from a single molecular cloud surface filling the beam, it corresponds to a far-UV flux of $\sim 2 \times 10^3$ times the mean interstellar radiation field in the solar neighborhood (Draine 1978). Given the likely number of clouds along each line of sight and various geometric effects, the likely far-UV field is 500–1000 times the solar neighborhood value. For this range of UV field strengths and densities between 3×10^3 and $3 \times 10^4 \text{ cm}^{-3}$, Black & van Dishoeck (1987) predict H₂ $v = 1 \rightarrow 0 S(1)$ line intensities in the range $(1.2\text{--}4.2) \times 10^{-5} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$, bracketing our typical observed, dereddened value.

The H₂ emission from the circumnuclear disk appears to be collisionally excited [$I_{v=2-1 S(1)}/I_{v=1-0 S(1)} \simeq 0.1$; Gatley et al. 1984]. Gatley et al. suggested that shocks created by mass outflow from the Galactic nucleus might excite the H₂. Such thermal line ratios can also occur, however, in UV-excited gas if the UV fields and densities are sufficiently high (Sternberg & Dalgarno 1989; Luhman et al. 1996). Since the typical hydrogen density in the circumnuclear disk is large, i.e., $n_{\text{H}} \simeq 10^5 \text{ cm}^{-3}$, and the UV field is intense in the central 3 pc (the number of total Lyman continuum photons absorbed by the gas is $\sim 2 \times 10^{50} L_\odot$; Lacy et al. 1980), the strength and character of the H₂ emission from the circumnuclear disk are also consistent with UV excitation.

4.3. Shock-Excitation

Shock-excitation of the H₂ $v = 1 \rightarrow 0 S(1)$ transition must take place, at some level, in the inner 400 pc of the Galaxy. A large variety of dynamical activity may give rise to shocks with appropriate characteristics. Outflows around newly formed stars and shocks caused by supernova remnants' impinging on molecular clouds in the Galactic disk both produce H₂ emission and should be observable in the Galactic center. Bally et

al. (1987, 1988) surveyed the Galactic center region in the ^{12}CO and ^{13}CO $J = 1 \rightarrow 0$ and $\text{CS } J = 2 \rightarrow 1$ lines. The gas distribution is highly asymmetric about the center, and some negative-velocity gas is seen at positive longitudes, which is “forbidden” to gas in circular orbits. This gas and other clouds with eccentric orbits may collide with material in more circular orbits. For example, Hasegawa et al. (1994) suggested that in the Sgr B₂ complex a dense ($n_{\text{H}_2} \approx 1.4 \times 10^4 \text{ cm}^{-3}$), $10^6 M_\odot$ “clump” has collided with the extended, less dense “shell” of the cloud complex, producing a $\sim 30 \text{ km s}^{-1}$ shock. Finally, the internal velocity dispersion of the molecular clouds is in the range $\Delta V = 20\text{--}50 \text{ km s}^{-1}$ (Bally et al. 1988). If the internal collisions efficiently dissipate the relative kinetic energy by radiative cooling following shocks, there should be H_2 emission throughout the molecular clouds, much of it, however, heavily extinguished.

Depending on the context, shock-excited H_2 emission could result either from dissociative J-shocks (colliding clouds, supernova remnants) or from C-shocks (outflows, dissipation of turbulence). The J-shocks give rise to $\text{H}_2 v = 1 \rightarrow 0 S(1)$ intensities in the range 3×10^{-5} to $10^{-4} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$, with the intensity’s being fairly insensitive to density and shock velocity over the range $10^4 \text{ cm}^{-3} \leq n \leq 10^5 \text{ cm}^{-3}$ and $30 \text{ km s}^{-1} \leq V_{\text{shock}} \leq 150 \text{ km s}^{-1}$ (Hollenbach & McKee 1989). For $A_K = 2.5$ mag, the predicted intensity matches what we observe in the Galactic center fairly well. In order to explain the distribution of observed H_2 emission, however, the number of shock fronts times the area covered per beam must roughly equal the beam area along virtually every line of sight through the inner 400 pc of the Galaxy, an unlikely picture at best.

C-shocks can produce $\text{H}_2 v = 1 \rightarrow 0 S(1)$ intensities in the range of those observed in the Galactic center. A single C-shock with $n = 10^4 \text{ cm}^{-3}$ and $V = 20 \text{ km s}^{-1}$ gives $I_{S(1)} \approx 3 \times 10^{-5} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$ (Draine, Roberge, & Dalgarno 1983). The emergent intensity, however, is extremely sensitive to the shock’s velocity, varying (at $n_{\text{H}} = 10^4 \text{ cm}^{-3}$) by 3 orders of magnitude from $V_{\text{shock}} = 18$ to $V_{\text{shock}} = 40 \text{ km s}^{-1}$. A model that makes use of C-shocks to produce the observed uniform $\text{H}_2 v = 1 \rightarrow 0 S(1)$ distribution would have to be somewhat contrived. While there may be some shock-excited H_2 emission from the Galactic center, it is difficult to argue away the expected PDR emission and then construct a reasonably simple shock model capable of explaining the

observations. A reliable test of the excitation mechanism would be to observe H_2 transitions arising higher above ground than the $v = 1 \rightarrow 0 S(1)$ line.

4.4. Total H_2 Luminosity

To estimate the total H_2 luminosity of the Galactic center, we extrapolate from our 400 pc-long strip by assuming that the scale height of the H_2 emission equals that of the far-IR radiation ($h \approx 0.2$; Odenwald & Fazio 1984). For $A_K = 2.5$ mag and $f \leq 1$ (see § 4.1), the dereddened $\text{H}_2 v = 1 \rightarrow 0 S(1)$ luminosity in the inner 400 pc diameter of the Galaxy is $8.0 \times 10^3 L_\odot$. Joseph (1989) gives ranges of $\text{H}_2 v = 1 \rightarrow 0 S(1)$ luminosity in greater than 1 kpc regions for various classes of galaxies: (1) merging galaxies: $3 \times 10^6\text{--}3 \times 10^8 L_\odot$; (2) interacting galaxies: $10^5\text{--}10^7 L_\odot$; (3) barred spirals: $10^4\text{--}10^6 L_\odot$. Over its inner ~ 1 kpc, our Galaxy most likely falls within the range for barred spirals.

In ultraluminous IR-bright galaxies ($L_{\text{IR}} \approx 10^{12} L_\odot$), Goldader et al. (1995) show that $\log [L_{S(1)}/L_{\text{FIR}}] = -4.95 \pm 0.22$. For the nearby starburst M82, we can use $\text{H}_2 v = 1 \rightarrow 0 S(1)$ measurements of the inner $60''$ (S. Pak & D. T. Jaffe, unpublished) together with far-IR continuum observations (D. A. Harper, as quoted in Lugten et al. 1986) to derive $\log [L_{S(1)}/L_{\text{FIR}}] = -5.2$ for the inner 1 kpc. For the inner 400 pc of the Milky Way, the data presented here yield $\log [L_{S(1)}/L_{\text{FIR}}] = -4.8$.

There is evidence in some high-luminosity galaxies that the H_2 emission results from UV excitation. In NGC 3256, a merging galaxy, the $\text{H}_2 v = 2 \rightarrow 1 S(1)/v = 1 \rightarrow 0 S(1)$ line ratio in the 600 pc region ($3''.5 \times 3''.5$) is 0.39 ± 0.06 , which suggests that UV fluorescence is responsible for at least 60% of the $\text{H}_2 v = 1 \rightarrow 0 S(1)$ emission (Doyon, Wright, & Joseph 1994). If H_2 in the Galactic center is UV-excited, as we suggest here, this mechanism could be shared by H_2 emission from galaxies with an enormous range of nuclear conditions.

This work was supported by the David and Lucile Packard Foundation and by NSF grant AST 9117373. We thank John Lacy and the referee, Leo Blitz, for helpful comments, Michael Luhman and Al Mitchell for contributions to the Fabry-Perot Spectrometer Project, and Jacqueline Davidson and the staff of the McDonald Observatory for their assistance on the observing run.

REFERENCES

- Altenhoff, W. J., Downes, D., Pauls, T., & Schraml, J. 1978, *A&AS*, 35, 23
 Bally, J., Stark, A. A., Wilson, R. W., & Henkel, C. 1987, *ApJS*, 65, 13
 ———. 1988, *ApJ*, 324, 223
 Black, J. H., & van Dishoeck, E. F. 1987, *ApJ*, 322, 412
 Blitz, L., Binney, J., Lo, K. Y., Bally, J., & Ho, P. T. P. 1993, *Nature*, 361, 417
 Catchpole, R. M., Whitelock, P. A., & Glass, I. S. 1990, *MNRAS*, 247, 479
 Doyon, R., Wright, G. S., & Joseph, R. D. 1994, *ApJ*, 421, 115
 Draine, B. T. 1978, *ApJS*, 36, 595
 Draine, B. T., Roberge, W. G., & Dalgarno, A. 1983, *ApJ*, 264, 485
 Fullmer, L., & Lonsdale, C. 1989, *Catalogued Galaxies and Quasars Observed in the IRAS Survey: Version 2 (JPLD-1932)* (Pasadena: JPL)
 Gatley, I., Jones, T. J., Hyland, A. R., Beattie, D. H., & Lee, T. J. 1984, *MNRAS*, 210, 565
 Gatley, I., Jones, T. J., Hyland, A. R., Wade, R., Geballe, T. R., & Krisciunas, K. 1986, *MNRAS*, 222, 299
 Gatley, I., & Merrill, M. 1993, *Proc. SPIE*, 1946, 2
 Glass, I. S., Catchpole, R. M., & Whitelock, P. A. 1987, *MNRAS*, 227, 373
 Goldader, J. D., Joseph, R. D., Doyon, R., & Sanders, D. B. 1995, *ApJ*, 444, 97
 Güsten, R. 1989, in *IAU Symp. 136, The Center of the Galaxy*, ed. M. Morris (Dordrecht: Kluwer), 89
 Hasegawa, T., Sato, F., Whiteoak, J. B., & Miyawaki, R. 1994, *ApJ*, 429, L77
 Hollenbach, D. J., Chernoff, D. F., & McKee, C. F. 1989, in *Infrared Spectroscopy in Astronomy*, ed. B. Kaldeich (ESA SP-290) (Noordwijk: ESA), 245
 Hollenbach, D. J., & McKee, C. F. 1989, *ApJ*, 342, 306
 Joseph, R. D. 1989, in *Infrared Spectroscopy in Astronomy*, ed. B. Kaldeich (ESA SP-290) (Noordwijk: ESA), 439
 Lacy, J. H., Townes, C. H., Geballe, T. R., & Hollenbach, D. J. 1980, *ApJ*, 281, 132
 Lester, D. F., Carr, J. S., Joy, M., & Gaffney, N. 1990, *ApJ*, 352, 544
 Lis, D. C., & Carlstrom, J. E. 1994, *ApJ*, 424, 189
 Lugten, J. B., Watson, D. M., Crawford, M. K., & Genzel, R. 1986, *ApJ*, 311, L51
 Luhman, M. L., & Jaffe, D. T. 1996, *ApJ*, in press
 Luhman, M. L., Jaffe, D. T., Keller, L. D., & Pak, S. 1994, *ApJ*, 436, L185
 ———. 1995, *PASP*, 107, 184
 Luhman, M. L., Jaffe, D. T., Sternberg, A., Herrmann, F., & Poglitsch, A. 1996, in preparation
 Mezger, P. G., Smith, L. F., & Churchwell, E. 1974, *A&A*, 32, 269
 Moorwood, A. F. M., & Oliva, E. 1990, *A&A*, 239, 78
 Odenwald, S. F., & Fazio, G. G. 1984, *ApJ*, 283, 601
 Oort, J. 1977, *ARA&A*, 15, 295
 Puxley, P., Hawarden, T., & Mountain, M. 1989, in *Infrared Spectroscopy in Astronomy*, ed. B. Kaldeich (ESA SP-290) (Noordwijk: ESA), 517
 Sodroski, T. J., et al. 1994, *ApJ*, 428, 638
 Sofue, Y. 1985, *PASJ*, 37, 697
 Sternberg, A., & Dalgarno, A. 1989, *ApJ*, 338, 197
 Yusef-Zadeh, F., Morris, M., & Chance, D. 1984, *Nature*, 310, 557