

CALIBRATION AND SYSTEMATIC ERROR ANALYSIS FOR THE *COBE*¹ DMR 4 YEAR SKY MAPS

A. KOGUT,^{2,3} A. J. BANDAY,^{2,4} C. L. BENNETT,⁵ K. M. GÓRSKI,^{2,6} G. HINSHAW,² P. D. JACKSON,²
 P. KEEGSTRA,² C. LINEWEAVER,⁷ G. F. SMOOT,⁷ L. TENORIO,⁷ AND E. L. WRIGHT⁸

Received 1996 January 5; accepted 1996 May 21

ABSTRACT

The Differential Microwave Radiometers (DMR) instrument aboard the *Cosmic Background Explorer* (*COBE*) has mapped the full microwave sky to mean sensitivity 26 μK per 7° field of view. The absolute calibration is determined to 0.7% with drifts smaller than 0.2% per year. We have analyzed both the raw differential data and the pixelized sky maps for evidence of contaminating sources such as solar system foregrounds, instrumental susceptibilities, and artifacts from data recovery and processing. Most systematic effects couple only weakly to the sky maps. The largest uncertainties in the maps result from the instrument susceptibility to Earth's magnetic field, microwave emission from Earth, and upper limits to potential effects at the spacecraft spin period. Systematic effects in the maps are small compared to either the noise or the celestial signal: the 95% confidence upper limit for the pixel-pixel rms from all identified systematics is less than 6 μK in the worst channel. A power spectrum analysis of the $(A - B)/2$ difference maps shows no evidence for additional undetected systematic effects.

Subject headings: cosmic microwave background — instrumentation: detectors —
 radio continuum: general — space vehicles

1. INTRODUCTION

The large angular scale anisotropy of the cosmic microwave background (CMB) reflects the distribution of matter and energy in the early universe, before causal processes could operate to create the rich array of structures observed at the present epoch. Precise observations of the CMB anisotropy fix the initial conditions for different models of structure formation and have important implications for theories of the high-energy behavior at the earliest times (e.g., inflation, cosmic defects). The Differential Microwave Radiometers instrument on the *Cosmic Background Explorer* (*COBE*-DMR) has completed its 4 year observations of the microwave sky. It has detected statistically significant signals whose spatial morphology and frequency dependence are consistent with CMB anisotropy (Bennett et al. 1996; Bennett et al. 1994; Smoot et al. 1992). Microwave emission from the Galaxy is weak at high latitudes ($|b| > 30^\circ$) and, with the notable exception of the quadrupole, does not significantly contaminate the primordial CMB signal (Kogut et al. 1996a, 1996b; Bennett et al. 1992b). The CMB is isotropic to a high degree: the detected anisotropy $\Delta T/T$ ranges from 10^{-6} to a few parts in 10^{-5}

on angular scales larger than 7° . At this level of sensitivity, great care must be taken to demonstrate that the results are not affected by microwave emission from nearby objects hundreds to thousands of times hotter than the CMB (e.g., the spacecraft, Earth, Moon, and Sun) or from instrumental effects associated with changes in the orbital environment, telemetry, or data processing.

DMR is well characterized in terms of its response to various potential systematic effects. Kogut et al. (1992) describe the analysis techniques and upper limits on systematic artifacts in the first year sky maps. An important result of their analysis was that, owing to DMR's rapid scan pattern and good pixel-pixel connectedness, most systematic effects couple only weakly to the sky maps. Noncelestial signals from nearby objects or within the instrument need not be removed to microkelvin precision in the time domain for their effects in the map to be negligible. Bennett et al. (1994) provide similar analysis and upper limits for the 2 year sky maps.

In this paper, we present the instrument calibration and upper limits to systematic artifacts in the final 4 year sky maps using data from 1989 December 22 through 1993 December 21 UT. Absolute calibration from external sources observed in flight (the Moon and the Doppler dipole from Earth's orbital motion about the Sun) are in agreement with the more precise prelaunch calibration; the calibration is determined to 0.7% absolute with drifts smaller than 0.2% per year. We detect gain modulation $\Delta G/G \approx 10^{-5}$ at the orbit period during the 2 month periods surrounding the June solstice, when the spacecraft flies through the Earth's shadow each orbit. The resulting modulation of the radiometric offset explains a previously detected instrumental signal of unknown origin during these periods.

The largest uncertainties in the maps result from the instrument susceptibility to Earth's magnetic field, microwave emission from Earth, and upper limits to potential effects at the spacecraft spin period. We correct the data for

¹ The National Aeronautics and Space Administration/Goddard Space Flight Center (NASA/GSFC) is responsible for the design, development, and operation of the *Cosmic Background Explorer* (*COBE*). Scientific guidance is provided by the *COBE* Science Working Group.

² Hughes STX Corporation, Laboratory for Astronomy and Solar Physics, Code 685, NASA/GSFC, Greenbelt MD 20771.

³ E-mail: kogut@stars.gsfc.nasa.gov.

⁴ Current address: Max Planck Institut für Astrophysik, 85740 Garching, Germany.

⁵ Laboratory for Astronomy and Solar Physics, Code 685, NASA/GSFC, Greenbelt MD 20771.

⁶ On leave from Warsaw University Observatory, Aleje Ujazdowskie 4, 00-478 Warszawa, Poland.

⁷ LBL, SSL, and CfPA, Building 50-351, University of California, Berkeley, Berkeley, CA 94720

⁸ UCLA Astronomy, P.O. Box 951562, Los Angeles, CA 90095-1562.

the magnetic response; the resulting uncertainties are dominated by uncertainty in the magnetic field vector at each radiometer. With the full 4 year data set we detect emission from Earth when Earth is 4° below the Sun/Earth shields or higher (about 5% of the data); the detected signal is weak enough that correction is not required. There is no evidence for additional systematic effects at the spin period, but the precision with which we can rule out potential new effects is limited by the instrument noise. The quadrature sum of all systematic uncertainties in the 4 year maps, after correction in the time domain, yields upper limits $\Delta T < 4 \mu\text{K}$ rms over the full sky in the four most sensitive channels. Artifacts at this level would contribute less than 0.6% to the variance of the CMB signal at 7° resolution: systematic effects do not limit the DMR maps.

2. INSTRUMENT DESCRIPTION

DMR consists of six differential microwave radiometers, two nearly independent channels (labeled A and B) at frequencies 31.5, 53, and 90 GHz (wavelength 9.5, 5.7, and 3.3 mm). Each radiometer measures the difference in power between two regions of sky separated by 60° , using a heterodyne receiver switched at 100 Hz between two corrugated horn antennas with beam width 7° full width at half-maximum pointed 30° to either side of the spacecraft spin axis. The 31A and 31B channels share a common antenna pair in right and left circular polarizations; the other four channels have independent antenna pairs in a single linear polarization, for a total of 10 antennas. The DMR antennas are mounted in three boxes spaced 120° apart on the outside of the superfluid helium cryostat containing the Far Infrared Absolute Spectrophotometer (FIRAS) and the Diffuse Infrared Background Experiment (DIRBE). The E-plane of linear polarization in each of the 53 and 90 GHz antennas is directed radially outward from the spacecraft spin axis. A shield surrounds the aperture plane to block radiation from Earth and the Sun. The DMR antenna apertures lie approximately 6 cm below the shield plane. The combined motions of the spacecraft spin (75 s period), orbit (103 minute period), and orbital precession ($\sim 1^\circ$ per day) allow each sky position to be compared to all others through a highly redundant set of all possible difference measurements spaced 60° apart.

The switched signal in each channel undergoes RF amplification, detection, near-DC amplification, and synchronous demodulation. The demodulated signal is integrated for 0.5 s, digitized, and stored in an on-board tape recorder. A single local oscillator provides a common frequency reference for the mixer in the A and B channels at each frequency. The A and B channels also share a common enclosure and thermal regulation system but are otherwise independent. We also sample the switched signal after one stage of DC amplification but before synchronous demodulation; the resulting “total power” signal serves as a low-precision check on the system temperature and gain of the RF and first DC amplifier with only minor contributions from celestial signals.

COBE was launched from Vandenberg Air Force Base on 1989 November 18 into a 900 km, 99° inclination circular orbit that precesses to follow the terminator. Attitude control keeps the spacecraft pointed away from Earth and nearly perpendicular to the Sun. Solar radiation never directly illuminates the aperture plane. The Earth limb is below the shield for 95% of the mission. During the 2

months surrounding the June solstice, the attitude control cannot simultaneously block both terrestrial and solar emission, and the Earth limb rises above the shield as the spacecraft flies over the Arctic. During the same 2 months, the spacecraft flies through Earth’s shadow over the Antarctic. The resulting eclipse modulates spacecraft temperatures and voltages.

Several authors provide a more complete description of *COBE* and DMR. Boggess et al. (1992) provide a mission overview. Smoot et al. (1990) describe the DMR instrument. Toral et al. (1990) and Wright et al. (1994) describe the DMR beam patterns. Bennett et al. (1992a) describe the instrument preflight calibration and provide a schematic of the *COBE* aperture plane.

3. DATA PROCESSING

The digitized data from the on-board tape recorders are transmitted once per day to a ground station. A software program merges the uncalibrated DMR differential data with housekeeping (temperatures, currents, voltages, and relay states), spacecraft attitude, and selected spacecraft archives (magnetometers, momentum wheels, and electro-magnet currents). The uncalibrated time-ordered data $S_{ij}(t)$ may be represented as

$$S_{ij}(t) = \frac{1}{\mathcal{G}(t)} \left[T_i - T_j + O(t) + N(t) + \sum_k Z_k(t) \right],$$

where T_i and T_j are the antenna temperatures of the two regions of sky observed at time t , $\mathcal{G}(t)$ is the gain factor providing calibration between antenna temperature and telemetry units, $O(t)$ is the radiometric offset produced by small imbalances in the differential radiometer, $N(t)$ is the random instrument noise, and $Z_k(t)$ are the time-dependent systematic signals from noncosmological sources and instrumental effects. A second program calculates the gain factor $\mathcal{G}(t)$, the instrumental baseline $B(t)$, and corrections for known systematics $W_n(t)$ to determine the calibrated differential temperatures

$$D_{ij}(t) = \mathcal{G}(t)[S_{ij}(t) - B(t)] - \sum_n W_n(t), \quad (1)$$

which we refer to as the DMR time-ordered data set. We distinguish between the noncosmological signals $Z_k(t)$ and our models for these signals, $W_n(t)$, to account explicitly for modeling errors in these signals.

The instrument gain, beam patterns, and environmental susceptibilities of each radiometer have been measured in preflight testing (Smoot et al. 1990). Based on these and on in-flight data, we reject data taken when the uncertainty in the systematic error model $W_n(t)$ becomes unacceptably large. Table 1 lists cuts and corrections made in the 4 year DMR data processing. We flag as unusable any datum for which the telemetry is unavailable or of poor quality or which deviates from the mean by more than 5 standard deviations. Such “spikes” are rarely noise outliers but result instead from observations of the Moon or instrument changes such as the end of the noise source calibration sequence. The exponential decay of the noise source power contaminates the first few samples after the noise sources have been commanded off; as a result, we conservatively discard the entire 32 s frame after each in-flight calibration.

We flag as unusable for sky maps any datum taken with

TABLE 1

CUTS AND CORRECTIONS MADE TO DMR TIME-ORDERED DATA

Effect	Cut ^a	Correction
No telemetry	Yes (0.4%)	No
Spike in data	Yes (0.3%)	No
Offscale data	Yes (0.3%)	No
Bad attitude	Yes (0.7%)	No
Unstable noise	Yes (4.2%) ^b	No
Calibration	Yes (2.2%)	No
Earth emission	Yes (5.0%)	No
Lunar emission	Yes (4.6%)	Yes
Planetary emission	No	Yes
Lock-in amplifier memory	No	Yes
Magnetic susceptibility	No	Yes
Earth Doppler	No	Yes
Satellite Doppler	No	Yes
Seasonal effects	Yes (26.0%) ^c	Yes

^a Numbers in parentheses are the fraction of data rejected for each cut.

^b 31B channel only.

^c 31A and 31B channels only.

the Moon within 21° of an antenna beam center, although we do use these data for lunar calibration and to map the beam pattern in flight. All other data are corrected for a model of lunar emission. We reject data taken when the Earth limb is 1° below the shield or higher (3° for the 31.5 GHz channels) but do not otherwise correct for Earth emission. We correct data taken with Mars, Jupiter, or Saturn within 15° of an antenna beam center but do not reject these data as unusable. We correct the time-ordered data for the instrument response to Earth's magnetic field, the 3.2% correlation between successive observations caused by the lock-in amplifier low-pass filter, and for the Doppler dipole from the satellite motion about Earth and Earth's orbital motion about the solar system barycenter. An orbitally modulated signal is present during the 2 month "eclipse season" surrounding the June solstice, related to thermally induced gain variations in the amplification chain. An empirical model based on tracers with high signal-to-noise ratio removes approximately $\frac{2}{3}$ of this signal. Estimated residuals, after correction, are small compared to the increase in data and sky coverage in the 53 and 90 GHz channels (recall that orbitally modulated signals couple only weakly to the sky maps, so that a small systematic signal accompanied by a larger increase in sensitivity can be a worthwhile trade-off). The amplitude of the effect, and hence the residual uncertainty, is larger in the 31A and 31B channels; we reject data taken during the period May 1 through August 4 of each year for the 31A channel and May 21 through July 24 for the 31B channel. The 31B channel suffered a permanent increase in receiver noise on 1991 October 4. We flag as unusable data when the instrument noise is unstable and use the noise during stable periods to weight all analyses. As a result, the 31B results are approximately 30% noisier than the 31A channel.

We fit the time-ordered data to a pixelized sky map and systematic terms by minimizing the χ^2 sum

$$\chi^2 = \sum_i \left[\frac{D_{ij}(t) - (T_i - T_j)}{\sigma_{ij}(t)} \right]^2, \quad (2)$$

where $\sigma_{ij}(t)$ is the instrument noise (Bennett et al. 1996). Note that the calibrated differential temperatures $D_{ij}(t)$ in equation (1) include model corrections $W_n(t)$ for non-

cosmological signals. We may vary the amplitude of these signals in equation (2) to replace the term $\sum_n W_n(t)$ with the more general $\sum_n a_n W_n(t)$. A sparse-matrix algorithm then performs the χ^2 minimization to yield a set of pixel temperatures T_i and coupling coefficients a_n to specified systematic effects (Janssen & Gulkis 1992; Jackson et al. 1992; Torres et al. 1989).

Several methods exist to derive limits to systematic signals $Z_k(t)$ in the time-ordered data and their projection ΔT_{sys} in the sky maps. Given a time-dependent model $W_n(t)$, a linear least-squares method solves simultaneously for the systematic coupling coefficient a_n in the time domain and the sky temperatures T_i in the map (eq. [2]). This procedure automatically removes the best estimate of the particular systematic signal from the sky map. We make a map ΔT_{sys} of the removed systematic effect by subtracting two solutions to equation (2) that differ only in the presence or absence of the model function $W_n(t)$. The residual uncertainty δT_{sys} after correction is determined by multiplying the systematic map ΔT_{sys} by the fractional uncertainty in the coupling coefficient:

$$\delta T_{\text{sys}} = \frac{\delta a_n}{a_n} \Delta T_{\text{sys}}. \quad (3)$$

In some cases, the model function $W_n(t)$ may not be well specified a priori. We may rebin the time-ordered data into a coordinate system in which the signal will add coherently to obtain limits to both the shape and amplitude of the systematic function $Z_k(t)$. We then use the binned data and associated uncertainties, sampled according to the DMR observation pattern, as the model function $W_n(t)$ in equation (2) to derive the systematic error map and uncertainty. For example, microwave emission from Earth diffracted over the shield has a complicated time dependence whose exact form depends sensitively upon the relative geometry of the Earth, shield, and horn antennas. We bin the data by the location of Earth relative to the *COBE* aperture plane and use the binned data both to evaluate various models W_n of the diffraction and to derive model-independent limits to Earth emission in the DMR maps. We also bin the time-ordered data by the spacecraft orbit and spin angles to search for signals at those periods.

Several tests limit systematic effects without requiring any a priori information. Fourier transforms of the calibrated $D_{ij}(t)$ provide a powerful tool to limit potential effects. Common-mode signals, particularly celestial emission, cancel in the $(A - B)/2$ "difference map" linear combination of the A and B channels at each frequency, or in similar difference maps from different time ranges in a single channel. Analysis of the difference maps provides model-independent limits to the combined effects of all systematic effects which are not identical in different channels or at different times.

The instrument noise is well described by a Gaussian probability distribution with standard deviation depending on the number of discrete observations N_i per pixel,

$$n_i = \frac{\sigma_0}{\sqrt{N_i}},$$

where σ_0 is the rms noise per 0.5 s observation (Bennett et al. 1996). Figure 1 shows the instrument noise sorted by the number of observations linking two pixels at 60° separation. There is no evidence for a "noise floor" from effects that do not average out with time. Figure 2 shows the




FIG. 1.—Instrument noise for the 53B radiometer, sorted by integration time. The time-ordered data are first sorted by the orientation of the two antennas, and all observations with the spacecraft in a fixed position in inertial coordinates are combined. Temperature patterns fixed on the sky affect the mean temperature difference of each such pixel combination but do not affect the standard deviation. The plot shows the uncertainty in the mean values (“instrument noise”). There is no evidence for a noise floor caused by noncelestial signals.

observation pattern in the 53B channel for the 4 year mission after applying all cuts. Correlations between pixels caused by the 60° antenna separation are negligible (Lineveaver et al. 1994). Because *COBE* is in a polar orbit, regions approximately 30° from the celestial poles are

observed each orbit, while regions near the celestial equator are only observed for 4 months of each year. The lunar cut further reduces usable data near the ecliptic plane. Since the model functions $W_n(t)$ are typically not well described by a set of temperatures T_i fixed on the sky, gradients in the sky




FIG. 2.—Noise pattern of the 53B channel for 4 years of data. The map is a Mollweide projection in Galactic coordinates. The lunar cut causes the striping in the ecliptic plane. The map also shows the north-south asymmetry near the ecliptic poles.

TABLE 2
CALIBRATION SUMMARY^a

CHANNEL (1)	ABSOLUTE CALIBRATION		RELATIVE A/B (%) (4)	LINEAR DRIFT (% yr ⁻¹) (5)	ORBIT DRIFT (10 ⁵ Δ \mathcal{G} / \mathcal{G}) (6)	SPIN DRIFT (10 ⁷ Δ \mathcal{G} / \mathcal{G}) (7)
	Ground (%) (2)	Flight (%) (3)				
31A	0.0 ± 2.5	+2.2 ± 3.0	0.40 ± 0.05	+0.23 ± 0.33	6.7	15
31B	0.0 ± 2.3	+1.6 ± 3.8		+0.03 ± 0.81	22.5	55
53A	0.0 ± 0.7	-0.8 ± 1.2	0.27 ± 0.02	-0.12 ± 0.03	2.6	20
53B	0.0 ± 0.7	-0.1 ± 1.4		-0.11 ± 0.11	2.4	13
90A	0.0 ± 2.0	+2.6 ± 2.5	0.01 ± 0.03	-0.12 ± 0.04	5.4	13
90B	0.0 ± 1.3	-1.7 ± 1.8		-0.20 ± 0.11	7.2	8

^a These corrections to the ground calibration have *not* been applied to the data. Uncertainties are 68% confidence level except for the spin and orbit drifts, which are 95% confidence upper limits.

coverage are apparent in many of the systematic error maps.

4. CALIBRATION

The primary DMR calibration consists of the radiometric comparison, before launch, of cold (~ 77 K) and warm (~ 300 K) full-aperture blackbody targets. In addition, noise sources inject ~ 2 K of broadband power into the front end of each radiometer between the horn antenna and the Dicke switch. Near-simultaneous observations of the blackbody targets and the noise sources calibrate the antenna temperature of each noise source and permit the transfer of the blackbody calibration standard to the flight observations (Bennett et al. 1992a). Observations of the Moon and CMB dipole provide additional in-flight calibration. Table 2 summarizes the DMR calibration over the 4 year mission.

4.1. Absolute Calibration

The preflight absolute calibration has been adjusted for two effects observed in the first year of the mission: the 31B gain was decreased by 4.9%, and the 90B gain was increased by 1.4% (Kogut et al. 1992). Errors in the absolute calibration create systematic artifacts in two ways: errors in the amplitude of detected structures in the sky and artifacts in maps from which a model of celestial emission (e.g., lunar emission) have been removed. Observations of the Doppler dipole from Earth's orbital velocity provide an independent absolute calibration in flight. Motion with velocity β through an isotropic blackbody radiation field at temperature T_0 creates an observed thermodynamic temperature distribution

$$T(\theta) = \frac{T_0(1 - \beta^2)^{1/2}}{1 - \beta \cos(\theta)} \approx T_0[1 + \beta \cos(\theta) + O(\beta^2)] \quad (4)$$

(Peebles & Wilkinson 1968). The CMB spectrum is well described by a blackbody with $T_0 = 2.728$ K (Fixsen et al. 1996). The dipole caused by Earth's orbital velocity about the solar system barycenter (29.27 to 30.27 km s⁻¹) provides a calibration signal $\Delta T \approx 270$ μ K whose known spatial and temporal dependence allows simple separation from other astrophysical signals. Figure 3 shows the modulation of the CMB dipole observed in the 53B channel throughout the 4 year DMR mission. We fit the time-ordered data for a fixed sky map plus a Doppler calibration term, including the change in the orbital speed from the eccentricity of Earth's orbit. Table 3 compares the Doppler calibration to the primary calibration. Values larger than unity indicate a channel in which the observed Doppler dipole is larger than

predicted. The Doppler calibration is in agreement with the more precise prelaunch absolute calibration and shows no evidence for calibration shifts at the few percent level.

If we accept the prelaunch calibration as accurate, the Earth Doppler dipole provides a determination of the absolute CMB temperature T_0 . We may use equation (4) and the Doppler calibration factors from Table 3 to infer the CMB absolute temperature: $T_0 = 2.83 \pm 0.07$ K at 31.5 GHz, 2.71 ± 0.03 K at 53 GHz, and 2.72 ± 0.04 K at 90 GHz for a weighted mean $T_0 = 2.725 \pm 0.020$ K (68% confidence limit [CL]), in excellent agreement with the FIRAS spectrum.

Figure 4 shows the Doppler dipole resulting from COBE's 7.5 km s⁻¹ orbital velocity about Earth. We have mapped the time-ordered data in a coordinate system comoving with the satellite velocity vector and have binned the resulting map temperatures by the angle from the velocity vector. The attitude control system prevents DMR from observing all angles in this coordinate system. Within the range of angles observed, we find good agreement with the predicted 70 μ K dipole. The COBE velocity dipole demonstrates the sensitivity to signals $\Delta T/T \sim 10^{-5}$.

The Moon provides a second external absolute calibration source. DMR observes the Moon in the antenna main beam for ~ 6 days each at both first and third quarters (Bennett et al. 1992a). The antenna temperature of the Moon within integration time τ is given by

$$T_{A, \text{Moon}} = \frac{1}{\tau} \int_t^{t+\tau} dt \int d\Omega T_{\text{Moon}} P[\theta(t), \phi(t)] \Big/ \int d\Omega P(\theta, \phi),$$

where $P(\theta, \phi)$ is the DMR beam pattern. We integrate the physical temperature and microwave emission properties across the lunar disk to estimate T_{Moon} as a function of lunar phase and Sun-Moon distance (Keihm 1982; Keihm & Gary 1979; Keihm & Langseth 1975). The gain factors derived from observations of the Moon show a pronounced dependence on lunar phase, with peak-to-peak variation of 3.7%, 5.3%, and 6.2% at 31.5, 53, and 90 GHz, respectively.

TABLE 3
ABSOLUTE CALIBRATION COMPARED TO NOISE SOURCES

Channel	Ground	Doppler	Moon
31A	1.000 ± 0.025	1.022 ± 0.030	1.021 ± 0.038
31B	1.000 ± 0.023	1.116 ± 0.058	1.016 ± 0.038
53A	1.000 ± 0.007	0.992 ± 0.012	1.021 ± 0.054
53B	1.000 ± 0.007	0.999 ± 0.014	1.018 ± 0.054
90A	1.000 ± 0.020	1.026 ± 0.025	1.014 ± 0.063
90B	1.000 ± 0.013	0.983 ± 0.018	1.013 ± 0.063




FIG. 3.—Modulation of the CMB dipole resulting from the Doppler effect of Earth's orbital motion about the Sun (channel 53B). Each datum represents 5 days. The amplitude of the modulation provides an independent absolute calibration.




FIG. 4.—53 GHz (A+B)/2 summed data mapped in a coordinate system fixed with respect to the satellite orbital velocity vector. The data have been binned by angle relative to the *COBE* orbital velocity vector and corrected for the 0.5° smoothing of the DMR beam. The attitude control system prevents full coverage in this coordinate system. The Doppler effect provides a known signal with amplitude $\Delta T/T \sim 10^{-5}$.

Longer term analysis shows an additional annual modulation with peak-to-peak amplitude 2%. These apparent gain modulations are not present in any method that does not involve lunar observations (i.e., on-board noise sources or the CMB dipole). We thus ascribe them to real time-dependent effects in T_{Moon} that are not duplicated in the Keihm model. See Jackson et al. (1996) for further analysis of T_{Moon} .

The lunar calibration modulation is repeatable over the 4 year mission and may be removed empirically to provide a stable “standard candle” for relative gain analysis. We adopt the combined peak-peak modulation as an estimate of the uncertainty of the lunar model for the absolute calibration. Table 3 shows the mean lunar calibration compared to the noise source calibration. The channel-averaged lunar absolute calibration is 1.8% larger than the prelaunch absolute calibration, well within the systematic limitations of the lunar model.

The Moon also serves to cross-calibrate the A and B channels. Systematic uncertainties in the model for $T_{A, \text{Moon}}$ cancel in the A/B ratio at each frequency. The resulting ratio (Table 4) places a limit to how well celestial emission will be expected to cancel in the $(A-B)/2$ “difference maps.” A similar analysis for the Earth velocity Doppler effect is consistent within much larger uncertainties. The A and B channels are cross-calibrated within 0.4%, well within the accuracy of the preflight calibration.

4.2. Relative Calibration

The noise sources are commanded on for 128 s every 2 hr. Provided the power broadcast by each device is constant in time, they provide a standard to monitor time-dependent changes in the instrument calibration. Figure 5 shows the calibration $\mathcal{G}(t)$ derived from the noise sources for the 53A and 53B channels over the full 4 year mission. The gain is stable to better than 3% throughout the 4 year mission. Gain drifts per se do not create systematic artifacts provided the noise sources track the true calibration $\mathcal{G}(t)$.

Each noise source is observed in both the A and B channels. The ratio of the two noise sources in a single channel provides information on the noise source stability, since the instrument calibration cancels. The ratio of a single noise source observed in two channels provides information on gain stability, since the noise source performance cancels. Based on these ratios, we correct the data for three step changes in noise source broadcast power: a 0.69% increase in power for the 90 GHz “down” noise source on 1990 March 17, a 0.69% increase in power for the 31 GHz “down” noise source on 1992 February 11, and a 0.34% increase in power for the 90 GHz “down” noise source on 1993 November 26. An additional anomaly occurred on 1993 September 1 in the 90 GHz data, when both noise sources showed step changes in a pattern inconsistent with a simple change in power or instrument calibration. A 0.29% step increase occurred for the 90A channel “up” noise source that was not mirrored by the same noise source

observed in the 90B channel. At the same time, the “down” noise source in the 90B channel decreased in power by 0.66%, unaccompanied by a similar change in the 90A channel. Since a calibration change would show up as a step in both the “up” and “down” noise source signals, while a noise source power change would show up in both the 90A and 90B channels, we can rule out simple models involving a single component. We currently have no explanation for this event but remove its effects in software by increasing the “up” noise source by 0.29% for the 90A channel only and decreasing the “down” noise source by 0.66% for the 90B channel only.

4.2.1. Long-Term Calibration Drifts

The gain solutions in the 90A and 90B channels are corrected for linear drifts of $0.81\% \text{ yr}^{-1}$ and $0.87\% \text{ yr}^{-1}$, respectively (Kogut et al. 1992; Bennett et al. 1994). We place limits on uncorrected long-term drifts in noise source power by examining the ratio of the A and B noise sources in each channel. The instrument calibration cancels in this ratio, which places a lower limit to the long-term accuracy of the calibration solution (which assumes that the noise sources do not change in time). Figure 6 shows this ratio for both the 53A and 53B channels. Channel 53A shows a stable ratio for the first year followed by a linear drift, while channel 53B shows an approximately linear drift throughout the mission. The fact that the ratio of the same physical devices does not have the same shape in both channels indicates that the observed changes are caused by a process more complicated than a simple change in noise source broadcast power. Table 5 shows the limits to linear drifts in all 6 channels based on the noise source ratios.

The CMB dipole provides a continuously observed signal of 3 mK amplitude, which we use to limit long-term errors in the noise source calibration. We fit the time-ordered data to the form

$$\Delta T = (1 + bt) \sum_{\ell=1}^2 \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m},$$

i.e., a spatially fixed dipole and quadrupole whose amplitudes change linearly with time. Since the CMB is effectively a constant for the 4 years of DMR observations, we can interpret the fitted parameter b in terms of a linear drift in the true calibration $\mathcal{G}(t)$ relative to the noise source solution $\mathcal{G}'(t)$. Table 5 shows the resulting limits to calibration drifts. There are no drifts significant at 95% confidence level.

The Moon also serves as a standard candle once the annual and phase variations are empirically removed. We have analyzed the ratio of the corrected lunar calibration to the noise source calibration to search for long-term drifts in the noise source solution (Table 5). The drifts inferred from the noise sources, dipole, and Moon are in general agree-

TABLE 4

RELATIVE A/B CALIBRATION COMPARED TO NOISE SOURCES

Frequency	Ground	Doppler	Moon
31	1.000 ± 0.026	0.915 ± 0.054	1.0040 ± 0.0005
53	1.000 ± 0.003	0.993 ± 0.018	1.0027 ± 0.0002
90	1.000 ± 0.010	1.044 ± 0.032	1.0001 ± 0.0003

TABLE 5
LINEAR CALIBRATION DRIFTS

Channel	NS Ratio (% yr ⁻¹)	CMB Dipole (% yr ⁻¹)	Moon (% yr ⁻¹)
31A	$+0.468 \pm 0.004$	$+0.365 \pm 0.278$	-0.140 ± 0.024
31B	-0.576 ± 0.015	$+0.947 \pm 1.281$	-0.293 ± 0.029
53A	-0.139 ± 0.001	-0.091 ± 0.122	-0.131 ± 0.009
53B	-0.191 ± 0.001	$+0.023 \pm 0.139$	-0.147 ± 0.011
90A	-0.099 ± 0.003	-0.173 ± 0.209	-0.098 ± 0.018
90B	-0.131 ± 0.001	-0.325 ± 0.174	-0.140 ± 0.015




FIG. 5.—Calibration factor $\mathcal{G}(t)$ derived from on-board noise sources for the 53A and 53B channels. Each datum represents 1 week. The gray bands show the “eclipse season” surrounding the June solstice. The gain is stable to better than 3% throughout the 4 year mission.

ment and provide some evidence that calibration drifts are dominated by changes in the power emitted by the noise sources. The significance of the coefficients is difficult to evaluate. The uncertainties in Table 5 are statistical only. Since the lunar results are possibly affected by long-term artifacts related to the annual and phase variations and since the noise source ratios are clearly more complicated than a simple linear drift, we do not use a weighted estimate of the three techniques. Instead, we adopt the unweighted mean and use the scatter among the three techniques as an estimate of the uncertainty. Table 2 shows the resulting limits on long-term calibration drifts in the 4 year DMR data set. Drifts from any source are small compared to the absolute calibration uncertainty.

4.2.2. Orbital Calibration Drifts

The noise sources provide direct calibration information every 2 hr. We interpolate the noise source gain solution $\mathcal{G}(t)$ using 48 hr of data fitted to a cubic spline with one interior knot. The resulting gain solutions are smooth at the daily boundaries but can not respond to gain variations with periods shorter than about 8 hr. We estimate gain variations at the orbit period by fitting a long-term baseline

to the noise source calibrations and binning the residuals by the spacecraft orbit angle relative to the ascending node. Long-term plots of the calibration (Fig. 5) show effects related to the “eclipse season” surrounding the June solstice. Figure 7 shows the binned calibration residuals during eclipse season and for “noneclipse” data (the rest of each year). The noise sources (*top panels*) show gain variation with amplitude $\Delta\mathcal{G}/\mathcal{G} \approx 1.6 \times 10^{-4}$ during the eclipse season. The shape of the gain variation is nearly identical to the spacecraft temperature variations, here (*superimposed solid line*) represented by a thermistor in the Instrument Power Distribution Unit (IPDU). Since the IPDU is not thermally controlled, it shows larger temperature variations that are less affected by the housekeeping digitization. The total power (*middle panels*), binned in a similar fashion, shows similar modulation, supporting the existence of a real orbitally modulated gain variation during the eclipse season. The amplitude of the total power variation, $\Delta P/P \approx 6.1 \times 10^{-5}$, is smaller than the noise source variation, as expected since the total power does not sample the entire amplification chain.

The mixer/preamp assembly and lock-in amplifier are maintained in a thermally controlled box. Figure 7 also




FIG. 6.—Ratio of “up” and “down” noise sources vs. time for 53A and 53B channels. Each datum represents 1 week. The gray bands show the “eclipse season” surrounding the June solstice. Uncorrected linear drifts are smaller than 0.2% per year at 95% confidence.

shows the lock-in amplifier temperature (*bottom panels*) during eclipse and noneclipse data. Thermal variations of ~ 10 mK amplitude at the orbit period are observed during eclipse season, compared to ~ 6 K for the unregulated IPDU. The thermal susceptibility of the amplifiers, measured prior to launch, is $\Delta\mathcal{G}/\mathcal{G} \approx 1\% \text{ K}^{-1}$ (Bennett et al. 1992a). The observed orbital gain variations are consistent with thermal modulation at the 10 mK level.

The existence of gain modulation $\Delta\mathcal{G}/\mathcal{G} \approx 10^{-4}$ with waveform similar to the IPDU thermistor provides a plausible explanation for a previously detected signal during eclipse season (§ 5.2; see also Kogut et al. 1992 and Bennett et al. 1994). An empirical fit to the IPDU thermistor and voltage monitors removes most of this signal; we do not explicitly correct the noise source solutions to model the orbital gain variation during eclipse season. Outside of eclipse season, the orbital environment is stable. The amplifier temperatures and direct noise source calibration show no modulation at the orbit period. The total power shows slight orbital modulation linked to variations in the input celestial signal from the CMB dipole and Galactic plane. There is no evidence for orbitally modulated gain variation

outside of the eclipse season at the level of a few parts in 10^5 (Table 2, col. [6]).

The noise source calibrations require 128 s, longer than the 73 s spin period. We obtain limits on calibration modulation at the spin period by binning the total power signal (sampled every 8 s) and scaling the resulting limits using the ratio of total power to gain variations observed at the orbit period, $\Delta\mathcal{G}/\mathcal{G} = 3.1\Delta P/P$. We find no variation in either the total power or the amplifier temperature at the spin period (Table 2, col. [7]), with limits $\Delta\mathcal{G}/\mathcal{G} < 2 \times 10^{-6}$ (95% CL).

Gain modulation can create systematic artifacts in two ways: through observations of the sky at different relative calibration (“striping”) and by modulating the radiometric offset. The relative importance of the two effects depends on the timescale. Over long periods, the offset signal $\Delta\mathcal{G}/\mathcal{G} \times O(t)$ is removed by the fitted baseline, leaving striping as the primary gain artifact. On timescales comparable to the spin or orbit periods, offset modulation becomes dominant. Since the offsets can approach 1 K, a gain modulation as small as 10^{-4} can create a 100 μK signal. We derive upper limits on the systematic artifacts in the DMR sky maps resulting from calibration drifts at the spin, orbit, or longer




FIG. 7.—Calibration signals binned at the orbit period for eclipse data (*left-hand panels*) and noneclipse data (*right-hand panels*). *Top*: Noise source calibration residuals. *Middle*: Total power residuals. *Bottom*: Lock-in amplifier temperature. The noise source and total power plots during eclipse season are overlaid with the scaled IPDU thermistor temperature to demonstrate the similarity of the waveforms (see text).

periods by adding terms to equation (2) of the form $W_{\text{cal}} = \Delta\mathcal{G}[S_i(t) + B(t)]$, where $\Delta\mathcal{G}(t)$ is taken from the 95% confidence upper limits in Table 2. That is, we simulate the effect of an uncorrected calibration modulation in the striping of the sky, the modulation of the instrumental offset, and possible cross-talk with other systematic effects. Systematic artifacts from residual calibration drifts and modulation are negligible, creating rms variations 0.4 μK or smaller (95% CL) in the most sensitive sky maps.

5. ENVIRONMENTAL EFFECTS

The *COBE* orbit provides a generally benign environment for the DMR instrument. The radiometers are above Earth's atmosphere, and the terminator-following orbit prevents large temperature changes. Kogut et al. (1992) review various effects associated with the orbital environment. The largest signals result from the response of the Dicke switch to Earth's magnetic field and the thermal response of the amplifiers to temperature changes when the orbit passes into Earth's shadow. We correct for both of these effects; residual artifacts in the sky maps are small.

5.1. Magnetic Susceptibility

The amplification chain in each channel is connected to the antennas using a latching ferrite circulator switched at 100 Hz by an applied magnetic field. An external magnetic field (from Earth or the electromagnets used to control the spacecraft angular momentum) modulates the insertion loss of the switch and creates a time-dependent signal described by the vector coupling

$$W_{\text{magnetic}}(t) = \boldsymbol{\beta} \cdot \mathbf{B}(t), \quad (5)$$

where $\mathbf{B}(t)$ is the magnetic field vector. We express the magnetic susceptibility vector $\boldsymbol{\beta}$ in an orthonormal coordinate system fixed with respect to the spacecraft: β_X is the susceptibility along the X -axis (antiparallel to the spin axis), β_R is the susceptibility along the radial axis (directed outward between two antennas), and β_T is the susceptibility along the transverse axis (from the positive horn to the negative horn). The antennas are pointed 30° to either side of the X -axis; magnetic signals from the β_X susceptibility are not spin modulated and produce a signal at the orbit period only. Both the R and T signals are spin modulated. The β_R

susceptibility produces an apparent temperature gradient oriented across the magnetic field (east-west) but at right angles to the antenna pointing. The β_T susceptibility produces an apparent temperature gradient oriented along the magnetic field (north-south) in phase with the antenna pointing. The inclination of the *COBE* orbit with respect to the magnetic poles breaks the degeneracy that would otherwise make the β_T susceptibility indistinguishable from a celestial dipole aligned with Earth's magnetic field.

Figure 8 shows the model magnetic signal $W_{\text{magnetic}}(t)$ for the 53B radiometer over the course of several orbits. The spacecraft spin modulates both the radial and transverse field components, causing the rapid variation at the spin period. The spacecraft orbital motion samples Earth's field at different latitudes, causing an orbital drift and modulating the amplitude envelope of the spin variations. The resulting signal is quite distinct from that expected from a fixed celestial signal, represented in Figure 8 by a model of lunar emission. The different time-dependent signatures greatly reduce the required accuracy of the magnetic model.

We simultaneously fit the time-ordered data in each channel for the temperature in each pixel and the magnetic susceptibility vector β (eqs. [2] and [5]). We obtain a significant improvement in χ^2 in each channel, demonstrating

that the DMR magnetic signals are not well described by *any* set of fixed pixel temperatures. More complicated models of the magnetic coupling (e.g., tensor or nonlinear terms) do not further reduce the χ^2 : a linear vector model (eq. [5]) is sufficient. We also fit β independently to each month of data and search for time dependence in the fitted coefficients. If the data are not corrected for the eclipse-related effects, all channels show anomalies in the β_X coefficients during the eclipse season. After correcting the time-ordered data for this effect (§ 5.2), the 31A and 31B channels show additional modulation of the β_X susceptibilities with a period of one year. We modify equation (5) to include additional terms of the form $\beta_{\text{annual}} \cos \theta \cos \phi$, where θ is the orbit angle relative to the ascending node and ϕ is the angle of the orbit plane relative to the vernal equinox. Since the resulting signal is not spin-modulated, it has almost no effect on the sky maps but is included to reduce cross-talk with other orbitally modulated effects. The 53 and 90 GHz channels show no significant variation in the fitted magnetic coefficients.

Table 6 lists the magnetic coefficients derived from 4 years of data. Slow changes in the magnetic signal from the orbitally modulated β_X susceptibility may be removed as part of the baseline. The values in Table 6 refer to the




FIG. 8.—Magnetic and celestial signals vs. time for 53B channel. *Top*: Magnetic signal $W_{\text{magnetic}}(t)$ from Earth's field. The spin and orbit modulation are clearly apparent. *Bottom*: Celestial signal from an unresolved source (the Moon). The orbital modulation results from sweeping the antenna beam pattern across the source at varying angles from beam center. The magnetic signal is easily distinguished from fixed celestial sources.

TABLE 6
MAGNETIC SUSCEPTIBILITY (mK G⁻¹)

Channel	β_x	β_R	β_T
31A.....	-0.173 ± 0.018	+0.262 ± 0.039	-0.196 ± 0.067
31B.....	+0.284 ± 0.024	+0.224 ± 0.054	+0.011 ± 0.089
53A.....	-1.514 ± 0.006	-0.081 ± 0.013	-0.881 ± 0.022
53B.....	+0.087 ± 0.006	-0.408 ± 0.015	-0.192 ± 0.025
90A.....	-0.135 ± 0.010	-1.175 ± 0.024	-0.334 ± 0.040
90B.....	-0.003 ± 0.007	+0.140 ± 0.017	-0.136 ± 0.028

two-orbit running mean baseline (Bennett et al. 1994) for which no such subtraction takes place.

We model the magnetic field $\mathbf{B}(t)$ using the spacecraft attitude solution and the time-dependent 1985 International Geomagnetic Reference Field (Barker et al. 1986) to order $\ell = 8$. We neglect the field components at higher ℓ and any contribution from the electromagnets used to dissipate spacecraft angular momentum. The amplitude of the field model varies from 196 to 402 mG over the *COBE* orbit. We limit deviations from the true field and our field model by using on-board magnetometers and find good agreement between the magnetometers and our application of the field model. We have examined the residuals for coherent behavior with respect to an inertial coordinate system (e.g., if the electromagnets preferentially fired with the spacecraft in a fixed location and attitude) and find none. The rms difference between the magnetometers and the field model is 8.9 mG averaged over the 4 year mission, which we adopt as the 68% CL uncertainty in the application of the model.

The simultaneous fit for the pixel temperatures and magnetic coefficients removes the fitted magnetic signals from the sky maps. The uncertainties in the fitted coefficients are dominated either by instrument noise (if $\beta \lesssim 0.3$ mK G⁻¹) or by the uncertainty in the magnetic field model. We obtain a map of the removed signal by subtracting the corrected sky map from a similar map for which no magnetic terms were fitted. We estimate the residual uncertainties after correction by multiplying this “effect” map by the fractional uncertainty in each fitted coefficient (eq. [3]). Figure 9 shows the 95% CL uncertainties in the 53B channel from magnetic effects. The magnetic residuals are among the largest systematic uncertainties in the DMR sky maps, but the amplitudes are small: the residual magnetic uncertainty, after correction, is less than 3 μ K rms in any channel.

5.2. Seasonal Effects

Previous analysis of the 1 and 2 year data sets showed the presence of an orbitally modulated signal during the

“eclipse season” surrounding the June solstice when the spacecraft repeatedly flies through Earth’s shadow (Kogut et al. 1992; Bennett et al. 1994). This “eclipse effect” is strongly correlated with various housekeeping signals, particularly the unregulated spacecraft temperatures and bus voltages which show the largest variation during eclipses and are thus least affected by the telemetry digitization (see Fig. 4 of Kogut et al. 1992). We model the effect empirically by fitting the time-ordered data to the form

$$W_{\text{eclipse}} = a\Delta T_{\text{IPDU}} + b\Delta V_{28},$$

where ΔT_{IPDU} is the temperature of the IPDU box and ΔV_{28} is the 28V bus voltage. We remove an orbital mean from both ΔT_{IPDU} and ΔV_{28} prior to fitting since long-term drifts are removed as part of the instrument baseline. The resulting peak-peak changes in the housekeeping signals are $\Delta T_{\text{IPDU}} \sim 5.3$ du, $\Delta V_{28} \sim 4.3$ du during eclipse season, and $\Delta T_{\text{IPDU}} \sim 0.3$ du, $\Delta V_{28} \sim 0.1$ du excluding eclipse season, where the housekeeping signals are expressed in digitized telemetry units (du). Table 7 shows the fitted coefficients for this empirical model during and excluding eclipse season. We detect the effect in all channels during eclipse season. Outside of eclipse season, the spacecraft temperatures and voltages are stable and the effect vanishes.

The detection of thermal gain variations at the orbit period during eclipse season (§ 4.2.2) provides a plausible mechanism for the eclipse effect. However, neither the observed gain variations nor the empirical housekeeping correlations remove the entire eclipse signal in all channels. The magnetic β_x coefficients for the 31A and 31B channels remain anomalously large during eclipse season even after correction for W_{eclipse} , indicating that the empirical model removes only $\frac{2}{3}$ of the signal in those channels. The eclipse effect creates artifacts in the maps in two ways: the direct projection of the signal into the maps and cross-talk with other orbitally modulated effects (particularly the β_x magnetic susceptibility). Both effects are important for the 31A and 31B channels; accordingly, we do not use data during eclipse season for these channels. The 53 and 90 GHz channels have smaller eclipse coefficients and show no β_x anomalies after correction. We correct the data using the empirical model and estimate the residual uncertainties in the sky maps using equation (3). Since the eclipse signal does not vary at the spin period, its projection onto the maps is small. Including the eclipse data in the 53 and 90 GHz channels adds less than 0.3 μ K rms artifacts to the maps (95% CL), much less than the 10 μ K reduction in noise gained by adding the 8 months of eclipse data to the 4 year data set.

TABLE 7
ECLIPSE COEFFICIENTS FOR EMPIRICAL MODEL

CHANNEL	THERMAL (mK du ⁻¹) ^a		VOLTAGE (mK du ⁻¹) ^a	
	During Eclipse	Excluding Eclipse	During Eclipse	Excluding Eclipse
31A.....	+0.442 ± 0.006	+0.046 ± 0.029	+0.311 ± 0.015	+0.018 ± 0.019
31B.....	+0.147 ± 0.008	-0.011 ± 0.038	+0.262 ± 0.019	-0.008 ± 0.027
53A.....	-0.022 ± 0.002	+0.001 ± 0.007	+0.013 ± 0.004	-0.002 ± 0.007
53B.....	-0.069 ± 0.003	-0.010 ± 0.010	-0.011 ± 0.004	+0.001 ± 0.007
90A.....	+0.042 ± 0.004	+0.032 ± 0.018	+0.024 ± 0.008	+0.009 ± 0.011
90B.....	0.000 ± 0.003	-0.018 ± 0.011	0.000 ± 0.005	+0.008 ± 0.011

^a Temperature and voltage housekeeping signals are processed in digitized telemetry units (du).




FIG. 9.—Full sky maps of 95% CL upper limits to the residual effect, after correction, of the magnetic susceptibility in the 53B channel. The maps are Mollweide projections in Galactic coordinates. *Top*: β_X susceptibility. *Middle*: β_R susceptibility. *Bottom*: β_T susceptibility. A fitted dipole has been removed from each map to show the higher order structure.

5.3. Orbit and Spin Effects

The spacecraft orbit and spin provide a natural period for any environmental effects. We test for the presence of additional effects, independent of an a priori model, by binning the corrected data $D_{ij}(t)$ by the orbit angle with respect to the ascending node and the spin angle with respect to the solar vector. With the exception of eclipse residuals for the 31A and 31B channels, there is no evidence for additional

effects; the binned data are compatible with instrument noise. Table 8 shows the resulting upper limits to the combined effects of any systematics at the spin and orbit periods, after correction for magnetic and seasonal effects. Artifacts in the maps from orbital effects at these levels are negligible (below $0.3 \mu\text{K}$ rms in any channel). Synchronous effects at the spin period couple more strongly to the sky maps. Figure 10 shows the artifacts in the sky maps

TABLE 8
95% CONFIDENCE UPPER LIMITS FROM ORBIT- AND
SPIN-MODULATED EFFECTS^a

CHANNEL	SPIN PERIOD All Data (μK)	ORBIT PERIOD	
		During Eclipse (μK)	Excluding Eclipse (μK)
31A	34	577	19
31B	41	489	47
53A	11	33	14
53B	10	19	12
90A	18	73	24
90B	8	21	18

^a All values are in units of antenna temperature.

resulting from spin-modulated signals at the 95% CL upper limit in Table 8. The noise limits on artifacts in the maps resulting from combined effects at the spin period, $\delta T_{\text{spin}} < 1.6 \mu\text{K}$ rms, are among the largest limits for the 4 year DMR maps.

6. FOREGROUND SOURCES

Emission from foreground sources within the solar system can create artifacts in maps of the microwave sky. We reduce artifacts from foreground sources by shielding the radiometers from the brightest sources (Earth and the Sun), correcting the data using models of source microwave emission, and rejecting data when uncertainties in the model become unacceptably large (Table 1).

6.1. Earth

Earth is the largest foreground source, emitting approximately 285 K over one-quarter of the sky. Emission from

Earth must be attenuated by 70 dB to reduce it below the 30 μK level of typical CMB anisotropies. DMR achieves this attenuation by using horn antennas with good off-axis side-lobe rejection and by interposing a shield between the radiometers and Earth. Radiation from Earth must diffract over the top of the shield before affecting the DMR data. The beam pattern, evaluated at the top of the shield, is typically -65 dB or lower, so only a modest attenuation from the shield is required.

We evaluate Earth emission by binning the time-ordered data $D_{ij}(t)$ by the position of the Earth limb in a coordinate system fixed with respect to the *COBE* spacecraft. Since Earth subtends an angle much larger than the 7° DMR beam, the azimuthal variation of the signal with respect to the spacecraft spin should reflect the antenna beam patterns with a positive lobe at the azimuth of horn 1 and a negative lobe at the azimuth of horn 2 in each channel. The signal change with respect to elevation angle as Earth sets below the shield depends on the details of the diffraction over the shield edge, while the overall normalization is set by the beam response at the top of the shield.

For most of the mission, the attitude control keeps Earth well below the Sun/Earth shield. During the eclipse season, Earth rises as high as 8° above the shield. When the Earth limb is above the shield, the binned data show a clear detection of the expected dual-lobed signal. At lower elevations, the signal decreases rapidly and falls below the noise. We derive limits to Earth emission at various elevation angles by fitting the binned data to a model of Earth emission based on scalar diffraction theory, the relative geometry of Earth, antennas, and shield, and the measured beam patterns of each antenna. Over narrow ranges of limb elevation angle (e.g., a strip 1 pixel high), the details of the diffraction




FIG. 10.—Full sky map of 95% CL upper limits to spin-modulated effects in the 53B channel (Mollweide projection in Galactic coordinates). A fitted dipole has been removed to show the higher order structure. The amplitude of the structure in this map is dominated by the instrument noise binned at the spin period.

become irrelevant, and only the azimuthal variation from the beam pattern is important.

The model (with fitted amplitude as a free parameter) provides a good fit to the binned data. Figure 11 shows the fitted amplitude versus elevation angle. When Earth is 5° above the shield, we recover a fitted amplitude $Z_{\text{Earth}} \approx 200 \mu\text{K}$, approximately 50% of the expected amplitude. This is well within the precision of the Earth model, whose overall normalization depends on the exact position of the deployed shield: a 6 cm shift in shield position moves the shield edge as much as 4 dB in the antenna beam patterns.

We detect a mean signal $Z_{\text{Earth}} = 42 \pm 15 \mu\text{K}$ when Earth is 1° below the shield, falling to $1 \pm 10 \mu\text{K}$ when Earth is 7° below the shield. This detection is somewhat larger than the upper limit $Z_{\text{Earth}} < 30 \mu\text{K}$ established from the 2 year data (Bennett et al. 1994). The difference results from Galactic emission in the 2 year data. The Galactic plane and bright extended features (Ophiuchus and Orion) cross the antenna beam center when Earth is below the shield. The resulting Galactic signal is brighter than Earth emission, even after binning in an Earth-based coordinate system. Simulations using the DMR scan pattern show that the binned Galactic emission has the opposite phase as Earth; i.e., Galaxy crosses the positive antenna when the Earth limb is at the azimuth of the negative antenna, and partly cancels Earth emission in the binned data. We account for this effect in the 4 year binned data by rejecting data with either antenna pointed at Galactic latitude $|b| < 15^\circ$ and subtracting a Galactic model from the remaining high-latitude data.

Table 9 shows the 95% CL upper limits to Earth emission when Earth is 1° below the shield. We reject data when




FIG. 11.—Amplitude of Earth emission in time-ordered data as a function of Earth limb elevation angle. The azimuthal variation from the beam patterns has been fitted to the Earth-binned data in a strip 1 pixel high. Earth emission falls rapidly as the Earth sets below the shield. We reject data for which the Earth limb is 1° below the shield or higher (3° for the 31A and 31B channels).

TABLE 9

95% CONFIDENCE UPPER LIMITS FROM EARTH EMISSION IN TIME-ORDERED DATA^a

Channel (1)	1° below Shield ^b (μK) (2)	7° below Shield ^c (μK) (3)
31A.....	179	26
31B.....	183	27
53A.....	85	8
53B.....	89	10
90A.....	136	10
90B.....	113	7

^a All values are in units of antenna temperature.

^b Direct fit to binned data 1° below the shield.

^c Values at -7° are taken from the values at -1° (col. [2]) and are then scaled to -7° using a diffraction model.

the Earth limb is 1° below the shield or higher (3° for the 31.5 GHz channels) but do not otherwise correct the data for Earth. The Earth signal decreases rapidly as Earth sets below the shield and falls below the threshold of detection 7° below the shield. Earth emission is not detectable for the majority of the data set: the Earth limb is 7° or more below the shield for 85% of the 4 year mission. We use the elevation dependence of the diffraction model to scale the upper limits at -1° to lower elevation angles and use this scaled model to map the Earth artifacts in the 4 year sky maps (eqs. [2] and [3]). Figure 12 shows the 95% confidence level limits for Earth emission in the 53B sky map. Since the upper limit is approximately twice the value of the detected signal, the scaled limits at lower elevations are conservative estimates of Earth emission. Earth emission contributes less than $1.8 \mu\text{K}$ to the pixel-pixel rms in the sky maps.

An alternate approach is to use the Earth-binned data as the model of Earth emission without reference to any a priori model. This avoids dependence on diffraction estimates but injects the pixel noise of the binned data into the mapping routine. Limits set by this model-independent technique, $\delta T_{\text{Earth}} < 3.5 \mu\text{K}$, are still small compared to the cosmic signal in the maps.

6.2. Moon

The Moon is the brightest source observed by DMR and is the only source visible in the time-ordered data. Away from beam center, its signal is rapidly attenuated by the antenna beam pattern. The beam pattern falls to a local minimum of -39 dB at 21° from beam center, at which point the Moon contributes approximately $150 \mu\text{K}$ to the time-ordered data. We reject any datum taken with an antenna pointed within 21° of the Moon and correct all remaining data for lunar emission using an a priori model based on lunar microwave emission properties and the measured DMR beam patterns (§ 4.1). Uncertainties in the lunar correction are dominated by $\sim 5\%$ systematic uncertainties in the brightness temperature of the Moon and by $\sim 3\%$ uncertainties in the antenna beam pattern. Residual uncertainties, after correction, are less than $9 \mu\text{K}$ in the time-ordered data and less than $0.3 \mu\text{K}$ in the maps.

6.3. Sun

A reflective shield protects the radiometers from the Sun, which never directly illuminates the aperture plane. A




FIG. 12.—Full sky map of 95% CL upper limits to Earth emission in the 53B channel (Mollweide projection in Galactic coordinates)

simple model of solar emission diffracted over the shield yields a limit $Z_{\text{Sun}} < 2\mu\text{K}$ in the time-ordered data (95% CL). We test for solar emission by binning the data by the solar position relative to the spacecraft axes, similar to the Earth binning in § 6.1. We find no evidence for solar emission in the time-ordered data. Solar effects are likely to be dominated by the heating of thermally sensitive components and not by the direct microwave emission. All solar effects (emission and thermal) are spin modulated and are subsumed in the spin-binned limits in Table 8 (§ 5.3).

6.4. Planets

We correct the time-ordered data for emission from Jupiter, Saturn, and Mars when those planets are within 15° of an antenna beam center. Uncertainties in the applied corrections are dominated by $\sim 10\%$ uncertainties in the brightness temperature of each planet (Kogut et al. 1992). Residual artifacts in the maps, after correction, are less than $0.4\mu\text{K}$. We test for planetary emission by mapping the time-ordered data in a coordinate system centered on the planet Jupiter. Accounting for the change in apparent diameter averaged over the 4 year mission, Jupiter should appear in such a map as an unresolved source with peak amplitude $200\mu\text{K}$. We fit the Jupiter-centered maps to a 7° FWHM Gaussian profile and recover amplitudes $209 \pm 139\mu\text{K}$, $195 \pm 40\mu\text{K}$, and $183 \pm 55\mu\text{K}$ antenna temperature at 31.5, 53, and 90 GHz, respectively.

6.5. Radio-Frequency Interference

Radio-frequency interference (RFI) from ground-based radars, if sufficiently strong, will appear as spikes in the time-ordered data. We have binned the flagged spikes by subsatellite point and find no correlation with position. We test for RFI from geostationary satellites by mapping the time-ordered data in a coordinate system focused on the geostationary satellite belt, including the effects of parallax.

Satellite RFI should appear as an unresolved source on the equator of this coordinate system. We find no such sources to limit $Z_{\text{RFI}} < 7\mu\text{K}$ in the time-ordered data and $\delta T_{\text{RFI}} < 0.02\mu\text{K}$ in the sky maps.

7. MISCELLANEOUS EFFECTS

Kogut et al. (1992) place stringent limits on a large number of potential systematic artifacts for the first year of DMR data. We have repeated these analyses for the 4 year data and include their effect in the combined limits to all systematic artifacts; however, we will not discuss each item separately. These effects include the solution convergence of the sparse matrix algorithm, pixel-pixel independence, discrete pixelization, nonuniform sky coverage, cross-talk with the DIRBE and FIRAS instruments, emission from the Sun-Earth shield, zodiacal dust emission, and cosmic-ray hits. Some of these effects are modulated at the spin or orbit periods and are absorbed in the limits in Table 8.

7.1. Attitude Solution

DMR uses the *COBE* fine aspect attitude solutions based on DIRBE stellar observations. For the first 45 months of the mission, residuals between the attitude solution and known stellar positions are less than $2'$ (68% CL). Fourier analysis of the residuals shows no periodic systematic uncertainties at either the spin or orbit periods; the frequency spectrum is close to white noise. A gyroscope failure on 1993 October 2 led to a wobble in spacecraft azimuth of amplitude $\sim 13'$, which is not included in the attitude solutions. We have simulated the effect of such a wobble in the 4 year DMR sky maps. Attitude artifacts in the maps are less than $0.2\mu\text{K}$ in the pixel-pixel rms.

7.2. Antenna Direction Vectors

The pointing of the 10 DMR antennas relative to the spacecraft body was measured prior to launch. We correct

TABLE 10
LOCK-IN AMPLIFIER MEMORY

Channel	Lock-in Memory Amplitude (percent of signal)
31A.....	3.220 ± 0.003
31B.....	3.146 ± 0.004
53A.....	3.203 ± 0.003
53B.....	3.172 ± 0.003
90A.....	3.110 ± 0.003
90B.....	3.139 ± 0.004

the pointing vectors of both 53B antennas for a 0.25 s telemetry timing offset (Kogut et al. 1992; Bennett et al. 1992a). Observations of the Moon provide a cross-check on the in-flight pointing of the antennas relative to the *COBE* spacecraft. We find no offsets larger than 16', well within our ability to model the centroid of the brightness distribution across the lunar disk. We have made sky maps using the lunar-derived antenna vectors and compared them to the mission maps made with the preflight antenna vectors. Systematic artifacts in the maps related to antenna pointing vectors are less than 1.4 μ K.

7.3. Lock-in Amplifier Memory

We correct the time-ordered data for the small "memory" of the previous datum caused by the low-pass filter on the lock-in amplifier input,

$$D(t) = D(t_i) - \alpha D(t_{i-1}),$$

where the coefficient $\alpha \approx 0.032$ (Kogut et al. 1992). In the absence of this correction, the lock-in memory creates a positive correlation between neighboring pixels in the sky maps. We compute α for each channel from the autocorrelation of the time-ordered data after correcting the calibrated

data for signals from the magnetic susceptibility, eclipse effect, lunar and planetary emission, and the CMB and Earth Doppler dipoles. Table 10 lists the resulting memory coefficients. We simulate the effect of small uncertainties in the applied coefficients. Artifacts in the maps resulting from uncertainties in the memory correction are negligible (less than 0.05 μ K).

8. DISCUSSION

We use equations (2) and (3) to make a sky map of each effect before correction and its 95% confidence uncertainty after our best correction (if any) is applied. Note that the temperatures in these maps are highly correlated from pixel to pixel: the spatial pattern is fixed, so that the residual (if any) across the DMR sky maps will be given by scaling the entire upper limit map in the range $[-1, 1]$.

We expand each systematic error map in spherical harmonics,

$$T(\theta, \phi) = \sum_{\ell m} a_{\ell m} Y_{\ell m}(\theta, \phi)$$

and calculate the multipole amplitudes

$$(\Delta T_{\ell})^2 = \frac{1}{4\pi} \sum_m |a_{\ell m}|^2$$

for each map. Tables 11, 12, 13, 14, 15, and 16 present the rms, peak-peak amplitude, and multipole amplitudes ΔT_{ℓ} for each systematic effect. The uncertainties in the potential systematic artifacts are largely independent; we adopt the quadrature sum of each column as the upper limit for the combined effects of all systematics. Note that this is not equivalent to adding the individual maps and then deriving ΔT_{ℓ} . Tables 11, 12, 13, 14, 15, and 16 list only the largest uncertainties; the row labeled "Other" contains the quadrature sum for all other systematics. This entry includes

TABLE 11
SYSTEMATIC EFFECTS FOR CHANNEL 31A^a

Effect	P-P ^b (μ K)	rms ^c (μ K)	ΔT_1 (μ K)	ΔT_2 (μ K)	ΔT_3 (μ K)	ΔT_4 (μ K)	ΔT_5 (μ K)	ΔT_6 (μ K)	ΔT_7 (μ K)	ΔT_8 (μ K)
Channel 31A before Correction										
β_X	6.7	0.4	0.2	0.1	0.1	0.0	0.1	0.0	0.0	0.0
β_R	31.9	6.5	1.1	6.1	0.5	1.6	0.7	1.0	0.2	0.5
β_T	17.4	2.9	18.4	1.6	1.9	0.7	0.9	0.3	0.1	0.2
Earth.....	7.0	1.2	1.5	0.9	0.4	0.4	0.3	0.2	0.0	0.1
Moon.....	7.9	1.6	0.0	1.2	0.0	0.4	0.1	0.3	0.0	0.5
Doppler.....	85.6	13.4	65.3	5.6	10.4	1.8	3.5	1.4	0.8	0.7
Spin.....	11.1	1.5	5.3	0.6	1.0	0.4	0.4	0.3	0.0	0.2
Other.....	81.3	9.1	2.3	0.8	0.4	0.5	0.4	0.5	0.4	0.5
Total ^d	124.7	17.9	68.2	8.6	10.6	2.7	3.7	1.8	0.9	1.2
Channel 31A after Correction ^e										
β_X	1.6	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
β_R	10.3	2.1	0.4	2.0	0.2	0.5	0.2	0.3	0.1	0.2
β_T	12.1	2.0	12.8	1.1	1.3	0.5	0.6	0.2	0.1	0.1
Earth.....	7.6	1.3	1.6	1.0	0.4	0.4	0.3	0.2	0.0	0.2
Moon.....	0.9	0.2	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.1
Doppler.....	4.3	0.7	3.3	0.3	0.5	0.1	0.2	0.1	0.0	0.0
Spin.....	23.2	3.1	11.1	1.3	2.0	0.8	0.8	0.7	0.1	0.4
Other.....	15.6	1.4	2.5	0.5	0.5	0.3	0.2	0.2	0.1	0.3
Total ^d	33.4	4.7	17.5	2.8	2.5	1.2	1.1	0.9	0.2	0.6

^a All results are in units of antenna temperature.

^b Peak-to-peak amplitude in the map after best-fit dipole is removed.

^c Pixel to pixel standard deviation after best-fit dipole is removed.

^d Quadrature sum of the individual effects in each column.

^e 95% confidence upper limits.

TABLE 12
SYSTEMATIC EFFECTS FOR CHANNEL 31B^a

Effect	P-P (μ K)	rms (μ K)	ΔT_1 (μ K)	ΔT_2 (μ K)	ΔT_3 (μ K)	ΔT_4 (μ K)	ΔT_5 (μ K)	ΔT_6 (μ K)	ΔT_7 (μ K)	ΔT_8 (μ K)
Channel 31B before Correction										
β_X	8.4	0.5	0.3	0.1	0.1	0.1	0.1	0.1	0.0	0.0
β_R	26.3	5.1	0.6	4.5	0.7	1.6	0.6	0.5	0.1	0.4
β_T	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Earth	11.5	1.6	0.5	1.5	0.3	0.3	0.3	0.2	0.0	0.1
Moon	8.5	1.6	0.1	1.3	0.0	0.4	0.0	0.3	0.0	0.5
Doppler	97.1	15.6	66.5	6.1	12.0	2.7	4.6	2.2	1.2	1.2
Spin	8.6	1.1	4.1	0.5	0.8	0.3	0.3	0.3	0.0	0.2
Other	95.9	10.9	0.7	0.7	0.3	0.5	0.3	0.5	0.4	0.5
Total	140.2	19.9	66.6	7.9	12.1	3.2	4.7	2.3	1.3	1.5
Channel 31B after Correction										
β_X	1.7	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
β_R	13.1	2.5	0.3	2.3	0.4	0.8	0.3	0.3	0.1	0.2
β_T	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Earth	13.0	1.8	0.6	1.6	0.4	0.3	0.3	0.2	0.1	0.1
Moon	0.9	0.2	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.1
Doppler	4.5	0.7	3.1	0.3	0.6	0.1	0.2	0.1	0.1	0.1
Spin	28.2	3.7	13.4	1.5	2.5	0.9	1.0	0.9	0.1	0.5
Other	22.1	2.6	2.8	0.5	1.1	0.4	0.5	0.3	0.2	0.3
Total	40.6	5.6	14.1	3.2	2.8	1.4	1.2	1.0	0.2	0.6

^a All results are in units of antenna temperature. Columns are the same as Table 11.

over a dozen individual effects, such as attitude and pointing errors, calibration drifts, lock-in amplifier memory, orbitally modulated effects, planetary emission, and artifacts from the map algorithm and pixelization. See Kogut et al. (1992) for a complete listing of these minor effects.

Upper limits on the combined effect of all systematics in the pixel-pixel rms of the 4 year DMR sky maps range from 5.4 μ K in the 31B channel (which contains only 21 months of “quiet” data) to 1.9 μ K in the 90B channel. Upper limits on the rms quadrupole amplitude ΔT_2 range from 3.8 to 1.1 μ K. The power drops rapidly at higher multipole moments ℓ .

We test for additional systematic artifacts in the sky maps by analyzing various “difference map” linear combinations in which celestial emission cancels, leaving instrument noise and (potentially) systematic effects. Examples are the (A – B)/2 differences between the A and B channels at each frequency, or similar maps made by differencing a single channel mapped in different time ranges (second year minus first year). Figure 13 shows the power spectrum of the 4 year (A – B)/2 difference maps compared to the expected instrument noise and the 95% confidence upper limits on combined systematic effects. The gray band represents the 95% CL range of power from instrument noise, determined by

TABLE 13
SYSTEMATIC EFFECTS FOR CHANNEL 53A^a

Effect	P-P (μ K)	rms (μ K)	ΔT_1 (μ K)	ΔT_2 (μ K)	ΔT_3 (μ K)	ΔT_4 (μ K)	ΔT_5 (μ K)	ΔT_6 (μ K)	ΔT_7 (μ K)	ΔT_8 (μ K)
Channel 53A before Correction										
β_X	32.9	2.2	5.1	0.4	0.8	0.4	0.4	0.3	0.1	0.1
β_R	6.3	1.0	0.6	0.6	0.2	0.7	0.2	0.2	0.0	0.1
β_T	101.0	15.7	92.8	7.6	11.9	3.0	2.7	1.7	0.5	1.0
Earth	6.6	0.7	0.5	0.6	0.2	0.2	0.1	0.1	0.0	0.1
Moon	7.1	1.5	0.0	1.2	0.0	0.4	0.0	0.3	0.0	0.5
Doppler	70.9	10.9	50.5	6.8	6.7	1.8	3.0	1.9	0.5	1.3
Spin	3.3	0.4	1.6	0.2	0.3	0.1	0.1	0.1	0.0	0.1
Other	37.3	3.7	1.0	1.0	0.4	0.6	0.3	0.5	0.4	0.5
Total	133.6	19.7	105.7	10.4	13.7	3.7	4.1	2.6	0.8	1.8
Channel 53A after Correction										
β_X	4.3	0.3	0.7	0.1	0.1	0.1	0.1	0.0	0.0	0.0
β_R	2.2	0.4	0.2	0.2	0.1	0.3	0.1	0.1	0.0	0.0
β_T	14.3	2.2	13.2	1.1	1.7	0.4	0.4	0.2	0.1	0.2
Earth	7.9	0.9	0.6	0.8	0.2	0.2	0.1	0.1	0.0	0.1
Moon	0.9	0.2	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.1
Doppler	1.2	0.2	0.7	0.1	0.1	0.0	0.1	0.0	0.0	0.0
Spin	7.6	1.0	3.6	0.4	0.7	0.3	0.3	0.2	0.0	0.1
Other	10.6	1.0	0.9	0.7	0.1	0.2	0.1	0.2	0.1	0.1
Total	21.5	2.8	13.7	1.6	1.8	0.6	0.5	0.4	0.2	0.3

^a All results are in units of antenna temperature. Columns are the same as Table 11.

TABLE 14
SYSTEMATIC EFFECTS FOR CHANNEL 53B^a

Effect	P-P (μK)	rms (μK)	ΔT_1 (μK)	ΔT_2 (μK)	ΔT_3 (μK)	ΔT_4 (μK)	ΔT_5 (μK)	ΔT_6 (μK)	ΔT_7 (μK)	ΔT_8 (μK)
Channel 53B before Correction										
β_X	1.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
β_R	31.5	5.3	2.8	3.2	1.2	3.6	0.9	0.8	0.2	0.5
β_T	22.1	3.4	20.3	1.7	2.6	0.7	0.6	0.4	0.1	0.2
Earth	6.6	0.8	0.6	0.7	0.2	0.2	0.1	0.1	0.0	0.1
Moon	7.2	1.6	0.0	1.2	0.0	0.4	0.0	0.2	0.0	0.5
Doppler	70.5	10.9	50.5	6.8	6.7	1.8	3.0	1.9	0.4	1.3
Spin	2.1	0.3	1.0	0.1	0.2	0.1	0.1	0.1	0.0	0.0
Other	41.9	4.3	2.7	1.3	0.4	0.6	0.3	0.5	0.4	0.5
Total	91.2	13.4	54.6	7.9	7.3	4.1	3.2	2.2	0.6	1.5
Channel 53B after Correction										
β_X	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
β_R	11.1	1.9	1.0	1.1	0.4	1.3	0.3	0.3	0.1	0.2
β_T	6.4	1.0	5.8	0.5	0.8	0.2	0.2	0.1	0.0	0.1
Earth	8.7	1.0	0.8	0.9	0.2	0.2	0.2	0.1	0.0	0.1
Moon	0.9	0.2	0.0	0.2	0.0	0.1	0.0	0.0	0.0	0.1
Doppler	1.2	0.2	0.7	0.1	0.1	0.0	0.1	0.0	0.0	0.0
Spin	7.1	0.9	3.4	0.4	0.6	0.2	0.2	0.2	0.0	0.1
Other	17.3	1.5	2.6	1.0	0.2	0.3	0.1	0.2	0.2	0.2
Total	24.3	2.9	7.4	1.9	1.1	1.4	0.5	0.4	0.2	0.3

^a All results are in units of antenna temperature. Columns are the same as Table 11.

Monte Carlo simulation using the observation pattern of each channel. The power spectra of the difference maps are in agreement with the expected instrument noise and show no statistically significant excursions. The upper limits to power from combined systematics are well below the noise limits: systematic artifacts do not limit analysis of the DMR sky maps.

9. CONCLUSIONS

We use in-flight data from the full 4 year mission to obtain estimates of potential systematic effects in the DMR time-ordered data and use the mapping software to create

maps of each effect and its associated 95% confidence level uncertainty. The largest effect known to exist in the time-ordered data is the instrument response to an external magnetic field. We correct for this magnetic susceptibility using a linear vector coupling between the radiometer orientation and Earth's magnetic field. The uncertainty in the correction is dominated by the instrument noise in the case of weak coupling and uncertainty in the local magnetic field for larger couplings.

With 4 years of data, we detect emission from Earth at the level $42 \pm 15 \mu\text{K}$ when Earth is 1° below the Sun/Earth shield. We reject data when the Earth is 1° below the shield

TABLE 15
SYSTEMATIC EFFECTS FOR CHANNEL 90A^a

Effect	P-P (μK)	rms (μK)	ΔT_1 (μK)	ΔT_2 (μK)	ΔT_3 (μK)	ΔT_4 (μK)	ΔT_5 (μK)	ΔT_6 (μK)	ΔT_7 (μK)	ΔT_8 (μK)
Channel 90A before Correction										
β_X	4.9	0.4	1.6	0.1	0.2	0.1	0.1	0.0	0.0	0.0
β_R	89.3	15.1	7.9	9.2	3.4	10.4	2.7	2.1	0.5	1.3
β_T	37.8	5.9	35.0	2.9	4.5	1.1	1.0	0.6	0.2	0.4
Earth	4.8	0.6	0.5	0.5	0.1	0.2	0.1	0.1	0.0	0.1
Moon	8.4	1.8	0.1	1.4	0.0	0.5	0.0	0.3	0.0	0.6
Doppler	62.3	9.6	44.3	5.9	5.9	1.6	2.7	1.6	0.4	1.1
Spin	4.9	0.6	2.3	0.3	0.4	0.2	0.2	0.2	0.0	0.1
Other	51.8	5.3	1.7	0.9	0.3	0.5	0.3	0.5	0.3	0.4
Total	126.9	19.7	57.1	11.4	8.2	10.6	3.9	2.8	0.8	2.0
Channel 90A after Correction										
β_X	1.0	0.1	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
β_R	12.3	2.1	1.1	1.3	0.5	1.4	0.4	0.3	0.1	0.2
β_T	10.3	1.6	9.5	0.8	1.2	0.3	0.3	0.2	0.1	0.1
Earth	9.4	1.2	1.0	1.0	0.2	0.3	0.2	0.2	0.1	0.1
Moon	1.4	0.3	0.0	0.2	0.0	0.1	0.0	0.1	0.0	0.1
Doppler	2.6	0.4	1.8	0.3	0.2	0.1	0.1	0.1	0.0	0.1
Spin	12.1	1.6	5.8	0.7	1.1	0.4	0.4	0.4	0.0	0.2
Other	15.4	1.0	1.6	0.7	0.1	0.2	0.1	0.2	0.1	0.2
Total	27.2	3.5	11.5	2.0	1.7	1.6	0.7	0.6	0.2	0.4

^a All results are in units of antenna temperature. Columns are the same as Table 11.




FIG. 13.—Power spectrum of the 4 year $(A-B)/2$ difference maps compared to the instrument noise and upper limits on the combined systematic uncertainties. Points represent the power spectrum of the $(A-B)/2$ map, while the light gray band represents the 95% CL range of power from Monte Carlo simulations of instrument noise. The dark gray band shows 95% confidence level upper limits to the quadrature sum of systematic effects.

or higher but do not otherwise correct for this emission. Emission from Earth agrees well with a simple diffraction model over the limited range of elevation angles for which we detect Earth. We use the elevation dependence of the diffraction model to scale the detection at -1° to lower elevation angles when deriving upper limits to Earth artifacts in the 4 year sky maps.

We correct the calibrated time-ordered data for the known systematic effects (Table 1) and bin the data by the spacecraft spin angle relative to the Sun. We find no evidence for additional systematic effects modulated at the spin period. Our ability to detect weak signals with this method is limited by the instrument noise; upper limits to spin-modulated artifacts in the maps determined from the spin-binned data are determined by the instrument noise and not by a detected signal.

We detect an orbitally modulated signal in all channels during the 2 month “eclipse season” surrounding the June

solstice, when the spacecraft flies through Earth’s shadow over the Antarctic each orbit. The detection of thermal gain variations during the same time range provides a plausible mechanism for this effect. We correct the time-ordered data using an empirical model based on variations in unregulated spacecraft temperature and voltage signals, which provide templates for the time variation less affected by telemetry digitization than the housekeeping signals on the amplifiers themselves. The model removes about $\frac{2}{3}$ of the signal. The residual is large enough in the 31A and 31B channels that the 4 year data would be adversely affected by including the corrected data with the residual signal. We do not use data during the eclipse season for the 31A and 31B channels.

Observations of the Moon and the Doppler dipoles from the motion of the satellite about Earth and Earth about the solar system barycenter demonstrate that the preflight calibration is accurate within its uncertainties. On-board noise

TABLE 16
SYSTEMATIC EFFECTS FOR CHANNEL 90B^a

Effect	P-P (μK)	rms (μK)	ΔT_1 (μK)	ΔT_2 (μK)	ΔT_3 (μK)	ΔT_4 (μK)	ΔT_5 (μK)	ΔT_6 (μK)	ΔT_7 (μK)	ΔT_8 (μK)
Channel 90B before Correction										
β_X	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
β_R	10.7	1.8	0.9	1.1	0.4	1.2	0.3	0.3	0.1	0.2
β_T	15.4	2.4	14.2	1.2	1.8	0.5	0.4	0.3	0.1	0.2
Earth	5.3	0.6	0.5	0.5	0.1	0.2	0.1	0.1	0.0	0.1
Moon	9.4	2.0	0.1	1.5	0.0	0.6	0.0	0.3	0.0	0.6
Doppler	62.7	9.6	44.3	5.9	5.9	1.6	2.7	1.6	0.5	1.2
Spin	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other	43.1	4.3	1.4	0.8	0.3	0.5	0.3	0.4	0.3	0.4
Total	79.1	11.1	46.5	6.4	6.2	2.2	2.7	1.8	0.6	1.4
Channel 90B after Correction										
β_X	0.5	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
β_R	2.9	0.5	0.3	0.3	0.1	0.3	0.1	0.1	0.0	0.0
β_T	6.7	1.0	6.2	0.5	0.8	0.2	0.2	0.1	0.0	0.1
Earth	7.8	0.9	0.7	0.8	0.2	0.2	0.1	0.1	0.0	0.1
Moon	1.5	0.3	0.0	0.3	0.0	0.1	0.0	0.1	0.0	0.1
Doppler	1.8	0.3	1.2	0.2	0.2	0.1	0.1	0.1	0.0	0.0
Spin	5.4	0.7	2.6	0.3	0.5	0.2	0.2	0.2	0.0	0.1
Other	11.2	0.8	1.3	0.5	0.2	0.2	0.1	0.1	0.1	0.1
Total	16.6	1.9	7.0	1.2	1.0	0.5	0.3	0.3	0.1	0.2

^a All results are in units of antenna temperature. Columns are the same as Table 11.

sources track the gain correctly within $0.2\% \text{ yr}^{-1}$. Artifacts from calibration errors are negligible. The $70 \mu\text{K}$ Doppler signal from the orbital motion of the spacecraft about Earth and the $200 \mu\text{K}$ emission from Jupiter serve as convenient test patterns in the appropriate specialized coordinate systems. DMR observes both signals with good signal-to-noise ratio.

We map the pattern of artifacts in the sky from the 95% confidence level uncertainty in each potential systematic effect and analyze these maps as though they were maps of

the CMB. The quadrature sum of the combined upper limits ranges from 5.4 to $1.9 \mu\text{K}$ in the pixel-pixel rms, and from 3.8 to $1.1 \mu\text{K}$ for the rms quadrupole amplitude. The power drops rapidly at higher multipole moments ℓ . The power spectra of the 4 year $(A - B)/2$ difference maps are consistent with the distribution of instrument noise in the maps and show no evidence for systematic artifacts. Upper limits to power from combined systematics in the DMR 4 year sky maps are well below the noise limits: systematic artifacts do not limit analysis of the DMR sky maps.

REFERENCES

- Barker, F. S., et al. 1986, EOS, 67, 523
 Bennett, C. L., et al. 1992a, ApJ, 391, 466
 ———. 1992b, ApJ, 396, L7
 ———. 1994, ApJ, 436, 423
 ———. 1996, ApJ, 464, L1
 Boggess, N. W., et al. 1992, ApJ, 397, 420
 Fixsen, D. J., Gales, J. M., Mather, J. C., Shafer, R. A., & Wright, E. L. 1996, ApJ, submitted
 Jackson, P. D., et al. 1992, in ASP Conf. Proc. 25. Astronomical Data Analysis Software and Systems. I, ed. W. Worrall, C. Biemesderfer, & J. Barnes (San Francisco: ASP), 382
 ———. 1996, in preparation
 Janssen, M. A., & Gulkis, S. 1992, in The Infrared and Submillimeter Sky after COBE, ed. M. Signore & C. Dupraz (Dordrecht: Kluwer), 391
 Lineveaver, C. H., et al. 1994, ApJ, 436, 452
 Keihm, S. J. 1982, Icarus, 53, 570
 Keihm, S. J., & Gary, B. L. 1979, in Proc. 10th Lun. Planet Sci. Conf., Vol. 3 (New York: Pergamon), 2311
 Keihm, S. J. & Langseth, M. G. 1975, Icarus, 24, 211
 Kogut, A., et al. 1992, ApJ, 401, 1
 Kogut, A., Banday, A. J., Bennett, C. L., Górski, K. M., Hinshaw, G., & Reach, W. T. 1996a, ApJ, 460, 1
 Kogut, A., Banday, A. J., Bennett, C. L., Górski, K. M., Hinshaw, G., & Smoot, G. F. 1996b, ApJ, 464, L5
 Peebles, P. J. E., & Wilkinson, D. T. 1968, Phys. Rev., 174, 2168
 Toral, M. A., Ratliff, R. B., Lecha, M. C., Maruschak, J. G., Bennett, C. L., & Smoot, G. F. 1990, IEEE Trans. Ant. Prop., 37, 171
 Smoot, G. F., et al. 1990, ApJ, 360, 685
 ———. ApJ, 391, L1
 Torres, S., et al. 1989, in Data Analysis in Astronomy III, ed. V. Di Gesu, L. Scarsi, & M. C. Maccarone (New York: Plenum), 319
 Wright, E. L., et al. 1994, ApJ, 420, 1