

GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

INTRODUCTION.

I. *Personal Establishment and Arrangement.*

At the beginning of the year 1908 the Established Staff in the Astronomical Department of the Observatory consisted of the following persons :—

Chief Assistants,—Philip Herbert Cowell, M.A., F.R.S., formerly Fellow of Trinity College, Cambridge; Arthur Stanley Eddington, M.A., M.Sc., Fellow of Trinity College, Cambridge.

Assistants,—Edward Walter Maunder; Thomas Lewis; William Grasett Thackeray; Henry Park Hollis, B.A.; Andrew Claude de la Cherois Crommelin, B.A.

Clerical Assistant,—Henry Outhwaite.

Established Computer (Higher Grade),—Charles Davidson.

Established Computers,—William Bowyer; Herbert Henry Furner; William Moody Witchell, B.Sc.; John Storey; Philibert Melotte; William Stevens; William Wood Burkett; Joseph Edward Evans; Richard Thomas Cullen.

GREENWICH OBSERVATIONS, 1908.

a

ii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Mr. J. Storey resigned in May 1908, having been appointed Assistant in Edinburgh Observatory.

In addition to the Assistants and Established Computers, twenty-two Supernumerary Computers were employed in the Astronomical Department.

Mr. W. W. Bryant, Superintendent of the Magnetic and Meteorological Branch, and Mr. D. J. R. Edney, Established Computer in that branch, have also taken part in the Astronomical Observations.

The duties of the establishment are distributed in the following manner:—

The Chief Assistants, in the absence of the Astronomer Royal, are empowered to act in all respects as his representatives, and to conduct confidential as well as routine business. In the ordinary transactions, the Chief Assistants superintend the calculations generally; and observe occasionally with any of the instruments. The Chief Assistants are charged with the adoption of clock-rates, instrumental zeros, and instrumental errors.

Mr. Thackeray is charged with the deduction of the results of the meridian observations, and with the general superintendence of the Computers of the Astronomical Reductions Branch.

Mr. Crommelin has charge of the altazimuth and of the Sheepshanks equatorial, and of the reduction of the observations made with them.

The charge of the sidereal and solar clocks, the distribution of time-signals, the dropping of the time-signal ball at 13^h, the management of the galvanic apparatus and connections, the receiving, rating, and issuing of chronometers, and all the computations relating to the time-department, are confided to Mr. Lewis. Mr. Lewis also has charge of the 28-inch equatorial.

Mr. Hollis superintends the measurement and reduction of the photographs taken for the Astrographic Chart and Catalogue.

Mr. Davidson has charge of the measurement of the photographs taken with the Thompson Equatorial and the reduction of the measures.

The measurement of the solar photographs and the superintendence of the reductions connected with them are entrusted to Mr. Maunder.

Mr. Outhwaite has charge of the money-accounts, the stores and stationery, and distribution of the publications of the Observatory; and undertakes part of the miscellaneous correspondence. The care of the library is also entrusted to him.

The course of observations and the succession of observers are arranged every Monday by one of the Chief Assistants and sanctioned by the Astronomer Royal. It is established as a rule, to be adhered to as closely as circumstances permit, that no Assistant be occupied on two successive days with astronomical observations. In general, the Assistant who makes the observations with the transit-circle is charged with all the observations that may occur from 19^h mean solar time (7 a.m.) to the next 14^h. During the latter part of the lunation, when the moon passes the meridian in the early morning between 14^h and 19^h, it is found necessary to appoint separate observers for the morning and the evening.

The observations with the transit-circle were generally made by Mr. Witchell, Mr. Storey, Mr. Evans, and Mr. Cullen. When the moon passes the meridian in the early morning, or in the absence of the regular observers, one of the supernumerary Computers is charged with the observations.

Observations of occultations were generally made by the observer on duty with the transit-circle.

Observations with the reflex zenith-tube were made by Mr. Witchell, Mr. Storey, Mr. Evans and Mr. Cullen.

Observations with the 28-inch refractor were made by Mr. Lewis, Mr. Bryant, Mr. Bowyer and Mr. Furner.

Observations with the Altazimuth were made by Mr. Crommelin, Mr. Furner, or by one of the Computers.

Photographs with the astrographic equatorial were generally taken by Mr. Hollis, or Mr. Stevens.

Photographs with the Thompson Equatorial (26-inch refractor and 30-inch reflector) were generally taken by Mr. Davidson, Mr. Edney and Mr. Melotte.

iv INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Photographs of the Sun with the Dallmeyer photoheliograph were taken by Mr. Maunder, or by one of the Computers under his direction.

The following are the signatures of those persons who have made observations in 1908 :—

W. H. M. Christie	WC	D. J. R. Edney	- DE	J. Shepperd	- S
P. H. Cowell	- C	W. Bowyer	- WB	B. D. Evans	- BE
A. S. Eddington	- SE	H. H. Furner	- HF	H. Acton	- HA
E. W. Maunder	- M	W. M. Witchell	- W	A. Witney	- AW
T. Lewis	- L	J. Storey	- JS	G. Cody	- GC
W. G. Thackeray	- T	W. Stevens	- WS	H. E. Green	- G
H. P. Hollis	- H	P. Melotte	- PM	F. Jeffries	- J
A. C. D. Crommelin	AC	J. E. Evans	- E		
W. W. Bryant	- B	R. T. Cullen	- RC		
C. Davidson	- CD	S. Daniels	- SD		

II. *Instruments.*

The principal instruments used by Halley, Bradley, Bliss, and Maskelyne are still preserved in the Royal Observatory : namely, Halley's transit, with pivots unequally distant from the telescope ; Bradley's transit ; Bradley's small equatorial ; the zenith-sector of Bradley and Maskelyne ; and the two mural-quadrants, which till 1887 November 17 were both mounted on their pier, the quadrant on the western side of the pier being included in the smaller fire-proof room of the Observatory. On 1887 November 18, in the course of operations for the extension of the computing rooms, the quadrant on the western side of the pier was removed to the Transit-circle room, that on the eastern side being now included (without having been disturbed) in the lower computing room ; and an extension of the pier was built to carry an equatorial. The ancient instruments first mentioned, together with the telescopes for investigating the parallaxes of α Aquilæ and α Cygni, formerly attached respectively to the west side of the pier of Troughton's mural-circle, and to the Quadrant pier, are now, with proper labels, suspended on brackets on the western wall of the Transit-circle room.

Troughton's transit instrument and mural-circle, used to the end of the year 1850, were dismantled in the year 1851, and are suspended, the one on the west and the other on the east wall of the Transit-circle room. The object-glass of the transit instrument is now inserted in the reflex zenith-tube.

TRANSIT-CIRCLE.

v

The 25-foot zenith-sector was dismantled, and its tube was divided (at its joints) into several parts, in May 1848; and the separate portions of the instrument are stored away in frames constructed to hold them. The object-glass was removed from the tube in 1850, to be used for the collimator of the old altazimuth.

Jones's Cape circle was transferred in November 1851 to the Observatory of Queen's College, Belfast.

The Instruments now in use are the following :—

The Transit-circle, constructed by Messrs. Ransomes and May, as engineers, and by the late Mr. William Simms, as optician, erected in the year 1850, and brought into use at the beginning of 1851.

The room in which this instrument is mounted occupies the site of the old circle-room, but is extended to the south, so that its entire length is 36 feet. The ridge of the roof is in the north-and-south direction. The opening in the roof, along the ridge, is 3 feet wide, and is covered by four shutters. The vertical openings in the north and south walls are also 3 feet wide, and each is covered by a single shutter. Any one of the shutters can be opened without disturbing the others.

A detailed description of the instrument, illustrated by plates, is given in Appendix I. of the volume for 1852: a reprint of which, with some modifications, is attached to the volume for 1867. The following particulars may be given here. The centre of the instrument is about $5\frac{1}{2}$ feet south and 19 feet east of the old transit instrument. The focal length of its object-glass is 11 feet 7 inches, and the clear aperture 8.1 inches. It was repolished by Messrs. Troughton and Simms in 1891, and again in 1906. The full aperture of eight inches has been used in all observations, including those of the Sun. The power of the eyepiece ordinarily employed is 195, and of that used for observations of the Sun 180. The axis of the instrument is of cast-iron, in two similar pieces, the length between the extremities of the pivots being 6 feet. The mould of the pivot was made of iron, for the purpose of hardening it by the process technically called *chilling*; the moulds for the other parts were sand. The two halves are connected by bolts through flanges at the junction-plane in the middle of the cube. The two portions of the telescope-tube (also of cast-iron, with the exception of the object-glass cell and the eyepiece-work) are bolted on the central cube. The pivots are of "chilled" iron and their diameter is 6 inches, the bearing of each being upon two portions of a concave cylinder, forming the Y. The Y's

are firmly screwed down to the massive piers between which the instrument is mounted.

For examination of the form of the pivots, each is perforated; within the hollow of the eastern pivot there is fixed a plate of metal perforated with a hole about 0.01 inch in diameter, behind which a light can be placed for illumination; at the distance of 6 inches from this hole is a lens of 1 inch focal length, producing an image of the hole about 0.002 inch diameter, which in fact is the real mark for collimation; and in the hollow of the western pivot there is fixed an object-glass at a distance from that image equal to its focal length. This combination forms a reversed telescope revolving with the instrument. It is viewed by a telescope of 7 feet focal length, which, when required, is placed on Y's, one of them planted in the opening of the western pier, and the other in a hole made for that purpose in the western wall of the room. During 1905 October 31 to November 18, observations were made for the determination of the relative errors of the form of the pivots of the transit-circle. The process employed is fully explained in *Appendix I. of Greenwich Observations* for 1852. The following table contains the residual errors, which define the apparent changes of position of a certain line in the material axis of the instrument:—

N. P. D. Pointer Reading.	Horizontal Error, Microme- ter Head to Left.	Vertical Error, Microme- ter Head Below.	N. P. D. Pointer Reading.	Horizontal Error, Microme- ter Head to Left.	Vertical Error, Microme- ter Head Below.	N. P. D. Pointer Reading.	Horizontal Error, Microme- ter Head to Left.	Vertical Error, Microme- ter Head Below.	N. P. D. Pointer Reading.	Horizontal Error, Microme- ter Head to Left.	Vertical Error, Microme- ter Head Below.
0	"	"	0	"	"	0	"	"	0	"	"
5	-0.07	-0.10	95	-0.10	-0.13	185	+0.05	+0.05	275	-0.13	+0.31
15	-0.05	-0.08	105	-0.16	+0.03	195	+0.13	0.00	285	-0.18	+0.39
25	-0.05	0.00	115	-0.05	+0.08	205	+0.18	+0.05	295	-0.18	+0.28
35	+0.03	-0.10	125	+0.08	0.00	215	+0.13	-0.03	305	-0.13	+0.18
45	+0.10	-0.13	135	+0.03	0.00	225	+0.13	+0.05	315	-0.08	-0.05
55	+0.03	-0.21	145	+0.16	+0.08	235	+0.08	+0.13	325	-0.03	-0.08
65	0.00	-0.23	155	+0.18	+0.03	245	0.00	+0.16	335	-0.05	-0.16
75	-0.10	-0.23	165	+0.16	-0.05	255	+0.03	+0.13	345	0.00	-0.08
85	-0.08	-0.18	175	+0.13	-0.05	265	0.00	+0.21	355	-0.03	-0.21

No correction has been applied on account of error in the form of the pivots.

The wire frame contained originally (besides the horizontal wire, to be noticed

shortly) seven vertical wires, adapted to observations of transits by eye-and-ear; to which six were added in the spring of 1854, for more convenient use in the observations of transits by galvanic contact. In 1891 October the galvanic wire system was rearranged: there are now ten wires so placed that the mean of the ten coincides nearly with the middle wire, which is also the middle wire of the eye-and-ear system. The intervals between the wires are given in the *Transit-Circle Tables* at the end of this *Introduction*.

The whole frame and whole system of vertical wires are moved horizontally by a micrometer-screw, whose graduated head is locked up in a small box attached to the eyepiece, to prevent inadvertent disturbance of the micrometer after it has been set to the reading which is adopted for the line of collimation. The micrometer-head is on the eastern side of the eyepiece, and the readings increase as the wire is moved towards the micrometer-head. In 1891 October a new screw was fitted to the micrometer by Messrs. Troughton and Simms, one revolution of which is nearly equal to the distance between two close wires, viz. 37".

From 1908 April 27 the field of view has been illuminated by the light from a small electric lamp carried near the end of the telescope tube beyond the object-glass. The light, after passing through a condensing lens, falls in a parallel beam on a disk of finely ground opal glass, attached at the centre of the object-glass and inclined at 45° to it, and is scattered down the axis of the telescope. The intensity of the illumination is controlled by a rheostat. The annular illumination formerly in use is still available, and observations are sometimes made by it for comparison of the two methods.

For determining the error of collimation, two horizontal telescopes or collimators of about 6 feet 10 inches focal length and 7 inches aperture, with their axes in the same horizontal line through the centre of the instrument and their object-glasses turned towards it, are mounted on Y's carried by massive brick piers, one on the north and the other on the south side of the transit-circle. Each collimator is furnished with wires in its principal focus, to be used as collimating marks; and each may be used as presenting a distinct mark for the other, or for the transit-circle telescope. For adjustment of the collimators accurately on each other, the plates carrying the wires are moveable by micrometer-screws: that of the south collimator in altitude only, and that of the north collimator in azimuth only. In order to view either collimator by the transit-circle telescope, it is only necessary to direct the transit-circle telescope towards that collimator; but in order to view one collimator by the other, it is necessary to place the transit telescope vertical, and to uncover the perfora-

tions in the opposite sides of its central cube, which permit one collimator to view the other through eight holes of sector-form. The opposite sides of the central cube were pierced with these perforations in August and September 1865. This arrangement was brought into use on 1866 December 17, when the collimators of 7 inches aperture were mounted in place of those of 4 inches aperture, used up to that time. To obtain a somewhat more perfect view of one collimator by the other, it is necessary to raise the transit-circle so far that there shall be no impediment to a direct view, by means of a mechanical apparatus provided for the purpose.

Since the end of 1907 October the system of wires in each collimator has been a pair of close vertical wires and a horizontal wire. With this arrangement the adjustment of the collimators for use in determining the line of collimation is as follows,—the transit telescope is placed in the proper position and, the south collimator being illuminated by reflected skylight, the eye is applied to the eyepiece of the north collimator and the systems of wires in both are seen. The separation of the vertical wires in the north collimator is less than that of the wires in the south collimator, and the reading for coincidence of collimators is determined by moving the north wires until they are seen symmetrically between the south wires. The operation is repeated six times, the micrometer being read each time, and the micrometer is left in the position indicated by the mean. The daily observation for collimation of the transit telescope is then made by bringing, by means of its R.A. micrometer, the central vertical wire several times in succession midway between the vertical wires of the south collimator and several times midway between the vertical wires of the north collimator, and reading the micrometer for each observation. The mean of the readings for the north and south collimators gives the reading for the position of the line of collimation.

In June 1882 a new form of mounting the collimators was adopted, in order to obtain a greater range of observations of stars by reflexion. In the new arrangement, the two piers having been cut down, the collimators are mounted on upright cast-iron arms which turn about centres below, thus allowing them to be swung on one side when not in use. An alteration was also made in the mounting of the mercury trough, by which it is raised about a foot. It is thus found practicable to observe stars by reflexion (with the full aperture of the object-glass) from Z.D. 20° to Z.D. $67\frac{1}{4}^{\circ}$, an increase of range of nearly 30° on each side of the zenith. To test the stability of the collimators during each determination of collimation-error, it is the usual practice, since the alteration, to observe the coincidence of the corresponding wires of the two collimators immediately after as well as before

the observation of the two collimators with the transit-circle. When the two sets of micrometer-readings differ, a correction corresponding to the half-difference is applied to the concluded reading for line of collimation.

Since 1884 June a reversion-prism made by Messrs. Troughton and Simms has been used from time to time in observations with the collimators as well as with the transit-circle to reverse the apparent direction of measurement or of motion, a movement towards the left (as in transits of south stars) being converted into a movement towards the right, or downwards or upwards, according to the position of the plane of reflexion of the reversion-prism.


The error of level is ascertained by the use of a Bohnenberger's eyepiece, with three lenses, and a transparent glass reflector at an angle of 45° with the axis of the eyepiece, which is placed between the lowest lens and the wires of the telescope. A beam of light being thrown horizontally upon the reflector, and a trough of quicksilver (at first of iron, but on 1889 September 26 furnished with a bottom of amalgamated copper, with very beneficial results as regards steadiness of the images; and on 1890 June 5 replaced by an amalgamated copper trough) being placed below the object-glass of the telescope, the image of the central wire is seen by reflexion at the same time as the wire itself; by means of the micrometer-screw, the images are made to coincide, or to touch alternately on the two sides; the mean of the micrometer-readings is taken, and the difference between the mean reading and the reading corresponding to the line of collimation is the error of level.

The graduated vertical circle for zenith-distance observations is fixed on the cylindrical base of the axis-cone on the west side of the central-cube. It is shielded from the Sun's rays by the steps which are used by the observers for ascending to the upper part of the pier. It is of cast-iron, 6 feet in diameter, and has two sets of divisions: one set, on its western side, cut upon a band of silver, which is let into the internal surface of a very flat cone, is accurately divided to five-minute spaces and is read by the microscopes; the other set, on its eastern side, is roughly divided by points to every $5'$, and is intended for setting the telescope to any object by means of two pointers reading respectively north polar distances and zenith distances. The tubes of the reading microscopes are inclined perforations through the western pier (not furnished with any metallic tube), pointing to the graduations on the flat internal conical surface of the graduated circle. Their eyepieces are all carried by one massive brass plate at the back of the pier, and are arranged in a circle, whose centre is 5 feet 2 inches above the floor, and whose diameter is about 21 inches. Their object-glasses are separately attached to the inner or eastern side of the pier,

GREENWICH OBSERVATIONS, 1908.

b

and are arranged in a circle of about 5 feet in diameter. Each of the microscope perforations through the pier is accompanied with a perforation for illumination: these illumination-perforations all diverge from one central electric light near the western face of the pier. Each is furnished with a lens, $3\frac{1}{2}$ inches in diameter, by adjustment of which the light from the electric lamp, after specular reflexion from the graduated surface, is thrown up through the microscope-perforations to the microscope eyepieces. A lining of double tin plates, inserted in the central opening of the pier, screens the stone pier, the brass plate, the micrometers, and the eyepieces. Provision is made for ten micrometer-microscopes and a pointer-microscope, arranged in the following order:—



L is the electric light for illumination. P is the pointer-microscope furnished with an eyepiece of low power for reading the integral graduations; A, C, E, B, D, F are the six micrometer-microscopes at intervals of 60° , used in ordinary observations; a and b are supplementary micrometer-microscopes at 20° distance from A and B respectively, and α and β similar microscopes at 25° distance from A and B; these supplementary microscopes are used for determinations of the errors of graduation of the circle, or occasionally in observations of zenith distance during repair of the six ordinary microscopes.

The errors of division of the circle were investigated for every 5° in 1851, 1856, 1871, and 1898, two independent determinations having been made in the last year. The mean of the determinations made in 1851 and 1856 was adopted for use up to the end of 1896, and in the sections of Zenith Distances and Star Ledgers in 1897. For 1898 and subsequent years, and in the Star Catalogue and Planetary Results for 1897 the mean of the four determinations made in 1856, 1871, and 1898 respectively was adopted, giving a result in which each of the 5° divisions is equally well determined. The determination made in 1851 was rejected as being discordant. The errors of the single degrees were determined in 1851, 1856, and 1898, the mean of the first two being used to the end of 1896 and in the earlier sections in 1897, and the mean

of the three for 1898 and subsequent years, and in the final sections for 1897. Full information on the subject is given in a paper by Mr. Dyson and Mr. Thackeray in the *Memoirs of the Royal Astronomical Society*, vol. liii., and in the *Transit-Circle Tables* at the end of the *Introduction* for 1897.

At the end of 1875 new micrometer-screws were applied by Mr. Simms to the six ordinary micrometers A, B, C, D, E, F. In March 1878 the micrometers A, C, and F were reversed (their heads being at the same time re-figured to read in the opposite direction), so that any effect of wear in the micrometer-screws might be eliminated for each pair of microscopes. At the end of 1885 new screws made of steel instead of gun-metal were applied by Mr. Simms to the six ordinary micrometers A, B, C, D, E, F; and were brought into use on 1886 January 1. Observations for determining the errors of these screws are given in the *Transit-Circle Tables* at the end of the *Introduction* for 1886 (p. cxxxv). Similar observations were made in 1893, 1896, and 1906. The observations are given in the *Transit-Circle Tables* at the end of the *Introductions* for 1894, 1896, and 1906. They show that the wear of the screws is small, and its effect almost entirely eliminated by the reversal of three of the screws.

There are two clamps attached to the eastern pier, at the same height as the centre of the circle (one on the north side, the other on the south side), which can take hold of the clamping circle of the instrument; but they have no slow motion. The end of the telescope contains only one horizontal wire, moveable by a micrometer, with which all observations of zenith-distance are made. For reducing every observation, therefore, it is necessary to combine the value of the reading of the telescope-micrometer with that of the mean of readings of the microscope-micrometers. When the telescope points vertically upwards, the head of the zenith-distance-micrometer is on the north side; and the zenith-distance-micrometer-readings increase as the wire is moved towards the head.

A new screw was applied to the micrometer by Messrs. Troughton and Simms in 1906 July. Its pitch is the same as that of the right ascension micrometer. Observations for determination of the value of one revolution of the screw and of its errors are given in the *Transit-Circle Tables* at the end of the *Introduction* for 1906. The observations show that the new screw is sensibly uniform, and no corrections for its errors are applied.

In 1873 an apparatus was attached to the zenith-distance-micrometer for mechanical registration of its readings. In this arrangement, punctures corresponding to each

bisection of an object, in its passage across the field, are made on a strip of paper, fixed on a light drum immediately above the divided head of the micrometer, and turning with it. To distinguish the several bisections, the pricker by which the punctures are made, is, after each puncture, moved through a definite space in the direction of the axis of the drum, by turning a screw, which carries it, through a quarter turn. After a set of bisections the punctures are successively brought up to a straight edge, $0\cdot050$ from the pricker, and the micrometer-head is read off; each is then marked with a pencil to distinguish it from any which may be made afterwards. By this arrangement several bisections of an object can be made without the observer having to move his eye from the telescope, and a permanent record is obtained, giving facilities for the correction of mistakes.

The reading (consisting of the combination of zenith-distance-micrometer-reading and mean of microscope-readings), corresponding to the nadir-position of the telescope, is found by the use of the Bohnenberger's eyepiece and the trough of mercury; the direct and reflected images of the horizontal wire being made to coincide or to touch alternately on the two sides, and the mean of the readings of the micrometer being taken.

Other explanations necessary for the understanding of the process of reduction of the observations will be given under the proper heads. In this place it may be useful to explain briefly the methods employed in making the various classes of observations which are required.

For star-observing generally, the telescope is directed approximately towards the object by the indications of the N.P.D. pointer, and, the telescope being moved by hand till the star is brought near the horizontal micrometer-wire at or near the reading of 20^{rev} , the clamping circle is fastened. The transit of the star is then observed over the vertical wires in the usual way. For stars within 15° of the pole, it has been the practice since 1899 to take transits over one of the wires at different settings of the transit micrometer, as stated in the footnotes, the slide carrying the wires being moved by means of the micrometer-screw and the readings noted by the observer. For stars within 3° of the pole (except the nine azimuth stars), the transits are observed by eye-and-ear, the chronograph being used for all other stars, including the nine azimuth stars. During the transit, an observation of zenith distance is made by means of the zenith-distance-micrometer, the horizontal wire being made to bisect the star near the passage over one or more of the vertical wires, and punctures being made on the drum of the micrometer. The corresponding readings of the micrometer are then taken, and the vertical wires to which they correspond noted and the pointer and

the six microscope-micrometers read. The practice of observing very faint stars in a perfectly dark field with the wires illuminated has been discontinued.

For observations of stars by reflexion, the telescope is placed, some minutes previously to the transit of the star, in the direction which is proper for viewing the reflected image of the star, and is clamped; the mercury-trough is placed in the requisite position, and the microscope-micrometers are read. The observer then ascends to the eyepiece of the telescope, and by means of the micrometer-screw which carries the horizontal wire, bisects the reflected image of the star at the passage of the first and second vertical wires (if the star has large polar distance), or of two or three of the galvanic wires (if the star is circumpolar), and reads the revolution-counter of the zenith-distance-micrometer. He then descends rapidly into the pit, unclamps the instrument, turns the telescope to the position proper for direct view of the star, and fastens the clamping circle; bisects the star near its passage over one or more of the vertical wires by means of the micrometer-screw, and completes the direct observation in the usual way, by reading the microscope-micrometers for the second observation, and the punctures on the zenith-distance-micrometer register for both observations.

In observing the Sun, the transits of both limbs are taken for right ascension, and observations are made both of the upper and lower limbs for north polar distance. The assistance of a second observer is required to read the microscope-micrometers while the observer at the eye-end of the telescopes makes the horizontal wire touch the limb with the zenith-distance-micrometer and observes the transits. In observing the Moon two observers are required when the north polar distances of both limbs are observed. When the Moon is past full the observer reads the circle beforehand so that as soon as the punctures are made on the micrometer register he is able to unclamp and move the telescope and observe the transit of the second limb.

A detailed account of the *Chronograph* used with the Transit-Circle illustrated by engravings, is given in the *Appendix to Greenwich Observations* for 1856, and a brief description of the chief parts will be found in the *Introduction* for 1880 and preceding years.

The Personal Equation Machine is fully described in the *Introduction* for 1899 and previous years.

The Old Altazimuth, constructed by Messrs. Ransomes and May (as engineers), and by the late Mr. William Simms (as instrument-maker and optician), and erected in 1847, was in use till 1897 November 29, when observations with it were discontinued.

The New Altazimuth was constructed and erected by Messrs. Troughton and Simms in 1896, but was not brought into regular use till 1899, as it was found that various structural modifications were required particularly in regard to the arrangements for relieving the friction on the Y's.

This Altazimuth is virtually a reversible transit-circle (of Messrs. Troughton and Simms' well known form with some improvements), which can be planted in any definite azimuth (say 0° , 45° , 60° , 70° , 80° , 90° E. or W.), and then used for a complete set of observations (including stars for clock and azimuth error) essentially as a transit-circle, giving directly azimuth and zenith distances for the same instant of time. This is arranged by mounting the iron supports of the transit-circle on a circular base resting at three points of its circumference on a circular casting planed on its upper surface. The change from one azimuth to another is effected by turning the instrument about a central pivot on which it is slightly raised, the weight being relieved by friction rollers near the circumference of the circular base.

For observations in the meridian, the instrument is used exactly in the same way as a transit-circle.

The principle of the method of observing when out of the meridian is to secure observations of azimuth and zenith distance such that the mean of the times is as nearly as possible the same for the two sets of observations. Transits are taken over vertical wires and over horizontal wires so arranged as to secure complete observations in both elements in the different azimuths. The near agreement of the means of the times of transit is obtained by providing in addition to the systems of vertical and horizontal wires (each carried on a micrometer slide) a wire carried by a position circle micrometer which is set before the observation to the approximate inclination of the path of the object to be observed. When the star or other object enters the field the telescope is moved in zenith distance till it is on the position wire; the telescope is then clamped, and the transits are observed across a series of vertical and horizontal wires symmetrically with respect to the centre, as the star moves in its oblique course through the centre of the field, the position wire (which is only used for setting) having been previously moved away from the star.

The tube and axis of the telescope are made of phosphor-bronze, the axis and central part of the telescope tube being in one casting. This central part is barrel-shaped (instead of being in the form of a cube as in the transit-circle), and is stiffened inside by diaphragms so as to minimize flexure of the axis. A clear aperture of 6 inches diameter is made in it on each side, to admit of one collimator being viewed by the other when the telescope is in a vertical position. The two portions of the telescope tube,

which are exactly alike and are slightly conical, are bolted to the central casting. Two zenith distance circles are provided, one of which is fixed while the other can be turned on the axis, and, by means of a clamp and slow motion, fixed in any position, so that it may be made available for the determination of division errors by comparison of corresponding angular intervals on the two circles. These circles are discs of phosphor-bronze 3 feet in diameter, the divisions being on thin bands of palladium silver alloy let into the face of the bronze discs.

The pivots are of steel, and of 6 inches diameter. They rest on segmental bearings of bell-metal. The weight of the instrument on the Y's is relieved by live rings (ball bearings) round the axis, suspended freely by strong springs carried by the iron supports of the instrument, an arrangement which was substituted with success for the friction-rollers with upward thrust, originally supplied, as these were found to cause torsion in the axis and bearings.

Each circle is read by four microscopes. To carry the microscopes two broad wheels made of bronze are fixed to the iron supports of the instrument, the centres of the wheels being in the axis of the telescope. These wheels are screwed to the iron supports and are in addition pinned near their centres to prevent any possibility of movement.

As previously stated the instrument is reversible in its Y's. The counterpoise springs with the live rings are carried by the reversing gear during reversal so as to avoid the necessity for removal and readjustment of these springs. Another form of reversal of the instrument is obtained by turning it through 180° in azimuth, so that there are in all four essentially different positions in which it may be used.

The object glass is of 8 inches aperture and 8 feet focal length. The full aperture is used except for the Sun, when it is reduced to 6 inches. The magnification employed is 168 for stars, and 180 for the Sun.

The instrument is provided with two collimators in the meridian, each of 6 inches aperture and of 69 inches focus. They are each mounted on two pairs of parallel arms (counterpoised) turning about centres below, so that they can be brought up into the position for observation, or lowered to be clear of the telescope and dome, when they are housed under hinged flaps.

The situation of the Altazimuth is about half-way between the Old Observatory and the New Building. Its exact position relative to the centre of the transit-circle is 141 feet ($= 0^s.15$) E. and 166 feet ($= 1''.67$) S. It is mounted on a pier of concrete, 20 feet above the ground and has a clear view all round. This pier, which is quite clear of

the observing floor, rises from a large concrete foundation, 6 feet deep, on which the whole building rests. The dome is built in two halves which can be opened to a clear width of $4\frac{1}{2}$ feet, and is capable of rotation, so that the opening may be in any required azimuth.

The Chronograph used with the New Altazimuth since 1900 March, was made by Sir Howard Grubb. The barrel is of the same dimensions as for the Transit-Circle Chronograph, $19\frac{3}{4}$ inches long and $11\frac{3}{4}$ inches in diameter, revolving once in 2 minutes, so that 1 second of time is represented by a space of 0.3 inch on the sheet; one sheet will take a run of about 6 hours. The registration both of clock-signals and of observations is made by means of a single stylographic pen which, through the action of an electro-magnet, taps the slowly rotating barrel lightly, and leaves a dot of ink on the paper, when the circuit is completed, whether by the observer or by the Standard Sidereal Clock. The barrel is turned smoothly by a driving-clock of the usual form, without electric control, uniform motion between the clock-taps being the essential condition in this case. Four chronograph barrels, which can readily be interchanged, are provided. The chronograph is mounted in the ground floor of the Altazimuth Pavilion.

The Reflex Zenith-Tube, constructed under the direction of Sir G. B. Airy by the late Mr. William Simms.—A detailed description of this instrument is given in *Appendix I.* to the *Greenwich Observations for 1854*; in this place the following account will suffice. The object-glass (aperture 5 inches, focal length about 9 feet 8 inches) is mounted upon a fixed tube, which is in a vertical position; upon this tube it can be turned in azimuth. To the cell of the object-glass is firmly fixed the micrometer-frame (with its plane horizontal), revolving with the object-glass when the latter is turned. The bearing of the screw of micrometer A is on the fixed micrometer-frame. The moving frame of micrometer A carries the bearing of the screw of micrometer B. Micrometer B carries the wires by which the star's image is bisected. The eyepiece is a four-glass diagonal eyepiece bent at right angles between its third lens and its fourth lens (counting from the eye), and having a diagonal-prism-reflector at the place where it is bent; and so placed that the fourth lens can receive rays of light from the direction of the nadir, and can, by means of the diagonal prism, transmit them horizontally to the eye. The fourth lens (which looks vertically downwards) and the diagonal prism are placed nearly over the centre of the object-glass, being carried by arms which are fixed to the large tube; the small horizontal tube which carries the remaining lenses of the eyepiece is supported by the large tube of the telescope, and does not project over any part of the object-glass. Below the

REFLEX ZENITH TUBE: SHUCKBURGH EQUATORIAL: SHEEPSHANKS EQUATORIAL. xvii

object-glass, at a distance nearly equal to half its focal length, is a trough of quicksilver, which till the end of 1888 was protected from tremors, with only moderate success, by an arrangement of caoutchouc supports, described in previous Introductions to 1880. From 1889 February 13 an amalgamated copper trough has been used, with the result that the steadiness of the images has greatly improved. A prismatic focussing rod with scale reading to 1-200th of an inch and ivory point for contact with the surface of the mercury has been applied to the instrument in place of the former light brass rod and wooden float; the focussing rod and the micrometer-frame are both of gun-metal. The image of the star, as it passes near to the zenith, is formed in the plane of the micrometer wires. It will easily be seen that, to ensure accuracy of result, no firmness of construction is requisite in this instrument except in the connection between the micrometer and the object-glass. The instrument is mounted about 18 feet south-west of the transit-circle.

The Shuckburgh Equatorial.—This instrument was constructed by Ramsden, and is one of that class usually called the “English equatorial.” It was originally the property of Sir George Shuckburgh, and is fully described by him in the *Philosophical Transactions* for 1793. It was presented to the Observatory in the year 1811. The telescope is 5 feet 4 inches in length, and has an object-glass of 4.1 inches aperture. It is mounted in the north dome, the situation of which is most unfavourable, as the south-western sky is concealed by the Octagon room.

The Sheepshanks Equatorial.—This instrument was erected in the year 1838. The aperture of the object-glass is about 6.7 inches, and its focal length about 8 feet 2 inches. The object-glass was made by M. Cauchoix, of Paris, and was presented to the Observatory by the Rev. R. Sheepshanks. Its definition is good: a small quantity of colour from the secondary spectrum, and a diffusion of light from brilliant objects, being the principal defects.

It is mounted in a dome, 19 feet east and $1\frac{1}{2}$ feet north of the transit-circle. The mounting, particulars of which are given in the *Introductions* to the *Greenwich Observations* for former years, is similar in general form to that known as the “German equatorial,” and was constructed by the late Mr. T. Grubb, of Dublin. Besides several negative eyepieces unfurnished with wires, this telescope has a wire micrometer, and a comet eyepiece with thick wires and position circle, which was supplied on 1890 November 10. On 1892 November 18 a new set of wires was mounted in this eyepiece, as the wires were found to be bent.

GREENWICH OBSERVATIONS, 1908.

The Great Equatorial.—This instrument now carries an object-glass of 28 inches aperture and 28 feet focal length on the mounting which was originally designed for the Merz object-glass of 12·8 inches aperture and 17 ft. 10 in. focal length. This mounting was constructed by Messrs. Ransomes and Sims (as engineers), and the late Mr. William Simms (as instrument-maker), and a complete description of it, with engravings, is attached as *Appendix to Greenwich Observations* for 1868. The form of the instrument is the “English equatorial,” the polar-axis turning on pivots at its extreme ends, and including between its two sides the telescope and declination-circle, which are mounted on a declination-axis with pivots at its opposite extremities. The object-glass of 28 inches clear aperture and 27 ft. 10 in. focal length was made by Sir Howard Grubb from discs supplied by Messrs. Chance, of Birmingham, and was brought into use at the end of 1893. It is of special form, adapted to photography as well as to eye-observation, on the plan proposed by Sir George Stokes (in a letter to me dated 1886 August 16) of reversing the crown lens to correct for the spherical aberration introduced by the further separation of the lenses necessary for photographic correction. Mechanical means are provided for readily effecting this reversal and separation without risk. It is found by trial that the further separation required for photographic correction is about 3·5 inches, and that the focus is thereby shortened by about 23 inches. When the crown lens is in the position for visual observation, the radii of curvature of the surfaces are:—

1st surface—146 in. convex	} Crown.	3rd surface— 138 in. concave	} Flint.
2nd „ 134 in. „		4th „ 1000 in. convex	

and the edges of the lenses are separated by 0·4 inch, the minimum focal length being for rays about midway between D and E.

The telescope tube, made of steel and cast-iron by Sir Howard Grubb, is specially adapted to the conditions, being provided with a sliding eye-end arranged for either direct view or diagonal view by means of a silver on glass plane reflector, and permitting of observations near the zenith and the pole, when the eye-end is shortened sufficiently by the sliding motion to clear the floor and the inside of the polar frame. The Corbett telescope, of 6½ inches aperture and 8 feet focal length, is mounted on the tube as a finder or guiding telescope. The position and transit micrometers and various eyepieces supplied for the 12·8-inch Merz refractor have been adapted to the 28-inch telescope, and have been fitted with electric illumination of the field and wires. A double-image micrometer is also used.

The dome was constructed by Messrs. T. Cooke and Sons and erected in 1893 April, replacing a cylindrical wooden dome. It is of a form specially adapted

to accommodate the 28-inch refractor, the diameter being greater than that of the tower on which it is carried. A description with diagrams is given in the *Monthly Notices, R. A. S.*, vol. li., p. 436. The diameter of the dome is 36 feet at a height of 7 feet above the rail, contracting to 31 feet at the base, and the shutter-opening is 7 feet wide throughout on both sides of the zenith down to the horizon. The framework is of T-iron vertical and horizontal ribs bolted to a circular girder forming the base, the opening being framed with curved vertical girders, and the whole is covered with papier-mâché. The shutter is in two parts, with a division right down the middle, each half being a balanced shutter 3 ft. 6 in. wide, extending right over the dome from the curb, through the zenith, to the opposite part of the curb, and carried by a wheel running on a rail at the zenith. The dome rotates on a system of wheels running freely on the horizontal surface of the wall-curb.

The Thompson Equatorial, made by Sir Howard Grubb.—This instrument, carrying a 26-inch photographic refractor and a 30-inch reflector on the same mounting, was presented to the Observatory by Sir Henry Thompson, and was brought into use at the end of 1898. It is erected on the central tower of the New Observatory building under the 30-feet dome, and is approximately 350 feet south and 224 feet east of the centre of the transit-circle. The equatorial mounting, which is of the German form, modified to allow of complete circumpolar motion without reversal, carries on one end of the declination axis the 26-inch photographic refractor, the Merz $12\frac{3}{4}$ -inch refractor (used as a guiding telescope), and the 9-inch photoheliograph; and on the other end the 30-inch Cassegrain reflector (made by Dr. Common), with a 6-inch refractor as a guiding telescope.

The instrument is driven by a clock with electrical control similar to that of the Astrographic Equatorial.

The 26-inch Refractor is a photographic telescope of 26 inches (0.66 metre) aperture and of 22 feet 5 inches (6.83 metres) focal length, so that its scale is double that of the Astrographic refractor, and one millimetre on the photographic plate represents $30''\cdot2$. The breech end of the instrument is arranged for a plate-carrier taking photographic plates 12 inches square, suitable adapters being inserted when smaller plates are used. The breech end is also arranged to carry a camera-box, 33 inches in length, fitted with a negative enlarging lens, 1.45 inches in diameter, and 3.3 inches in focus, and with a rapid exposing shutter, for photographing sun-spots and other objects on an enlarged scale. When used for this purpose, a diaphragm carrying a roller spring blind is placed before the object-glass, so that the sun's light may only be permitted to fall on the object-glass immediately before the exposure of the plate is to be made.

The aperture is usually stopped down to 12 inches. The image of the sun in the primary focus of the telescope is 2·5 inches in diameter at mean distance, and this is enlarged 12 diameters, *i.e.* to a scale of 30 inches to the solar diameter, for the image formed on the photographic plate. The exposing shutter is of the same form as that used in the Thompson Photoheliograph (*vide infra*). The plate-carrier for the camera-box is constructed to take plates 10 inches by 12 inches or, by use of suitable adapters, of smaller sizes. The whole sun is, therefore, not photographed at one time, but only limited areas; generally a single spot-group of special size or interest.

The *Merz Refractor*, of 12·8 inches aperture and 17 feet 10 inches focal length, is mounted as a guiding telescope to the 26-inch, and is above it when the instrument is pointing south, and the 26-inch is on the west of the pier of the equatorial. The eyepiece of the Merz is mounted on cross-slides (parallel and perpendicular to the equator) which permit of a guiding star being observed at a distance of 45' from the centre of the field, and are furnished with scales divided to minutes and read by verniers to 5".

The *Thompson Photoheliograph*, presented to the Observatory by Sir H. Thompson in 1891, is a photographic refractor by Sir H. Grubb of 9 inches aperture and 8 feet 10 inches focal length, with a Ross enlarging doublet of 4·3 inches focus. Its tube is mounted on the 26-inch refractor on the side opposite the Merz refractor. The image of the Sun in the primary focus is 1 inch in diameter, and this is enlarged by the doublet to about 7·4 inches on the photographic plate, the whole length of the instrument being about 12 feet 4 inches. At the principal focus cross spider-lines are placed, which give facilities for determining the position angles of sun-spots on the photographs. The exposure is given (as in the Dallmeyer photoheliograph, *see* page xxii) by a shutter at the primary focus having a slit in it of adjustable width; but, whilst the shutter of the Dallmeyer photoheliograph travels in a groove, and is drawn downwards by a strong spring, in the Thompson, the shutter, which is made of aluminium for the sake of lightness, turns about a pivot 18 inches from the exposing slit, under the action of a spring placed about midway between the pivot and the slit, and giving a very rapid motion to the latter. This telescope is also provided with a 9-inch object-glass prism by Mr. Hilger.

The 30-inch Reflector.—This instrument was constructed under the supervision of Dr. Common who undertook the figuring of the mirrors. It may be used as a simple reflecting telescope for obtaining photographs in the principal focus, or as a Cassegrain. The mirrors are made of silver on glass, the concave being of 30 inches diameter, with a circular hole of 6 inches diameter. The weight of this mirror is 265 lbs., the glass of which it is composed being 4 inches thick. When not in use, it is covered by a

plate-glass disc, with a hinge on one side which allows of its being turned so as to lie along the telescope tube (made D-shape to receive it) when observations are taken. The focal length of the large mirror is 3.48 metres or 11 feet 5 inches, so that 1^{mm}. at the principal focus corresponds to 59".3. Two convex mirrors of focal lengths 24 inches and 38 inches are provided, the equivalent focal lengths of the combinations being 76 feet and 49 feet respectively.

The convex mirror, or the photographic plate-holder, is carried by an arm mounted in a slide so that it can be fixed in a convenient position, and the final adjustment for focus can be made from the eye-end by means of a rod with geared wheels actuating a screw in the boss of the arm, which gives a small motion to the mirror or plate-holder.

The Hodgson 6-inch Refractor, of 7 feet 7 inches focal length, is mounted on the reflector as a guiding telescope. It is furnished with cross-slides like the Merz and Astrographic refractors. The scales on the slides are read by verniers to 5".

The dome was constructed by Messrs. T. Cooke and Sons, of York, in 1884, and was formerly used for the Lassell reflector. It is hemispherical, of 30 feet diameter, and is covered with papier-mâché on a framework of angle-iron. It is carried by nine wheels or rollers (each 2 feet in diameter) fixed to the iron curb of the dome, and running on a flat rail, horizontal rollers being provided to prevent excessive lateral motion from the force of the wind or other cause, and it can thus be turned with great ease. The shutter-opening extends from beyond the zenith to the horizon, and is closed by a single curved shutter (3 feet 6 inches wide at the zenith, and 6 feet wide at the horizon), which turns about a point in the dome-curb opposite to the opening, and runs on guiding rails at the horizon and near the zenith, the curved shutter being continued by an open framework to complete the semicircle. The dome was erected on the central tower of the New Observatory building in 1896 September.

The Astrographic Telescope, of 13 inches aperture with 10-inch guiding telescope, mounted in a dome over the upper computing room.—This instrument was constructed by Sir Howard Grubb, F.R.S., on the lines laid down by the *Congrès Astrophotographique International pour le Levé de la Carte du Ciel* in 1887, and was brought into use at the end of 1890. The 13-inch photographic telescope and a 10-inch visual telescope are firmly connected, the tubes being of iron. The apertures of the object-glasses are 0^m.33, or 13ⁱⁿ.0, and 10ⁱⁿ.0 respectively, and the focal lengths of both telescopes 3^m.43 or 135ⁱⁿ.1, so that 1^{mm}.0 on the plate corresponds to 1'.0. [The focal length of the photographic telescope is more exactly 3^m.441 or 135ⁱⁿ.4, so that 1^{mm}. represents 0'.9990.] The photographic telescope is corrected, as regards spherical and chromatic aberration, for rays near

xxii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Fraunhofer's line G. It is arranged to carry a plate 16^{cm.} square, with special provision for exact focussing and orientation. The eyepiece of the 10-inch visual telescope is mounted on cross-slides (parallel and perpendicular respectively to the equator) which are furnished with scales reading to 1-12th of a millimetre or 5'', and permit of the observation of a guiding star up to a distance of about 45' from the centre of the field. The mounting is of the German form, so arranged as to allow a range of $1\frac{1}{2}$ hours' motion on each side of the meridian without reversing the telescope. A large counterpoise is thus necessary, both because of the double weight, and of the distance from the polar axis of the two telescopes; but the movement in right ascension is very easy, owing to the arrangement adopted to relieve the friction of the polar axis. The greater part of both the transverse and end thrusts of the polar axis is received on a single anti-friction bearing, carried by a separate vertical column, disposed directly under the centre of gravity of the instrument. The rollers of the bearing turn round horizontal axes, and a bevelled collar on the polar axis rests on them. The column transmits the weight to counterweight levers. The driving clock is placed inside the stand and is controlled electrically by a seconds pendulum. The detector of the control is similar in principle to that used in Sir David Gill's form; and the system of correctors by differential wheels was devised by Sir Howard Grubb (vide *Proceedings, Institute of Mechanical Engineers*, 1888 July 31, page 311). The telescope is also provided with an electric hand control, by means of which the observer can accelerate or retard the motion in right ascension and thus keep the guiding star bisected in the guiding telescope.

The Dallmeyer Photoheliograph has an object-glass of 4 inches aperture and 5 feet focal length, forming an image of the Sun half an inch in diameter; this image is enlarged by a secondary magnifier to 8 inches on the camera screen, where the sensitive plate is inserted, the whole length of the instrument being about 9 feet 7 inches. The exposure is given by a shutter having a slit of adjustable width, which is carried by a spring across the primary image. The instrument is equatorially mounted and provided with a driving clock.

Two detached telescopes, one of which is 46 inches in focal length, and the other 30 inches.—The diameters of the object-glasses of these telescopes are 3·6 inches and 2·7 inches respectively. The telescope formerly belonging to the Western equatorial has also been mounted for use as a detached telescope; its focal length is 36 inches, and the aperture of its object-glass 3·6 inches. The 62-inch telescope mentioned in former years, and that of the transit-instrument formerly used as a collimator to Troughton's transit, are now mounted on tripod stands for general use.

Two Newtonian Reflectors, the largest being one of 10 feet, by Sir W. Herschel.

A portable transit-instrument with axis-view by Brauer of St. Petersburg and *a portable 18-inch altazimuth* by Troughton and Simms, obtained in the year 1866.

The following instruments, used in the Transit of Venus Expeditions, 1874 and 1882, are now transferred to the Observatory :—

Five 6-inch equatorials complete, viz.: the “Lee,” “Hodgson,” “Cooke,” “Simms No. 1” and “Simms No. 2”; two 4-inch portable telescopes by Troughton and Simms; five 3-inch portable transit-instruments by Troughton and Simms; a 14-inch altazimuth and two 14-inch vertical circles or altitude instruments also by Troughton and Simms; and five photoheliographs by Dallmeyer. Several of these are, at the present time, lent to other institutions. The Naylor 6-inch equatorial and a 14-inch altitude instrument, which had been employed at Bermuda for the observation of the Transit of Venus, 1882, were lost on the return voyage in the wreck of the “City of Brussels” at the entrance of the river Mersey. The “Cooke” 6-inch equatorial used in the Transit of Venus Expedition, 1882, was transferred to the Observatory in 1884 to replace the Naylor equatorial.

One of the portable transit-instruments is generally mounted in the *Transit Pavilion*, a brick building in the front court, the centre of which lies in Bradley’s meridian. The roof of the building, which was completed in 1891 October, consists of a pair of semi-domes, which slide apart to leave a clear opening of uniform width (2 feet 6 inches) in the meridian. The pier for the transit is 19 feet west and 67 feet 10 inches north of the centre of the Transit-circle. The instrument is available for longitude operations, or for practice observations by officers of the Royal Navy and others, or for time determinations on occasions when the Transit-circle is under repair.

The Sidereal Standard Clock, constructed by Messrs. E. Dent and Co., is fixed to the north wall of the Magnet-house basement, as in this apartment the temperature is kept nearly uniform. The escapement will be found described in vol. iii. of the *Transactions of the Cambridge Philosophical Society*: it is a detached escapement very closely analogous to the ordinary chronometer-escapement, the pendulum receiving impulse only at each alternate vibration; consequently, the escape wheel and seconds hand move only at alternate seconds (the even seconds). The pendulum is compensated in the following way. A central steel rod is encircled by a zinc tube, which rests on the rating nut on the steel rod; the zinc tube is in its turn encircled by another steel tube, which rests at its upper end on the zinc tube, and carries at

xxiv INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

its lower end the cylindrical leaden pendulum bob attached at its centre. Slots are cut in the outer steel tube, and holes made in the intermediate zinc tube, with the object of exposing equally all parts of the compound pendulum rod to the action of temperature. For final adjustment of the rate there is placed on the crutch rod a sliding weight, which can be raised or lowered by a nut at the level of the crutch-axis without disturbing the pendulum. The rate of the clock is so steady that, when first mounted, the barometric inequality was indicated with the greatest regularity, the daily losing rate of the clock being increased by $0^{\text{m}}.3$ for an increase of 1 inch of barometer reading.

In the autumn of 1873 an apparatus for correction of this inequality was applied to the clock by Messrs. E. Dent and Co., and has been in action regularly since that time. This compensating apparatus of the Sidereal Standard is founded on the magnetic principle, long previously in use for daily adjustment of the Mean Solar clock. Two bar magnets, each about six inches long, are fixed vertically to the bob of the clock pendulum, one in front, the other at the back, their lower ends being nearly level with the bottom of the pendulum bob. The lower pole of the front magnet is a north pole, and the lower pole of the back magnet a south pole. Below these a horseshoe magnet, having its poles precisely under those of the pendulum magnets, is carried transversely at the end of a lever, the opposite arm of which is attached by a connecting rod to a float in the lower leg of a syphon barometer, placed in one corner of the clock-case. The area of the cistern in which the float rests is four times as great as that of the upper tube. For change of one inch of barometer reading the horseshoe magnet is thus shifted two-tenths of an inch; and as the average distance between its poles and those of the pendulum magnets is about $3\frac{3}{4}$ inches, the change of rate produced by increase or decrease of the magnetic action is sensibly uniform. As the clock gained with low barometer reading, it was necessary to place the horseshoe magnet so that there should be attraction between its poles and the adjacent poles of the pendulum magnets. The action of this apparatus is found to be quite successful.

Galvanic contact for registration of the clock-beats is made by a wheel of 30 teeth on the escape-wheel arbor, a tooth of which at every beat of the clock presses together two light springs. Another pair of springs is also pressed together at the beginning of each minute by an arm on the same arbor, and a supplementary signal is thus sent through a galvanometer as a check on the numeration of seconds. This arrangement was substituted in the latter part of 1881 for that formerly employed (in which a pin on the pendulum was used to press together the contact springs) in order to avoid any effect on the pendulum, contact being made in the

part of the beat when the pendulum is quite detached from the clock-train, after the impulse has been given. The currents obtained from the Sidereal Standard at every alternate second are used to drive a relay, from which independent circuits are derived. One of these is appropriated to the seconds-magnet of the altazimuth chronograph; another drives a relay placed, in the computing room, on the desk of the Superintendent of the Time Department. From this relay are derived several circuits, one for driving a galvanic chronometer in the Time Department, another for use with the transit-circle chronograph, and a third for use in the Transit Pavilion. No practical difficulty is found in sub-dividing the interval of two seconds between the clock-beats on the chronograph. The numeration of seconds on the chronograph is readily obtained from an automatic signal from the Sidereal Standard clock on an independent circuit, or from occasional comparisons with the clock Hardy.

The clock Hardy, in the Transit-circle room, was originally furnished with Hardy's escapement, for which a dead-beat escapement was substituted by Dent in the year 1829. The jewelled holes were removed by Dent in 1836, and the pivots now turn in brass holes. It has a zinc and steel pendulum. The clock is provided with contact-springs for the purpose of occasional registration of its seconds on the chronograph. Upon the escape-wheel-arbor is mounted a wheel having 59 teeth (that is, 60 teeth with one cut away), and a tooth of this wheel at every beat of the clock presses together two light springs. The omitted tooth in the wheel corresponds to the 1^s of each minute.

The clock "Dent 2" has been fitted with springs and placed in the Record Room for use when the Sidereal Standard is under repair. There is also a clock marked "Arnold 1," having contact apparatus similar to that in the clock Hardy, in order that the clock may, if required, be temporarily used in the place of Hardy; one in the Sheepshanks dome, marked "Earnshaw," with a zinc and steel compensation pendulum; one in the New Altazimuth dome, marked "Graham 1," with a mercurial pendulum; in the Thompson Equatorial dome, one marked "Dent 2016," with a zinc and steel pendulum and seconds' contact, and one marked "Graham 2," fitted with contact springs and with a zinc and steel pendulum, which is available for use in longitude determinations; one in the Astrographic dome, marked "Dent 2017," with a zinc and steel pendulum; one marked "Dent 2009," with a wooden-rod pendulum in the Great Equatorial dome; one marked "Molyneux," with a wooden-rod pendulum in the Reflex Zenith-Tube room.

The Mean Solar Clock, "Dent 2012," is placed in the lobby at the foot of the Octagon room staircase, near to the trigger of the ball-apparatus. This clock (which

GREENWICH OBSERVATIONS, 1908.

d

has been substituted for the electrical clock "Shepherd" formerly in use) has a zinc and steel pendulum, and is fitted with springs for sending out hourly signals to the Post Office and for working the mean solar dials distributed throughout the Observatory. On the escape-wheel-arbor are two wheels, each containing 30 teeth, which close contact-springs alternately, one wheel at the even, the other at the odd seconds. The currents thus set up actuate two relays, by means of which various electromagnetic clocks in different parts of the Observatory are driven synchronously with the Mean Solar clock. Arrangements are made so that in the case of failure of the clock "Dent 2012," the circuit can be readily switched on to the electrical clock "Shepherd."

One of the synchronous clocks has a large dial exposed to public view on the east boundary wall of the Observatory; four are in the chronometer rooms, and are used in the daily comparisons of chronometers; one is in the upper computing room; one (of the dimensions of a chronometer) is upon the desk of the Superintendent of the Time Department; one is in the hall of the Astronomer Royal's dwelling-house and four are in the office rooms of the New Observatory building. A comparison of the mean time chronometer with the chronometer on the same desk synchronous with the sidereal standard clock (as mentioned in a preceding paragraph) is, in fact, a comparison of the mean solar clock with the sidereal standard. The time shown by the sidereal standard at comparison being corrected for its error (as ascertained from star-transits), the true sidereal time at the comparison is found; and this is converted, by calculation, into true mean solar time at the comparison. The difference between this and the time shown by the mean solar clock is the error of the latter. Then, by means of a commutator on the same desk, the circuit of a galvanic current, passing through a coil (with no iron core) in the clock-case, can be made either to attract or to repel a magnet attached to the pendulum, and thus to accelerate or retard the mean solar clock as long as may be necessary for its correction. The clocks connected with the mean solar clock necessarily receive the same correction. The arrangement for giving hourly signals is as follows. The mean solar clock closes a circuit precisely at each hour: that at 13^h, by the intervention of electro-magnets, pulls automatically the ball-trigger and drops the time-ball at Greenwich. Until 1885 October 27 there was an automatic arrangement for reversing the direction of the current through this apparatus; but, as this sometimes caused failure in the signal and was found to be unnecessary, it was removed on that day. At every hour also the primary clock-current by relay-action causes a galvanic current to pass on a line of wire to the central station of the Post Office Telegraphs, St. Martin's-le-Grand, London, where it is distributed by automatic action of the "Chronopher," on some of the principal

lines of railway and to important provincial towns. The hourly current passes also to the clock tower of Westminster Palace for the guidance of the superintendent of the clock. The clock is not regulated or corrected by any direct galvanic action, but signals are received at definite times from the clock, which show daily at Greenwich the deviation of the clock from true time. The 13^h current drops a time-ball at Deal, the property of the Admiralty; and from this ball a return galvanic current is automatically communicated to Greenwich at the instant that the ball has completed its descent. At Devonport and Portsmouth and Portland local clocks, giving hourly time signals, are corrected daily by the help of the current at 10^h. This time signal automatically starts an auxiliary seconds' pendulum, suspended freely just behind the clock pendulum, which is brought into coincidence of beat with the free pendulum by an attendant (by means of electro-magnetic action as in the Greenwich mean solar clock). The Devonport clock sends a return signal to Greenwich daily at 13^h. 0^m. 39^s. 0, the Portsmouth clock at 13^h. 0^m. 20^s. 0, and the Portland clock at 13^h. 0^m. 30^s. 0, to serve as a test of the accuracy with which the local clocks have been corrected.

The following clocks are also in use in the Observatory as Mean Solar Clocks :—
 “Mudge and Dutton,” with a gridiron pendulum, in the upper chronometer room ;
 “Dent 2014,” with a wooden-rod pendulum, in the lower chronometer room ;
 “Graham 3,” with a mercurial pendulum, in the ball lobby, occasionally used for facilitating the regulation of the Mean Solar Clock (described above). Several chronometers are also kept in the lower computing room for occasional use. Besides these there are two journeymen or assistant clocks ; an electric clock by Shepherd, formerly used at Ashford ; and a watchman's clock.

The following clocks are on loan :—“Dent 1916” and “Dent 2013” at the Cape Observatory ; “Dent 2011” at the Kew Observatory ; “Arnold 2” at the Oxford Observatory ; and “Dent 2010” at Devonport, where it is employed to drop the Time Ball.

III. *Subjects of Observation in the Year 1908.*

Transit-Circle.—The Sun, Moon, and Moon-culminating stars have been observed on the meridian at every practicable opportunity.

With regard to the planets, the following rules have been observed :—The inferior planets, Mercury and Venus, have been observed on the meridian at every practicable opportunity, excepting Sundays. The superior large planets, Mars, Jupiter, Saturn, Uranus, and Neptune, have not generally been observed when their solar time of meridian passage is greater than 14^h, excepting at times when the Moon passes the

xxviii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

meridian later than 14^h . The small planets have been observed only when the time of meridian passage is earlier than 13^h . The observations of all planets are intermitted on Sundays.

In the meridian observations of the stars, continuous observation is maintained of all stars in Prof. Newcomb's *Catalogue of Fundamental Stars* visible in this latitude.

The other stars observed are, since the beginning of 1906, taken from a new working catalogue of stars of magnitude 9.0 and brighter between the limits $+24^\circ$ to $+32^\circ$ of declination, serving as reference stars for the Oxford Astrographic Zones.

The Altazimuth has been used in the meridian for observations of the Sun, Moon, planets and fundamental stars. Out of the meridian the instrument was generally used for the Sun, Moon and planets, and the necessary adjustment stars.

The Reflex Zenith tube.—Regular observations with this instrument were recommenced in June 1902. A working catalogue has been constructed of stars down to magnitude 7.0 which transit within $50'$ of the zenith at Greenwich, and stars were observed from this list continuously throughout the year with the object of determining the variation of latitude as well as the constants of aberration and nutation.

The Sheepshanks Equatorial has been used for observing occultations of stars by the Moon.

The 28-inch Refractor has been in use with the object-glass in the visual position for micrometric observations of double stars.

Measures of the equatorial and polar diameters of Jupiter and of Saturn and its ring-system have also been made.

The Thompson Equatorial.—With the 26-inch refractor photographs have been taken of Neptune and its satellite, and also of Jupiter and Saturn for determining their positions in connection with the observation of the faint outer satellites with the 30-inch reflector.

With the 30-inch reflector photographs for position have been taken of Comet *c* 1908, of Minor Planets, of Jupiter's Sixth, Seventh, and Eighth satellites, and of Saturn's Ninth satellite, Phœbe.

In addition, an extensive series of photographs of Comet *c* 1908 of longer exposure were taken for the determination of the motions and changes taking place in the tail.

Photographs of the Sun have been taken on every day when the weather permitted, with the Thompson photoheliograph.

The Astrographic Equatorial.—The photography for the Greenwich Section of the Astrographic Chart having been completed, work with this instrument has been mainly confined to replacing Chart plates which were inferior to the general standard.

In connection with the observations of Jupiter's outer satellites and of Saturn's Ninth satellite, Phœbe, with the Thompson Equatorial, a number of photographs were taken of fields of the reference stars used for determining the plate-constants.

The 10-inch guiding telescope has also been used for observing occultations of stars by the Moon.

IV. *Explanation of the Printed Observations.*

§ 1. *Transits observed with the Transit-Circle, and Computations of Apparent Right Ascension.*

The observations here printed in detail are those of the Sun, Moon, planets, fundamental and adjustment stars. Commencing with the year 1897, the printing of the details of observations of other stars has been discontinued, the daily results only being given in the *Ledgers of Mean Right Ascensions and Mean North Polar Distances of Stars*.

The *first* column on each page contains the day, which is always supposed to commence with the transit of the Sun.

The *second* column contains the numbers for convenience of reference.

The *third* column contains the name of the object observed. With respect to the Sun, Moon, and planets, the limb whose transit is observed is always mentioned, and if the limb is defective the requisite correction for defect of illumination is given in the notes. If no limb is mentioned, it is to be understood that the estimated centre was observed. The centre is observed only when the planet's disc is so small and so round as to make it easier to estimate the centre than to determine with accuracy the place of the limb. In the case of Mercury (and more rarely of Venus), whose limbs are sometimes very badly defined, a correction for defective illumination is applied when the estimated centre of the illuminated portion is observed, to reduce to the centre of the disc, see p. xxxiii. In the case of the Moon, observations of the crater Mösting A have been made as well as of the limbs whenever observable. With

xxx INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

regard to the stars, the proper names which have commonly been used in the *Greenwich Observations* are adopted in preference to other names. For other stars, the names have been taken in the following order of preference :—

1. Flamsteed's constellation-No. and constellation, with Bayer's letter ; taken from Baily's edition of Flamsteed, or the British Association Catalogue.
2. The No. in Bessel's *Fundamenta*, &c., deduced from Bradley's Observations, referred to as "Bradley."
3. The No. in Auwers's edition of Mayer's Catalogue.
4. The hour and No. in Piazz's Catalogue, edition 1814.
5. The No. in Groombridge's Catalogue.
6. The No. in Baily's edition of Flamsteed's Catalogue, referred to as "B.F."
7. The hour and No. in Weisse's two Catalogues of the Stars in Bessel's Zones, referred to as "W.B." and "W.B. (2)" respectively.
8. The No. in Lalande's Catalogue, published by the British Association, or in that edited by Fedorenko, or in the Supplement, referred to as Lalande, Lalande F., and Lalande F. (2) respectively.
9. The No. in Lacaille's Catalogue of Southern Stars, published by the British Association.
10. The No. in Oeltzen's two Catalogues of Argelander's Zone Observations, referred to as "Oeltz. Arg. (N.)" or "(S.)" respectively.
11. The No. in the First Radcliffe Catalogue (1845).
12. The No. in Carrington's Red Hill Catalogue.
13. The Zone and No. in Argelander's Bonn *Durchmusterung* or in Schönfeld's continuation of the *Durchmusterung* (Bonn Catalogue, Sections I, II, III, and IV.; Bonn Observations, Vols. III., IV., V., and VIII.), referred to as "B.D."
14. The No. in the Greenwich Catalogues; in the British Association Catalogue, referred to as B.A.C.; in the Cape Catalogue for 1880; or in the Catalogues of the *Astronomische Gesellschaft* (1875).

In observing a double star, the brighter star (if it is not otherwise expressed) is always observed.

When no mark is attached to the name of the object observed, it is to be understood that the observations were made by the galvanic or chronographic method, which was generally in use throughout the year. In any exceptional case where the eye-and-ear method is employed, the observation is distinguished by having the letters (E & E) affixed to it. The letter R denotes that the object was observed by reflexion in the trough of mercury.

The *fourth* column contains the initials of the observer's name.

The next *ten* columns contain the clock-times (seconds and decimals of a second only) at which the object was observed to pass each of the wires. For eye-and-ear

transits, the wires used are denoted by the letters M, N, O, T, X, Y, Z. The wires used for galvanic transits are denoted by the letters O, P, Q, R, S, T, U, V, W, X; O, T, and X being wires of the eye-and-ear system, and the others new wires inserted between O, T, and X. The intervals O-P, P-Q, R-S, S-T, U-V, V-W, W-X are nearly equal; the interval Q-R is nearly double, and the interval T-U nearly treble the former. This system of ten wires for the galvanic transits was introduced on 1891 October 28, in substitution for the system of nine wires previously in use.

The nine azimuth stars are generally observed by turning the R.A. micrometer-screw with the right hand until the star is bisected by the middle wire T (or sometimes one of the other wires), the instant of accurate bisection being recorded on the chronograph with the left hand. The reading of the micrometer is then recorded, and the operation repeated until ten observations are obtained. The readings of the R.A. micrometer are always mentioned in the notes. As the "reduction to the centre wire" in this case depends on the value of successive revolutions of the micrometer-screw, the errors of this screw have been investigated. The results of the most recent investigation of these errors are given in the *Transit-Circle Tables* for 1906. It is considered that the errors are too small to necessitate correction.

The eye-and-ear method of observation is exclusively employed for stars, not being adjustment stars, within three degrees of the pole.

In observing by galvanic contact, the observer merely makes contact of the galvanic spring at the instant of the object passing the wire. Occasionally, when no object is under observation, he makes contacts in any arbitrary manner (usually by a commutator which puts the contact-springs of the clock Hardy into circuit with the "observations" pricker-magnet of the chronograph), recording in his book the time by the clock Hardy; and thus the numeration of hours, minutes, and seconds on the punctured sheets (referred to the Sidereal Standard clock) is easily established. The punctured sheets are examined the next day, the numeration of minutes and seconds is written upon them, and the transits are read into the transit-book.

In observing by eye-and-ear, it is the practice to take a second from the clock-face, before the transit over the first wire, and to preserve the counting by listening to the beats of the clock, and not to look again at the clock-face till the transit is completed. The fraction of the second is estimated by remarking the place at which the object is seen at the successive beats of the clock. Errors in the hours and minutes are

xxxii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

seldom alluded to in the notes of the printed observations; but every alteration of the seconds is carefully recorded.

All transits of stars observed by galvanic contact are usually referred to the Sidereal Standard clock; but for stars observed by eye-and-ear, the clock Hardy is necessarily still used. The times printed in the columns "Seconds of Transit over the Wires" are those of the observations as made, the letters (E & E) in the third column indicating when the clock Hardy was used. The difference between the indications of the two clocks is given in the notes.

The *fifteenth* column contains the *Concluded Transits*, giving the hour and minute of the time of meridian passage (which have not been given in the preceding columns), as well as the mean of seconds.

For stars observed by eye-and-ear by the clock Hardy, the *Concluded Transit* includes the correction for reduction of the transit by Hardy to the Sidereal Standard clock. The correction applied in each instance is given in the footnotes.

When transits have been observed over an unsymmetrical group of wires, corrections are applied to reduce each separately to the centre wire. When transits have been observed over all the wires, or over a selection of wires symmetrical with respect to the centre wire, the necessary small correction is applied to the mean of the observed transits.

The adopted equatorial intervals of the wires used are given in the *Transit-Circle Tables* at the end of this *Introduction*. These have been determined from observations of close polar stars.

For stars above the pole, the designations of Wires I., II., III., &c., at the heads of the columns of printed transits correspond to O, P, Q, &c. (or to M, N, O, &c., when the object is observed by eye-and-ear), in all cases excepting Polaris, &c. For stars below the pole, the designations I., II., III., &c., correspond to X, W, V, &c., or to Z, Y, X, &c.

For a planet, the correction to reduce to the centre wire, computed as for a star, is to be multiplied by

$$1 + \frac{\text{daily increase of R.A. in seconds of time}}{24 \times 60 \times 60};$$

but it has generally been found easier to add, to the computed interval between the mean of wires and the centre wire, the amount of the planet's increase of right-ascension for that interval, which is very readily determined from the numbers given in the *Nautical Almanac*.

For the Moon, the equatorial intervals for the wires observed are added together, and the sum is divided by the number of wires; this quotient is then multiplied by

$$\frac{3600 + I}{3600} \times \frac{\sin \text{Moon's geocentric } Z.D.}{\sin \text{Moon's apparent } Z.D.} \times \secant \text{ of Moon's geocentric declination,}$$

where I is the increase (in seconds of time) of the Moon's R.A. for the transit over a meridian upon the Earth distant by 1^h of terrestrial longitude, as given in the section *Moon-culminating stars* in the *Nautical Almanac*.

In cases where a defective limb of a planet has been observed, a correction for defect of illumination (negative to the time of passage of the first limb, or positive to that of the second limb, according as the planet had passed or had not passed the opposition in right ascension) is applied in forming the concluding transit. These corrections are computed from the tables given for this purpose in the *Nautical Almanac*.

Occasionally the limb of Mercury is so ill-defined that the observer records a transit of the "centre." This is considered to be the centre of gravity of the illuminated portion of the disc, and a correction of $\frac{4}{3\pi}$ ($=0.424$) $\times$ defect of illumination should be applied for reduction to the centre of the actual disc. In practice this is applied in forming the concluded transit as $0.4 \times$ correction for defect of illumination in R.A., calculated as for a transit of a defective limb.

The *sixteenth* column contains the numerical values, in seconds of arc, of the errors of collimation, level, and azimuth; the last two being distinguished from the first by a round and a square bracket respectively. These errors will be treated separately as follows:—

Error of Collimation.

The value of this error is given in seconds of arc. This error is considered positive when it implies an additive correction to the time of observed transits of stars above the pole.

The method of determining the position of the line of collimation by means of the north and south collimators has been fully explained above on page vii. The reading of the micrometer of the north collimator, for the setting of its system of wires on the image of that of the south collimator, is ascertained every day, and the micrometer-screw of the north collimator is then left at that reading. The reading of the R.A. micrometer of the transit-circle, for coincidence of the central wire

xxxiv INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

with the image of the wire of each collimator, is then found; the mean of these two readings gives the reading for the line of collimation for the day; and the mean of these determinations, usually for a week, is adopted as the reading for the geometrical line of collimation of the transit-telescope. A correction of $+ 0^{\circ}.011$ has been applied to the "Adopted mean of group," to reduce the collimation-readings resulting from the observations of the collimator made through the central cube of the transit-circle to those resulting from observations made when the instrument is raised. (See *Introduction to Greenwich Second Nine-Year Catalogue* for 1900.)

During the year 1851 the observations of the collimators with the transit-circle telescope were usually made twice in every day. Upon examination of the results it did not appear that there was any discoverable difference at different hours of the day, and therefore generally, since that time, the observations have been made only once a day.

The R.A. micrometer is left at a fixed convenient reading near to that of the geometrical collimation, but not necessarily coinciding with it. The algebraical excess of the reading for collimation above this is the error of collimation, with the sign as explained above. To the excess thus converted into arc is then applied $-0''.19$ for diurnal aberration; and the result is the quantity which is given in the sixteenth column of the *Transits* as the error of collimation, and this is adopted for correction of the observed transits.

The numerical correction to the time of observed transit, in seconds of time, is,

$$\text{Error of collimation} \times \frac{1}{15 \sin N.P.D.}$$

Error of Level.

The error of level of the axis of revolution is considered positive when the western end of the axis is too high.

The process of determining the error of level consists in taking the reading of the transit-micrometer which makes the image of the middle wire as seen by reflexion from the surface of quicksilver coincide with the wire as seen directly, as is explained in the description of the instrument above, and in comparing this reading with the reading for the geometrical line of collimation. The excess of the reading for the line of collimation above the reading for coincidence of direct and reflected

images gives the error of level, with the sign mentioned above. The mean of the results is adopted for a longer or shorter time, as the instrument appears to have been more or less steady in level.

The result adopted for the error of level is given in the sixteenth column of the *Transits*, in curved brackets.

The numerical correction to the time of observed transit, in seconds of time, is,

$$\text{Error of level} \times \frac{\text{cosine zenith distance}}{15 \sin \text{N.P.D.}}$$

The error of level has usually been observed three times every day, and in some instances where there appeared to be a considerable change, the separate results have been used for the morning and evening observations respectively.

As a check on the general system of correction for error of level, transits of stars have been occasionally observed by reflexion, and the results have been compared with those of direct observations. A table exhibiting this comparison, together with the apparent correction to the error of level deduced from each star observed, is given at the end of this *Introduction*.

Error of Azimuth.

This error is considered positive when the eastern pivot is too far north. It is determined from observations of the following stars :—Polaris, Cephei 51, Groombridge 1119, Bradley 1672, Groombridge 2283, δ Ursæ Minoris, λ Ursæ Minoris, Groombridge 3548, and Bradley 3147.

The error actually investigated is the error at the pole, usually denoted by the letter n , and not the error at the horizon or azimuth. Each of the nine circumpolar stars that may have been observed is reduced separately, and the following process is used :—The observation of the circumpolar star and a clock star (the nearest in right ascension that has been observed) being corrected for “reduction to centre wire” and for collimation error (but not level error), and denoting by m the instrumental errors at the equator and by E the clock error, then for each star—corrected time of transit $+ \frac{1}{15} n \cot \text{N.P.D.}$ $+ m + E$ = apparent tabular right ascension for the day, and hence by a subtraction it follows that n multiplied by the difference of value of the factor $\frac{1}{15} \cot \text{N.P.D.}$ for the

xxxvi INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

two stars is equal to the difference for the two stars of the excess of the tabular R.A. over the corrected time of transit. On a few rare occasions it is necessary to apply a correction for the change in clock error between the two observations. The level and n being thus found from observation and their values being adopted by taking means of the groups into which the observed values appear to fall, the azimuth error and m are found by the formulæ

$$\begin{aligned}\text{Azimuth error} &= -n \operatorname{cosec.} \operatorname{colat.} + l \cot \operatorname{colat.} \\ 15 m &= -n \cot \operatorname{colat.} + l \operatorname{cosec.} \operatorname{colat.}\end{aligned}$$

The corrections are applied by the formula $\frac{1}{15} n \cot \text{N.P.D.} + m$.

A table of factors for the three errors, having for argument the N.P.D. of the object, which may be found convenient for the verification of the computed corrections, is given in the *Transit-Circle Tables* at the end of this *Introduction*.

The *seventeenth* column contains the seconds of every transit, corrected for the three preceding errors, and is conceived to represent the clock-time at which each body passed the true astronomical meridian of Greenwich. The numbers to which a bracket is annexed are those resulting from the mean of the two limbs of the Sun or a planet.

The *eighteenth* column contains the seconds of the tabular R.A. of the stars which are used for determining clock and azimuth errors; the latter are given in order to enable the reader to judge of the general state of adjustment of the instrument, but not to assist in determining clock-errors.

From the beginning of 1906, the right ascensions are deduced from the *Standard Mean Right Ascensions for 1900.0 based on 12-hour groups*, printed in the Introduction to the *Second Nine-Year Catalogue for 1900*. The values of the precession and proper motions are taken from Prof. Newcomb's *Catalogue of Fundamental Stars*. The corrections necessary to bring up the stars' right ascensions to the day of observation are taken from the *Nautical Almanac*. In the *Transit-Circle Tables*, at the end of this *Introduction*, are given the adopted mean right ascensions for 1908.0 of all the stars which have been used for determining clock-errors or azimuth-errors, together with the corrections to the mean right ascensions given in the *Nautical Almanac*.

The *nineteenth* column contains the error of the clock, found by subtracting

the numbers in the seventeenth column from those in the eighteenth. The error is therefore positive when the clock is slow. The apparent clock-errors given by the nine close polar stars used for azimuth-error, as has been remarked above, are not employed in determining the clock-errors available for computations of right ascensions.

The *twentieth* column contains the adopted losing rate and the adopted error of the clock at 0^h sidereal, as found by the following process. The observations are divided into groups, defined by bars across this column (in all instances the same as the limits of the observations of each observer). The mean of the clock-errors given by all the clock-stars in each group is then taken, and this is considered to be the clock-error corresponding to the mean of all the times of transit of those stars. For the determination of clock-rate, the clock errors found by each observer are corrected for that observer's personal equation, Mr. Witchell being taken as the Standard Observer. The values of the personal equation are derived from the observations of the previous year. Each mean clock-error, properly modified, is now compared with that which precedes and with that which follows; a preceding rate and a following rate are thus found, and by these the Chief Assistant is guided in adopting the clock-rate to be used through the group of observations: this adopted clock-rate is set down in the twentieth column. For facility of calculation, the proportional part of this rate, corresponding to the mean sidereal time or mean of all the times of transit for the group, is applied with changed sign to the mean clock-error; and thus the clock-error at 0^h sidereal is found. If it happen that any observations included in the same group are made in the sidereal day preceding or following that for which the mean error has been found, the whole adopted daily rate is subtracted from or added to the error found for 0^h; and thus the error for 0^h of the preceding or the following sidereal day is obtained. These errors at 0^h for all the different sidereal days are also set down in the twentieth column. It is to be remarked that, in the application of these errors, if the observations of small stars and planets have been made by the same person as the observations of the clock-stars, no further computation is necessary for obtaining the right ascensions of those objects than the application, to the time of transit, of the clock-error at 0^h, and of a proportional part of the clock-rate corresponding to the right ascension of the object; but, if the clock-stars are observed by one observer, and the other object by another, a correction for personal equation is required. The personal equations are *not* applied in the column of *Apparent Right Ascensions from the observations* in this section.

The observations which are available in the year 1908 for determining personal equations in the use of the chronographic method of recording transits are discussed

two stars is equal to the difference for the two stars of the excess of the tabular R.A. over the corrected time of transit. On a few rare occasions it is necessary to apply a correction for the change in clock error between the two observations. The level and n being thus found from observation and their values being adopted by taking means of the groups into which the observed values appear to fall, the azimuth error and m are found by the formulæ

$$\begin{aligned}\text{Azimuth error} &= -n \operatorname{cosec.} \operatorname{colat.} + l \cot \operatorname{colat.} \\ 15 \ m &= -n \cot \operatorname{colat.} + l \operatorname{cosec.} \operatorname{colat.}\end{aligned}$$

The corrections are applied by the formula $\frac{1}{15} n \cot \text{N.P.D.} + m$.

A table of factors for the three errors, having for argument the N.P.D. of the object, which may be found convenient for the verification of the computed corrections, is given in the *Transit-Circle Tables* at the end of this *Introduction*.

The *seventeenth* column contains the seconds of every transit, corrected for the three preceding errors, and is conceived to represent the clock-time at which each body passed the true astronomical meridian of Greenwich. The numbers to which a bracket is annexed are those resulting from the mean of the two limbs of the Sun or a planet.

The *eighteenth* column contains the seconds of the tabular R.A. of the stars which are used for determining clock and azimuth errors; the latter are given in order to enable the reader to judge of the general state of adjustment of the instrument, but not to assist in determining clock-errors.

From the beginning of 1906, the right ascensions are deduced from the *Standard Mean Right Ascensions for 1900.0 based on 12-hour groups*, printed in the Introduction to the *Second Nine-Year Catalogue for 1900*. The values of the precession and proper motions are taken from Prof. Newcomb's *Catalogue of Fundamental Stars*. The corrections necessary to bring up the stars' right ascensions to the day of observation are taken from the *Nautical Almanac*. In the *Transit-Circle Tables*, at the end of this *Introduction*, are given the adopted mean right ascensions for 1908.0 of all the stars which have been used for determining clock-errors or azimuth-errors, together with the corrections to the mean right ascensions given in the *Nautical Almanac*.

The *nineteenth* column contains the error of the clock, found by subtracting

the numbers in the seventeenth column from those in the eighteenth. The error is therefore positive when the clock is slow. The apparent clock-errors given by the nine close polar stars used for azimuth-error, as has been remarked above, are not employed in determining the clock-errors available for computations of right ascensions.

The *twentieth* column contains the adopted losing rate and the adopted error of the clock at 0^h sidereal, as found by the following process. The observations are divided into groups, defined by bars across this column (in all instances the same as the limits of the observations of each observer). The mean of the clock-errors given by all the clock-stars in each group is then taken, and this is considered to be the clock-error corresponding to the mean of all the times of transit of those stars. For the determination of clock-rate, the clock errors found by each observer are corrected for that observer's personal equation, Mr. Witchell being taken as the Standard Observer. The values of the personal equation are derived from the observations of the previous year. Each mean clock-error, properly modified, is now compared with that which precedes and with that which follows; a preceding rate and a following rate are thus found, and by these the Chief Assistant is guided in adopting the clock-rate to be used through the group of observations: this adopted clock-rate is set down in the twentieth column. For facility of calculation, the proportional part of this rate, corresponding to the mean sidereal time or mean of all the times of transit for the group, is applied with changed sign to the mean clock-error; and thus the clock-error at 0^h sidereal is found. If it happen that any observations included in the same group are made in the sidereal day preceding or following that for which the mean error has been found, the whole adopted daily rate is subtracted from or added to the error found for 0^h; and thus the error for 0^h of the preceding or the following sidereal day is obtained. These errors at 0^h for all the different sidereal days are also set down in the twentieth column. It is to be remarked that, in the application of these errors, if the observations of small stars and planets have been made by the same person as the observations of the clock-stars, no further computation is necessary for obtaining the right ascensions of those objects than the application, to the time of transit, of the clock-error at 0^h, and of a proportional part of the clock-rate corresponding to the right ascension of the object; but, if the clock-stars are observed by one observer, and the other object by another, a correction for personal equation is required. The personal equations are *not* applied in the column of *Apparent Right Ascensions from the observations* in this section.

The observations which are available in the year 1908 for determining personal equations in the use of the chronographic method of recording transits are discussed

xxxviii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

in the *Transit-Circle Tables* at the end of this *Introduction*. They have been treated in the same manner as in preceding years, by comparing the clock-error at the same sidereal hour given by two different groups of stars observed by different persons. These personal equations have been applied to the collected results of transits for the Sun, Moon, and planets, whenever they were needed throughout the year 1908. But those used in the adoption of clock-rate as explained on page xxxvii, referred to Mr. Witchell as standard, have been deduced from the observations in 1907. The investigation of these is given in the *Transit-Circle Tables* for last year.

It has been the practice on favourable nights for the transit-circle observer to observe two "clock stars" by the eye-and-ear method, at the same time making a careful comparison between the clock "Hardy" (used in observing by this method) and the Sidereal Standard, which is used in chronographic transits. It is then an easy matter to compare the values of the quantity "clock slow" as found by the same observer using the different methods. The discussion of these results is given in the *Transit-Circle Tables*.

In the standing footnote on each page is given the duration of passage for the semidiameter of a planet when only one limb has been observed. It is found as follows :—

For the Sun, the following corrections, depending on the respective observers, are applied to the Nautical Almanac semidiameter :—

Mr. Eddington	$-0^{\circ}06_{10}^s$	Mr. Cullen	$-0^{\circ}02_{22}$
Mr. Witchell	$-0^{\circ}01_{26}$	Mr. Daniels	$-0^{\circ}01_7$
Mr. Storey	$-0^{\circ}12_7$	Mr. Shepperd	$+0^{\circ}04_6$
Mr. Evans	$+0^{\circ}03_{22}$	Mr. Acton	$-0^{\circ}02_8$

The subscript numbers denote the number of observations on which the respective corrections depend.

These corrections have been determined from the observations of the respective observers during the current year.

For the Moon and Mercury, the duration is taken without alteration from the *Nautical Almanac*.

For Mösting A the correction to reduce to the centre of the Moon has been obtained by interpolation with second differences from the *Berliner Jahrbuch*.

For Venus, a correction is applied to the Nautical Almanac semidiameter, determined from observations made with the transit-circle from 1851 to 1862. The investigation by Mr. Stone will be found in the *Monthly Notices of the Royal Astronomical Society*, vol. xxv. No. 3. The correction, which has been applied additively to the Nautical Almanac semidiameter during the year, is:—

$$+ 0\cdot026 + 0\cdot027 \times \text{tabular duration of passage of semidiameter.}$$

For Jupiter and Saturn, both limbs are always observed, if possible; but if, through accident or necessity, only one limb is observed, the duration of passage of the semidiameter is obtained by applying to the value given in the *Nautical Almanac* a correction derived from a few of the neighbouring observations. The elements of the computation are given in the notes.

The *twenty-first* column contains the right ascension of the centre of the body observed: it is formed by adding the time from the seventeenth column, the clock-error at 0^h next preceding from the twentieth column, the proportional part of the rate in the same column corresponding to the right ascension, and the duration of the passage of the semidiameter from the footnotes. No result is set down for a clock-star, unless at least six clock-stars, distributed over six hours of right ascension, have been observed: no result is set down for an azimuth star unless at least three of the nine azimuth stars have been observed above pole, and three below pole, and no result is set down for other close circumpolar stars, unless there has been a good determination of the azimuthal error on the day of observation.

Whenever the concluded result in this or other sections is obviously erroneous, it is enclosed within brackets, and is not used in subsequent computations.

§ 2. *Determinations of the Errors of Collimation, Level, and Azimuth of the Transit-Circle.*

With regard to these tables, the following remarks, in addition to the explanation which has been given, in the description of the instrument, and in the account of the methods of determining the errors of collimation and level, will be sufficient.

OBSERVATIONS FOR COINCIDENCE OF COLLIMATORS.

These observations are made on every week-day. On all days the view is taken through the pierced cube of the transit-circle. On Mondays the transit-circle is raised, and observations are taken with an uninterrupted view of the S. collimator by the N. collimator as well as through the cube.

xl INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Immediately before the observations of the collimators with the transit-circle for determination of collimation-error, six readings of the micrometer of the N. collimator, for setting its wire system on the wire system of the S. collimator, are taken, and the wire is then left at a reading corresponding to the mean of the six. Since the alteration in the form of mounting of the collimators, referred to on page vii, this observation has been usually repeated immediately after the observations of the collimators with the transit-circle. The means of both sets of micrometer-readings are printed in the table, and will give a satisfactory idea of the firmness of the collimators.

OBSERVATIONS OF THE COLLIMATORS FOR DETERMINATION OF THE ERROR
OF COLLIMATION.

The way in which these observations are made has been fully explained (pages vii and xxxiii). The means of the readings for the separate collimators are set down in the second and third columns. The mean of these gives, after the application of a correction " $+ 0.16 \times$ excess of first reading over second reading of N. collimator for coincidence with S. collimator," the concluded reading set down in the fourth column. The next column gives the reading which has been adopted for the period of time limited by the bars above and below; and the last column gives the adopted reading for the line of collimation after the application of a correction of $+ 0.011$ to reduce the result to that which would have been obtained if the instrument had been raised when the observation for coincidence of the collimators was made. By comparing the numbers in this column with the *Readings of Transit-Micrometer* given in the footnotes of the section *Transits observed, &c.*, and applying $- 0.19$ for diurnal aberration, the errors of collimation actually used will be deduced. For determination of the sign of the error, it must be observed that the micrometer-head is east, and that the readings of the micrometer increase as the wire moves towards the head.

Since the reversion-prism was applied in 1884 June, it was the practice for a number of years, in each set of observations for coincidence of the collimators with each other, and of the transit-circle with either of them, to take three readings with the direction for increasing readings apparently towards the right, and three with the direction apparently towards the left, in order to eliminate any personality depending on the apparent direction of measurement. A series of observations for the investigation of this personality was made by different observers, with the direction for increasing readings apparently towards the right, towards the left, downwards and upwards

respectively. The discussion of these observations, given in the *Transit-Circle Tables* for the year 1885, shows that the apparent direction of measurement has no appreciable influence on the results.

DETERMINATION OF THE LEVEL-ERROR.

The method of making these observations has been described (pp. ix and xxxiv). A comparison of the transit-micrometer reading for coincidence of the direct and reflected images of the central wire (printed in the second column) with the adopted reading for the line of collimation gives the error of level in terms of revolutions of the micrometer; and the numbers in the third column exhibit the errors of level in arc. The last column gives the adopted values of the error and the limits of time during which they have been used. The determination of the sign of the errors has been previously explained.

DETERMINATION OF THE ERROR OF AZIMUTH.

The method of finding the error of azimuth has been fully explained on p. xxxv. The first three columns require no explanation. In the fourth column is given the value of $\frac{1}{15} \cot N.P.D.$ for each polar and clock-star. The difference of the two quantities in the third column divided by the difference of the corresponding quantities in the fourth column is the value of n given in the fifth column. In the sixth column is given the adopted value, usually the mean of a number of separate determinations. The resulting error of azimuth in the seventh column is found from the adopted values of the level and n by the formula.

$$\text{Error of Azimuth} = \text{Error of level} \times \cot. \text{ colat.}, - n \operatorname{cosec} \text{ colat.}$$

§ 3. *Mean Errors and Adopted Rates of the Sidereal Standard Clock.*

In this section are given the errors of the Sidereal Standard Clock for mean midnight as deduced from the observations of each observer. The adopted losing rate (found as explained on p. xxxvii.) is also given.

§ 4. *Observations of Zenith Distance with the Transit-Circle.*

The observations here printed in detail are those of the Sun, Moon, planets, fundamental and adjustment stars. Commencing with the year 1897, the printing of the details of observations of other stars has been discontinued, the daily results only being

GREENWICH OBSERVATIONS, 1908.

f

given in the *Ledgers of Mean Right Ascensions and Mean North Polar Distances of Stars*.

In order to include the whole of the reductions belonging to a single observation on one page, the elements for those reductions which are not frequently changed, and which do not require tabular arrangement, are given in footnotes or notes at the beginning of the section. These are :—correction for runs; formulæ used in computing micrometer-corrections, corrections for curvature of path, and inclination of wire; and zenith-point corrections.

The corrections for planets, including change of N.P.D. for reduction to the meridian, parallax, and semidiameter, are exhibited in tabular arrangement at the lower part of each page, opposite to the reference numbers which correspond to the observations in the upper part of the page.

DAY.—The day of observation always begins with the Sun's transit.

NO. FOR REFERENCE.—This column requires no explanation.

NAME OF OBJECT.—The nomenclature of stars follows the same general rule as in the section of *Transits observed*, &c. The letters N. L. and S. L. denote north and south limbs of planets, and N. C. and S. C. the north and south cusps of Mercury or Venus. Mösting A denotes the crater of that name on the Moon. The words Wire R indicate an observation of the image of the horizontal or zenith-distance wire as seen by reflexion in a trough of mercury, as is described in page xii. For other objects, the letter R. denotes that the object is observed by reflexion in the trough of mercury.

READINGS OF THE MICROSCOPE-MICROMETERS.—The application of the letters for the six ordinary microscope-micrometers is the same as in the observations of preceding years: the observer, beginning with the northern horizontal microscope, and passing the upper microscopes, the southern horizontal, and the lower ones, in the direction of the circle's circumference, reads them in the order A, C, E, B, D, F (see diagram, page x). These readings in the observing books are placed in the order $\begin{smallmatrix} A, C, E \\ B, D, F \end{smallmatrix}$; from which form they are easily changed into the order of printing. Each pair of adjacent readings, therefore, is the pair of readings at the opposite ends of a diameter. The number of integral revolutions is given in the fifth column only.

Occasionally the number of revolutions expressed in this column does not apply unchanged to all the microscopes, on account of the difference of their readings; but since the readings never differ by more than three or four tenths of a revolution, this will occasion no ambiguity. On a few occasions the cross-wires have fallen on the negative side of the micrometer-zero; in such cases (which are always mentioned in the notes) the fractional part of the reading given is positive, being that taken from the micrometer-head, but the integer 9 represents -1 . The reading of the pointer-microscope is omitted for the sake of economy of space, but it may be inferred from the zenith-point given in full at the bottom of each page, and from the deduced zenith-distance.

It is proper now to explain how the revolutions and parts of revolutions of the microscope-micrometers are converted into arc, when the six ordinary microscopes are used.

First, it may be premised that, as one revolution of each microscope-micrometer does not differ extravagantly from $1'$ on the limb of the circle, we may consider each revolution as a nominal minute. Next, supposing the number of integral revolutions, as shown by each microscope, to be the same for all, we ought, in order to obtain the mean of the fractions of a revolution, to add together the subdivisions as shown by the different micrometer-heads, and divide the sum by 6. Thirdly, as the subdivisions are in the decimal scale, we should then multiply this mean by 60, to reduce it to sexagesimal seconds. It is evident thus that the number of nominal seconds to be attached to the nominal minute will be found by simply adding together the subdivisions on the micrometer-heads, and shifting the decimal point.

From the ordinary observations of runs it is known that the screws of the micrometers are so sensibly equal that, in the reduction of the observations, it is sufficient to take the mean of the six readings, and to apply to it the mean correction for runs. To do this, the following process is employed. The sum of the runs for the six micrometers never differs materially from $29^{\circ}3$. Now, if the correction, by the addition of which each revolution of a micrometer may be converted into a minute of arc, were exactly $\frac{1}{51}$ part of the reading, then the sum of the runs would be $\frac{50}{51} \times 30^{\circ}000$, or $29^{\circ}4118$. If the sum of the runs is not exactly equal to this quantity, there will be a variable correction after adding $\frac{1}{51}$ part of the mean of the readings of the microscopes. To determine this quantity, let the sum of the runs be $29^{\circ}412 - x$. Then the true reading in seconds of space, corresponding to a nominal

reading of r'' (including the value both of the nominal minutes and of the subdivisions of the nominal minute) will be

$$r'' \times \frac{30.000}{29.412 - x}$$

or

$$r'' \left\{ 1 + \frac{1}{50} + 0.03468 \times x \right\}$$

Hence the correction, after adding $\frac{1}{50}$ part of the mean of readings, will be $+ r x \times 0''.03468$. When the four supplementary microscopes are used these rules are modified.

CORRECTIONS FOR ERROR OF DIVISION.—The corrections for errors of division are the means of the determinations made in 1856, 1871, and 1898 for every 5° , and of those made in 1851, 1856, and 1898 for every 1° . They are printed in the *Transit Circle Tables*.

FLEXURE AND R—D DISCORDANCE.—The observations for determination of the horizontal astronomical flexure of the transit-circle telescope are given in the *Transit-Circle Tables* at the end of this *Introduction*. In making these observations, the nearly horizontal wire of the south collimator was brought upon the image of the corresponding wire of the north collimator by looking through the cube of the transit-circle; and the micrometer-head was set at the mean of 10 successive readings. After each complete set of 5 observations of the collimating wires, the observations for coincidence of these wires were repeated; and a correction has been applied, when required, to the mean circle-reading at bisection of the wire of the south collimator, for the half-difference in the two sets of observations for coincidence.

The investigation of the discordance of direct and reflexion observations is given in the *Transit-Circle Tables*.

From 1906 January 1, a constant flexure term of $+0''.60 \sin Z.D.$ has been adopted to include the effects of astronomical flexure and the discordance of direct and reflexion observations; but this correction is not applied in this section or in the *Ledgers*, being deferred until the results are collected in the *Catalogue* (§ 7).

To obtain information as to the cause of the R—D discordance, observations were made between 1893 December 3 and 1894 April 22 of the zenith-distances of stars directly and by reflexion, the direct observation being taken first. The observations were in other respects similar to the regular reflexion observations. These observations are printed in the *Transit-Circle Tables* for 1894 (pp. cxi-i), and a discussion of

TRANSIT-CIRCLE.—READINGS OF ZENITH-DISTANCE MICROMETER : VERTICAL WIRES. xlv

them is given in a paper by Prof. Turner and Mr. Thackeray in the *Memoirs of the Royal Astronomical Society*, vol. li.

During the years 1895 to 1898 a new series of observations were made with a view to obtaining further information on the cause of the R—D discordance. Pairs of stars were chosen of nearly the same zenith-distance and a few minutes apart in right ascension. One star of each pair was observed by reflexion, the other directly at the same transit; on the next night of observation the order was reversed, the star which had been observed previously by reflexion being observed directly. In this way a value of 2 (R—D) was derived. In order to test whether any change was taking place in the instrument, two readings of the circle were taken, one just before the star entered the field, and the second just after the bisections with the zenith-distance micrometer had been made. The results of the observations are given in the *Transit-Circle Tables* for the years 1895, 1896, 1897, and 1898, together with a comparison with the R—D discordance obtained in the usual way.

READINGS OF ZENITH-DISTANCE MICROMETER.—This column contains the reading of the micrometer on the eye-end of the telescope, corresponding to the several bisections of an object during its passage across the field of view. The instrument is furnished with two clamps, one north and the other south, for fixing it, but with no slow-motion screw; the object must therefore be bisected by means of the micrometer-screw, and the micrometer must be read for every observation. The reading of 20^r corresponds pretty nearly to the middle of the range of the screw, and the wire is then tolerably near the centre of the field of view; observations are, therefore, generally made with a position of the wire not greatly different from this.

In 1906 July a new steel screw was applied to the micrometer by Messrs. Troughton & Simms. Observations for the determination of the value of 1^{rev.} of this screw are given in the *Transit-Circle Tables* for 1906, the resulting value being 36[″].93. The errors of the screw were also investigated and found to be insensible.

VERTICAL WIRES IN THE TELESCOPE-FIELD AT WHICH CIRCLE-OBSERVATIONS ARE TAKEN.—Omitting, in this section, all reference to the wires P, Q, R, S, U, V, W, adapted to galvanic observation of transits, the telescope contains seven vertical or transit wires M, N, O, T, X, Y, Z, placed at sensibly equal intervals of about 14[°].8 for transit of an equatorial star; these wires are here called 1, 2, 3, 4, 5, 6, 7, in the case of a star observed above pole, the numbers being reversed for a star observed S. P. The central or 4th wire is so nearly in the meridian as to prevent the necessity of any correction to the observed circle-reading when the body is observed at the

passage over that wire. In that case, the only corrections which are required to the mean of the microscope-micrometer-readings (when converted into arc and combined with the equivalent for the zenith-distance micrometer-reading) are those arising from errors of division of the graduated circle, and from astronomical flexure of the telescope.

When the object has not been observed at the central wire, corrections are required for reduction to the meridian. These consist of three, viz. :—First, for want of horizontality of the micrometer-wire; secondly, for curvature of path of the object, or for difference between the small circle described by the body and the great circle of which the horizontal wire forms a part; and thirdly, in the case of the Sun, Moon and planets, for the change of N.P.D. in the interval between the time of observation and the meridian passage.

The correction due to inclination of wire was found by bisections of stars (by the zenith-distance micrometer) at the first and seventh, or second and sixth, vertical wires. Its numerical value, additive before transit and subtractive after transit, for stars above the pole, is given in the notes at the beginning of the section.

The general formula for the correction for curvature is given also in the notes at the beginning; its value is easily interpolated from a table constructed for convenient intervals of N.P.D. In the case of stars very near the pole, when the place of observation does not coincide with a vertical wire, the correction for curvature is computed from the formula :—

Correction for curvature = number ($\log. = 6.43569$) $\times \sin 2$ N.P.D. $\times t^2$, t being the interval between the time of meridian passage and the time of transit expressed in seconds of time.

The formulæ for correction for motion are the following: In the case of the Moon, the correction for the change of N.P.D. in passing from one wire to the next is computed from the formula, "Change of declination for one hour of terrestrial longitude (from the section of Moon-culminating stars in the *Nautical Almanac*) $\times \sec. \text{declination} \times \frac{\sin \text{Geoc. Z.D.}}{\sin \text{App. Z.D.}} \times \frac{14.8}{3600}$ (the logarithm of the last factor being 7.61396)." In the case of the other planets, the correction for motion is deduced immediately from the hourly motion in declination given in the *Nautical Almanac*; or, in the case of newly discovered planets, from their peculiar ephemerides.

SECONDS OF CONCLUDED MERIDIAN CIRCLE-READING.—The numbers in this column are found by adding together the mean of the microscope-readings corrected for

runs, the correction for division error, the equivalent for the mean of the zenith-distance micrometer readings, corrected for curvature and inclination of the wire, and the correction for motion in the case of the Sun, Moon, or planets. The degrees and minutes may be ascertained, when required, by subtracting the "Correction for zenith-point" given in the notes from the apparent zenith-distance.

ZENITH-POINT CORRECTION.—The observations for this purpose are those of the reflected image of the horizontal wire, and those of the direct and reflected images of stars made at the same transit.

From 1886 January 1 to 1896 December 31 no correction was applied to nadir-observations for discordance between the results of this observation and the mean of reflexion observations of north and south stars. From 1897 January 1 a correction of $-0''.25$ has been applied to the nadir-observations in forming the adopted zenith-point, when there are no reflexion observations of north and south stars; but when the observation of nadir is combined with reflexion observations of stars no correction is applied. From observations in former years a connexion was traced between this discordance and the errors of the screws of the microscope-micrometers, and it was hoped that the application of the new steel screws (one of each pair being reversed) would remove the discordance. This was found to be the case till 1892, the mean apparent correction to the nadir-observation as found from reflexion observations of stars since 1886 being as follows:—

1886	— 0.09	1894	— 0.27	1902	— 0.20
1887	+ 0.01	1895	— 0.31	1903	— 0.23
1888	— 0.12	1896	— 0.34	1904	— 0.41
1889	+ 0.07	1897	— 0.27	1905	— 0.22
1890	+ 0.08	1898	— 0.36	1906	— 0.40
1891	+ 0.07	1899	— 0.41	1907	— 0.24
1892	— 0.26	1900	— 0.39	1908	— 0.33
1893	— 0.34	1901	— 0.29		

In consequence of the discordance shown in 1892, the errors of the microscope-micrometer-screws were re-determined in 1893. The observations on which these depend are given in the *Transit-Circle Tables* for 1894. Although signs of wear were shown, the errors arising from this cause are so nearly eliminated by the reversal of three of the screws, that it has not been considered advisable to apply any correction to the circle-reading for apparent error of the screws. Similar observations made in 1896 and 1906 confirm these results. These observations are given in the *Transit-Circle Tables* for 1896 and 1906.

With a view of obtaining information as to the cause of this discordance, observations of the nadir have been taken since 1895 July three times each day when practicable.

The following differences were found from observations since 1895 :—

	2 ^h .—3 ^h . G.M.T.	3 ^h .—9 ^h . G.M.T.	9 ^h .—15 ^h . G.M.T.	No. of Days.
	"	"	"	
1895	+ 0.29	0.00	+ 0.27	54
1896	+ 0.20	0.00	+ 0.16	84
1897	+ 0.11	0.00	+ 0.17	108
1898	+ 0.16	0.00	+ 0.11	104
1899	+ 0.17	0.00	+ 0.19	152
1900	+ 0.14	0.00	+ 0.05	118
1901	+ 0.17	0.00	+ 0.09	123
1902	+ 0.20	0.00	+ 0.17	97
1903	+ 0.04	0.00	+ 0.09	103
1904	+ 0.06	0.00	+ 0.15	79
1905	+ 0.13	0.00	+ 0.04	74
1906	+ 0.19	0.00	— 0.01	40
1907	+ 0.17	0.00	+ 0.14	67
1908	+ 0.11	0.00	+ 0.05	82

The + sign indicates a greater circle-reading.

The details of the computation of zenith-point are given in a subsequent table. The adopted corrections for zenith-point are given in the footnotes to this section.

From an investigation made in 1903, with the observations of 1901, it appears that the probable error of a complete determination of zenith-point (four stars and three nadir observations) is about $\pm 0''.11$.

APPARENT ZENITH-DISTANCE.—The numbers in this column are found by adding the Zenith-point correction to the Concluded meridian circle-reading; when the sum exceeds 270° (denoting that the object is between the zenith and the north horizon), the difference from 360° is taken, and the negative sign is attached to it; for stars observed by reflexion, the sum is subtracted from 180° , and the remainder, with its algebraical sign, is the apparent zenith-distance.

When both limbs of the Moon are observed, it usually happens that one of them is slightly defective from the want of illumination by the Sun. The necessary correction (which is not applied in this column, but which must be applied with others in order to form the numbers in the column of *Geocentric N.P.D. of Centre*) is ascertained by computation in the following manner:—When the Moon is horned, let θ be the angle which the great circle joining the cusps makes with the meridian; δ_s the Sun's declination, δ_m the Moon's declination as seen at Greenwich; P the Sun's hour-angle at the Moon's transit: then

$$\tan \theta = \operatorname{Cosec} P. \cos \delta_m. \tan \delta_s. - \cot P. \sin \delta_m;$$

TRANSIT-CIRCLE.—APPARENT ZENITH-DISTANCE: CORRECTION FOR DEFECTIVE ILLUMINATION OF MOON AND PLANETS. xlix

and the correction to be applied to the Geocentric N.P.D. will be

$$\text{Moon's geocentric semidiameter} \times \text{versin } \theta.$$

For Mercury and Venus, when horned, the investigation of the correction for defective illumination assumes the same form as for the Moon.

If the Moon be gibbous, as is the case in the greater number of instances, draw a great circle through the centre of the Moon, at right angles to the meridian passing through the Moon, and let it meet the meridian passing through the Sun; then the intersection of this circle with the Sun's meridian will determine the place of a fictitious Sun, which would equally illuminate both limbs of the Moon when on the meridian; and the elevation or depression of the true Sun above or below the great circle joining the Moon and fictitious Sun, measured in a plane at right angles to that circle, will represent the angle by which the lowest or highest part of the illuminated hemisphere is distant from the limb, and therefore the angle whose versed sine multiplied into the Moon's semidiameter is the correction required.

Let, then, P be the north pole of the heavens, M the Moon on the meridian of Greenwich, and S and S_1 the true and fictitious Suns, and imagine triangles formed by great circles joining these. Let also δ_1 be the declination of the fictitious Sun, θ_1 the arc joining the true and fictitious Suns, and θ the perpendicular arc before mentioned.

$$\text{Then} \quad \tan \delta_1 = \tan \delta_m \cdot \cos MP S_1.$$

$$\text{And} \quad \theta_1 = \delta_s - \delta_1.$$

$$\begin{aligned} \text{Also} \quad \sin \theta &= \sin \theta_1 \cdot \sin MS_1 P. \\ &= \sin \theta_1 \frac{\sin MP}{\sin S_1 P} \\ &= \sin \theta_1 \frac{\cos \delta_m}{\cos \delta_1}. \end{aligned}$$

From which θ is found.

The correction required to Geocentric N.P.D. is

$$\text{Moon's geocentric semidiameter} \times \text{versin } \theta.$$

In either case the North limb is fully illuminated if θ is positive, and the South limb if θ is negative.

GREENWICH OBSERVATIONS, 1908.

g

1 INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

In the cases of Mercury and Venus when gibbous, or of Mars and other planets sensibly gibbous, the investigation takes the following form. The angle θ being found as above, the length of the straight line drawn from the Sun perpendicular to the plane represented by the great circle above mentioned is equal to Distance of Earth from Sun $\times \sin \theta$; and therefore if ϕ be the apparent angular distance of the Sun from that plane as seen from the planet, the value of ϕ is determined by the equation

$$\sin \phi = \frac{\text{Length of perpendicular line}}{\text{Dist. of Planet from Sun}} = \frac{\text{Dist. of Earth from Sun}}{\text{Dist. of Planet from Sun}} \times \sin \theta.$$

The angle ϕ represents in this case the angle by which the lowest or highest point of the illuminated hemisphere is distant from the limb as viewed by an observer anywhere in the plane of this great circle, and the correction therefore will be

$$\text{Planet's semidiameter} \times \text{versin } \phi.$$

In cases where the "centre" of Mercury is observed, a correction of $0.4 \times$ ordinary correction for defect of illumination in N.P.D. is applied for reduction to the actual centre of the disc. See page xxxiii.

These corrections for defective illumination of the Moon and tables for calculating the corrections for defective illumination of the planets are now given in the *Nautical Almanac*.

BAROMETER.—The barometer is by Horne and Thornthwaite, marked "No. 389," with metallic scale and graduation modified for change of mercury-surface in the cistern. Its scale is divided by vernier to 0.002 inch; but it is not usually necessary to read it to a smaller quantity than 0.01 inch for the computation of refraction. This barometer was substituted in April 1879 for the barometer by Newman, which had been in use to that date. From comparisons with the Standard Barometer made between 1879 March 19 and May 6 (allowance being made for the difference of temperature of the two barometers), it appears that the correction required to the barometer "Horne and Thornthwaite, No. 389," is only -0.007 inch. No correction has been applied to the readings of this barometer.

EXTERIOR THERMOMETER.—From 1897 January 1 the thermometer Negretti and Zambra, No. 70661, has been used as the Exterior Thermometer for all observations with the Transit-circle. This thermometer was mounted in the front court in a screen allowing a free circulation of air on 1896 July 18.

The following are the results of the latest comparisons with the Standard Thermometer which agree well with previous determinations.

1906 May 16.

Temperature.	Error.
32°·2	+ 0°·2
50°·2	+ 0°·2
66°·3	+ 0°·3
73°·9	+ 0°·3
86°·6	+ 0°·2
95°·7	+ 0°·3

A correction $-0^{\circ}\cdot 2$ has been applied to the exterior thermometer from 1897 January 1.

INTERIOR THERMOMETER.—This thermometer is No. 12264, by Negretti and Zambra.

The following are the results of comparisons with the Standard Thermometer made on 1906 May 16.

Temperature.	Error.
32°·6	+ 0°·6
50°·7	+ 0°·7
64°·1	+ 0°·8
70°·3	+ 0°·9
87°·2	+ 0°·8
95°·9	+ 0°·5

A correction of $-0^{\circ}\cdot 5$ has been applied to the readings of the Interior Thermometer from 1889 March 15 to 1894 December 31, and of $-0^{\circ}\cdot 6$ from 1895 January 1.

Previous to its removal to the new Magnetic Pavilion the Meteorological Standard Thermometer was read at noon by the Transit-circle observer for comparison with the Exterior Thermometer. Since 1899 July 12, readings of this thermometer have been made each day at Apparent Noon as well as Mean Noon by the Meteorological observer. A comparison of the readings of the Exterior Thermometer and the Meteorological Standard is given in the *Transit-Circle Tables*.

REFRACTION.—From the beginning of 1906, Pulkowa refractions have been used. Tables of these refractions are given in an Appendix to the *Greenwich Observations*, 1898.

lii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

PARALLAX.—The corrections for parallax of the Sun, Moon, and planets actually employed are given in the lower part of the page, as previously stated; the assumed ellipticity of the Earth being $\frac{1}{300}$ and the value of the solar parallax $8''\cdot80$.

For the planets the formula is :—

$$\log. \text{ parallax} = \log. \sin (\text{Z.D.} - 11'.12'') + \text{ar. co. log. distance} + 0.94360.$$

For the Moon, the horizontal equatorial parallaxes are interpolated, without alteration, from the *Nautical Almanac*, for the time of observation. The formula employed for the computation of the parallax to be applied to the observed zenith-distance is :—

$$\text{Parallax} = \text{Hor. Equat. Par. from N.A.} \times \text{Sin Dist. from Geoc. Zenith} \times \text{Geocentric Radius (Log. = 9.9991136)} + \text{Corr. derived from the following table :—}$$

Z.D.	Correction for South Limb.	Z.D.	Correction for South Limb.	Z.D.	Correction for North Limb.	Z.D.	Correction for North Limb.
30	—0".01	60	+0".10	30	—0".08	60	—0".10
35	0.00	65	+0".11	35	—0".09	65	—0".10
40	+0".02	70	+0".13	40	—0".09	70	—0".10
45	+0".03	75	+0".15	45	—0".10	75	—0".10
50	+0".05	80	+0".16	50	—0".10	80	—0".09
55	+0".07			55	—0".10		

The correction tabulated above is the sum of two corrections, of which the first is due to the difference in parallax between the limb and centre of the Moon; and the second is due to the employment of the arc of equatorial parallax instead of its sine, for which the formula is,

$$\frac{(\text{Seconds of hor. parallax})^3}{6} \times \sin \text{Z.D.} \times \cos^2 \text{Z.D.} \times \sin^2 1''.$$

For Mösting A the log sine horizontal equatorial parallax of the crater is obtained by interpolation from the *Berliner Jahrbuch*.

For the computation of the parallaxes of the small planets, the distances are taken from the Supplement to the *Nautical Almanac*, or from the *Berliner Jahrbuch*, for planets given in those works.

SEMI- DIAMETER.—The semidiameters of the Sun, Moon, and planets are given in the lower part of the page. For the Moon, Mercury, and Uranus, they are taken

unchanged from the *Nautical Almanac*. For the Sun, Venus, Mars, Jupiter, and Saturn, the semidiameters are also taken unchanged from the *Nautical Almanac*, except in cases where only one limb has been observed. In these cases, for the Sun, the following corrections have been applied to the tabular semidiameter, the subscript numbers denoting the number of observations on which they depend :—

Mr. Eddington	— 0".30 ₉	Mr. Cullen	— 0".31 ₂₂
Mr. Witchell	— 0".24 ₃₀	Mr. Daniels	— 0".01 ₉
Mr. Storey	— 1".24 ₈	Mr. Shepperd	+ 0".81 ₇
Mr. Evans	— 0".23 ₃₀	Mr. Acton	+ 0".71 ₉

For Venus the correction + 0".392 + 0.027 × tabular semidiameter, derived from an investigation mentioned at page xxxix, has been applied to the semidiameter given in the *Nautical Almanac*; and for Mars, Jupiter, and Saturn, a correction usually derived from neighbouring observations, and specified in the notes, has been similarly applied.

For Mösting A the correction to reduce to the centre of the Moon has been obtained by interpolation from the *Berliner Jahrbuch*.

GEOCENTRIC N.P.D. OF CENTRE.—The numbers in this column are found by combining the apparent zenith-distance with the refraction, parallax, semidiameter, and assumed colatitude 38°.31'.21".80.

§ 5. *Runs of each Microscope-Micrometer of the Transit-Circle; and Zenith-Points of the Transit-Circle.*

With regard to the *Runs of the Microscopes*, the columns under A, B, C, D, E, F contain respectively the number of revolutions of each micrometer which measures an arc of 5' on the circle; and the last columns give the sums of these numbers for the six microscopes, and exhibit the number of revolutions which, for the mean of the six microscopes, measures an arc of 30', on the supposition that the screws of the micrometers have sensibly equal values. In the first calculations for the reduction of the observations it is assumed that the space of 30' corresponds approximately to $\frac{50}{51} \times 30^{\text{r}}000$, or to 29^r4118; and the fiftieth part of the micrometer-reading for the mean of the microscopes being added to the micrometer-reading, the remaining correction, depending on the difference between the values in the last columns of the tables in question and 29^r4118, is computed in the way that has been explained (page xliii).

With regard to the table of *Zenith-Points*, the following explanation will be sufficient.

Whenever observations of stars have been made by reflexion, the results for zenith-point derived from them have been combined with those derived from the observations of the reflected image of the wire, see page xlvii. The results are always divided into three groups, viz. : those resulting from north stars, those from south stars, and those from nadir-observations, and the mean of each group is taken. In combining these a weight 2 is attributed to each of the means of the two groups of star-observations, and a weight 1 to the mean of the group of wire-observations, in order to prevent too much weight being given to those divisions of the circle which are always under the microscopes for the wire-observations. From the pains taken to equalize the number of observations of stars north and south of the zenith, no undue weight is, on the average, given to either of these positions ; and, on the whole, it is presumed that, by this method of combination, the best possible average result is obtained. In the few cases where reflexion observations of only north or only south stars have been obtained, a correction is applied for the R—D discordance in obtaining the zenith-point. The value of this correction, which is given in a footnote in the tables of zenith-points, is derived from the adopted value $0''.600 \sin Z.D.$ of the R—D discordance.

From 1897 January 1 a correction of $-0''.25$ has been applied to observations of the nadir, in forming the zenith-point on those days when no reflexion observations of stars were obtained. (See p. xlvii.)

§ 6. *Ledgers of Mean Right Ascensions and Mean North Polar Distances
for 1910·0 of Stars observed in 1908.*

These ledgers are divided into two parts. Part I. contains Fundamental and Zodiacal stars ; Part II. contains the stars of the Oxford Astrographic Zone.

The apparent right ascensions and north polar distances (obtained as explained in §§ 1 and 3) are reduced to mean place for the epoch 1910·0, with Newcomb's constants of precession. Since 1902 a further correction for latitude variation has been applied to the north polar distances. This correction is taken from provisional results sent by Professor Albrecht.

For stars in the lists of the national ephemerides, the correction to reduce to 1910·0 is found by subtracting the apparent R.A. or N.P.D. given in those ephemerides from the mean R.A. and N.P.D. for 1910·0 in the same ephemerides.

For the remaining stars of Part I. and for all the stars of Part II., star constants are formed for 1910·0, and the star corrections are computed by multiplying these by the day numbers with the help of Crelle's or Cotsworth's multiplication tables, thus avoiding the use of logarithms.

The logarithms of the day numbers, with Newcomb's constants, are given in the *Nautical Almanac* for mean midnight of each day and it has been the practice to interpolate values of these quantities to within two hours of the time of observation. The additional correction depending on the longitude of the moon has only been taken into account for those stars for which apparent places are given in the ephemerides for every day of the year.

The places for 1910·0 are not corrected for secular variation during the interval between the date of observation and 1910·0 (except in the case of stars contained in the ephemerides, which have been reduced as explained above). It is intended ultimately to apply mean corrections for secular variation when the annual results are collected in the *Greenwich Catalogue* for 1910.

Corrections for the proper motion in the interval between the time of observation and the epoch 1910·0 are applied to the stars of Part I. but *not* to those of Part II. The proper motion adopted in the former case is given in the *Catalogue of Mean R.A. and N.P.D.* (§ 7).

The corrections for personal equation have been applied in this section when the observations of stars for clock-error have been made by one observer and those of any other star requiring reduction on the same day by another observer.

Half weight has been given to a result for right ascension when the transit was observed at two or three wires only, observations at a single wire being rejected. A reflexion and direct observation at the same transit are considered as separate observations with full weight.

In Part I. the rules for the nomenclature of stars are the same as those in the section *Transits observed*, detailed on page xxx. In Part II. the Zone and No. in the *Bonn Durchmusterung* are used.

§ 7. *Catalogue of Mean R.A. and N.P.D. for 1910·0 of Fundamental and Zodiacal Stars observed in 1908.*

The results of observations of stars included in Part I. only of the ledgers are given in this catalogue.

The right ascensions in the catalogue are the means of the separate determinations for each star, and are identical with the means printed in the ledgers.

The north polar distances in the catalogue are obtained from the results in the ledgers by applying corrections for the discordance of direct and reflexion results,

and in the case of reflexion observations for inclination of the verticals at the surface of the mercury and at the centre of the transit-circle.

A constant flexure term of the R—D discordance, namely $+0''.60 \sin Z.D.$, has been adopted. This is sensibly the same as the mean values used in the 1880, 1890 and 1900 Catalogues.

For the circumpolar stars the following weights are assigned to the observations below pole—

N.P.D. not exceeding 15°	weight 1
N.P.D. 15° to 36°	weight $\frac{2}{3}$
N.P.D. 36° to 41°	weight $\frac{1}{2}$

Beyond N.P.D. 41° the observations are not combined. Direct and reflexion observations are treated as having equal weights.

As already explained Pulkowa refractions and an adopted colatitude $38^\circ 31' 21''.80$ have been used in the reductions, Newcomb's constants of precession have been adopted, and the results have been corrected for latitude variation.

The proper motions given are generally taken from Prof. Newcomb's *Catalogue of Fundamental Stars for 1900.0*, or from the *Catalogue of Zodiacal Stars for the epoch 1900.0 and 1920.0*. For the stars given in the *Berliner Jahrbuch*, but not in the *Nautical Almanac* or *Connaissance de Temps*, the proper motions of Prof. Anwers are used.

§ 8. *Horizontal and Vertical Diameters, Right Ascensions, and North Polar Distances of the Sun, Moon, and Planets, deduced from the Observations, and compared with the Nautical Almanac and other Ephemerides: with the inferred Position of the Ecliptic, the Geocentric Errors of the Sun, Moon, and Planets, in Longitude, and Ecliptic Polar Distance, and the Equations between the Geocentric Errors of the Planets and the Heliocentric Errors of the Earth and Planets.*

HORIZONTAL AND VERTICAL DIAMETERS OF THE SUN, MOON, AND PLANETS.

The duration of the passage of the Sun's diameter is found by subtracting the clock-time of transit of the first limb from that of the second limb in the *Transits observed*, &c., without any further correction. The tabular duration is found by doubling the time of the passage of the semidiameter given in the *Nautical Almanac*. The excess of the latter above the former is set down as the apparent error of the *Nautical Almanac*. The mean of 130 measures of the value of this error made during

the year is $+0^s.010$. The values obtained by the several observers are given on p. xxxviii.

The Sun's vertical diameter is found by subtracting the zenith-distance of the north limb, corrected for refraction and parallax, from that of the south limb similarly corrected. The tabular diameter is found by doubling the semidiameter of the *Nautical Almanac*. The excess of the latter above the former is set down as the error of the *Nautical Almanac*, and the mean of 139 measures of the value of this error made during the year is $+0''.04$. The values obtained by the several observers are given on p. liii.

For the duration of the passage of the Moon's diameter, the correction for defect of illumination is investigated as on p. xxxiii. The mean of two determinations of the error of the Moon's tabular duration of passage made in this year is $+0^s.085$.

The Moon's vertical diameter is found in the same manner as that of the Sun (the correction for defective illumination having been already applied, see page xlix). The mean of 12 determinations of the error of the Moon's tabular vertical diameter made during the year is $+0''.59$.

For the planets, the duration of passage of diameter is found by subtracting the clock-time of transit of the first limb from that of the second limb in the section *Transits observed*, the correction for defect of illumination being applied where necessary. This correction is double the amount applicable to the mean of limbs which is stated in the notes to that section.

The vertical diameters of Mercury, Venus, Mars, and Jupiter are found by subtracting the N.P.D. of the north limb from that of the south limb, after the correction for defective illumination has been applied, as explained at page 1. The vertical diameters of the other planets are obtained in a similar way, excepting that there is no correction for defective illumination.

RIGHT ASCENSIONS AND NORTH POLAR DISTANCES OF THE SUN, MOON, AND PLANETS.

As the annular illumination of the transit-circle was in use to 1908 Apr. 29; the central illumination with simple condensing lens from Apr. 30 to Sept. 16, and with compound corrected lens from Sept. 17 to the end of the year; and as the Sun, Mercury, and Venus are observed without and the clock stars generally with artificial illumination, the following corrections have been applied in this section to the observed Right

GREENWICH OBSERVATIONS, 1908.

h

lviii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Ascensions of the Sun, Mercury, and Venus to reduce to a uniform system of clock error, viz., that given by the annular illumination :—

	Date.	Correction.
Jan.	1-Apr. 29	$^s.00$
Apr.	30-Sept. 16	$+ ^s.06$
Sept.	17-Dec. 31	$+ ^s.02$

The observations to determine these corrections are given in the *Transit-Circle Tables*.

The observed right ascensions of the Sun's centre are generally the means of those deduced from the observations of the two limbs; but when one limb only has been observed, the R.A. of the centre is deduced from that of the limb, by application of the duration of transit of semidiameter given in the *Nautical Almanac*, corrected by the quantities given on page xxxviii. The right ascensions are transcribed from those in the section of *Transits observed*, without alteration, except that, when necessary, the correction for personal equation and the reduction to the annular illumination are applied.

The following are the values of the mean error of tabular right ascension of the Sun's centre (mean of two limbs) found by the different observers :—

	SE	W	JS	E	RC	SD	S	HA	Other Observers.
1908	$- ^s.11_{10}$	$- ^s.02_{24}$	$- ^s.07_7$	$- ^s.05_{19}$	$- ^s.10_{19}$	$- ^s.16_6$	$- ^s.08_5$	$- ^s.02_8$	$- ^s.07_{16}$

the subscript numbers denoting in each case the number of observations.

The observed north polar distances of the Sun are taken from the section of *Zenith-Distances*, &c., the corrections for discordance of direct and reflexion results and (since 1902) for latitude variation being applied. As has been previously explained, Pulkowa refractions and assumed colatitude $38^{\circ}.31'.21''.80$ have been used throughout in the reductions. The tabular right ascensions and north polar distances are taken without correction from the *Nautical Almanac*.

The following are the values of the mean error of tabular N.P.D. of the Sun's centre (mean of two limbs) found by the different observers :—

	SE	W	JS	E	RC	SD	S	HA	Other Observers.
1908	$+ ''03_9$	$+ ''12_{30}$	$- ''08_8$	$+ ''09_{20}$	$- ''39_{22}$	$+ ''21_9$	$- ''45_7$	$- ''78_9$	$- ''03_{25}$

As the microscope-micrometers for the observations of the north polar distance of the Sun are not read by the observer but by his assistant, whereas the determination of the zenith-point depends on the readings of the observer himself, any difference, depending on personality, in the readings of the observers and of the microscope-readers for the Sun would introduce a systematic error into the north polar distances of the Sun. With a view to determining the magnitude of this error, if it exists,

the microscopes for the midday nadir observation were read each day by the observer and microscope-reader. The results of this investigation are given in the *Transit-Circle Tables*.

The observed right ascensions of the Moon's centre are taken from the *Transits observed*, corrected, when necessary, for personal equation, and also for the Moon's motion in R.A. during the passage from the mean of wires to the true meridian. When both limbs of the Moon are observed, the correction for defective illumination (page xxxiii) is first applied to the proper limb, and the mean of the two is then taken without any further alteration.

The tabular right ascension of the Moon's centre is found by applying to the right ascension of the limb (given in the section, *Moon-culminating Stars*, of the *Nautical Almanac*) the sidereal duration of passage of semidiameter, which is given in the same section of the *Nautical Almanac*.

The following are the values of the mean error of tabular right ascension of the Moon's centre, from observations of the first and second limbs, found by the different observers :—

Moon 1 L.					
	W s.	JS s.	E s.	RC s.	Other Observers. s.
1908	-0°51 ₁₅	-0°49 ₄	-0°37 ₁₈	-0°55 ₁₇	-0°56 ₁₄
Moon 2 L.					
1908	-0°28 ₇	-0°39 ₄	-0°34 ₁₀	-0°31 ₁₅	-0°16 ₈

The observed north polar distances of the Moon's centre are taken from the section of *Zenith-Distances observed*, &c., corrected for discordance of direct and reflexion results and for the motion in N.P.D. during the passage from the mean of wires to the true meridian, and since 1902 for latitude variation.

The tabular north polar distances of the Moon's centre are taken from the section *Moon-culminating Stars* of the *Nautical Almanac*.

The following are the values of the mean error of tabular north polar distance of the Moon's centre, from observations of the north and south limbs, found by the different observers :—

Moon N. L.					
	W	JS	E	RC	Other Observers.
1908	-1°44 ₁₂	-1°00 ₈	-0°02 ₁₁	-1°84 ₁₆	-0°14 ₁₀
Moon S. L.					
1908	+1°36 ₁₁	+2°11 ₆	+1°70 ₁₈	+0°27 ₁₈	+1°54 ₁₁

From the beginning of 1905 the lunar crater Mösting A has been observed, as well as the Moon's limbs whenever practicable, the observations of transits serving to

connect the observations of the first and second limbs made before and after Full Moon respectively, while the observations of zenith distance in combination with similar observations made at the Cape Observatory serve to determine the parallax of the Moon.

For all the planets, the right ascensions are extracted from the twenty-first column of *Transits observed*, with no alteration, except for personal equation and reduction to annular method of illumination when necessary; the north polar distances are taken from the last column in the upper part of the page of the *Zenith-Distances observed*, &c., corrected for discordance of direct and reflexion results, and since 1902 for latitude variation. The tabular places are taken from the *Nautical Almanac*, for such planets as are given in that work and its appendices. For others the tabular places are taken from the *Berliner Jahrbuch*, or its Circulars; or occasionally from MS. ephemerides mentioned in the notes.

When only one limb of a planet is observed, the number I. or II., or the letter N, S, or C, in Columns 4 and 8, indicates that the first, second, north, or south limb, or the estimated centre, was observed.

ECLIPTIC INVESTIGATION.

The investigation of the position of the ecliptic has been made in the following manner. The mean of all the daily errors in each month of the R.A. and N.P.D. of the sun is taken to be the error for the day which is nearest to the mean of all the days of observation. When the same day is not found for R.A. and for N.P.D., an alteration of a unit or more has sometimes been made. From these, the error in the ecliptic polar distance is obtained by means of the factors R and S in the tables forming the second part of the *Appendix* to the *Greenwich Observations*, 1836. Supposing these errors to arise from an erroneous position of the ecliptic assumed in the *Nautical Almanac*, they may be expressed by the formula $x \cos l + y \sin l + z$, where l is the Sun's longitude. Weights are then assigned proportional to the number of observations of N.P.D., and the equations solved by the ordinary method of least squares. This method has been used since 1888, instead of the method of weighting and grouping the observations previously adopted.

MEAN ERRORS OF THE TABULAR PLACES OF THE SUN AND PLANETS.

For the *Mean Errors of the Tabular Geocentric Places of the Sun and Planets*, the observations have been collected into groups rarely exceeding a month in duration, and the mean of all the errors in each group, for R.A. and for N.P.D., giving half weight to observations of a single limb of the Sun in either element, is supposed to be the error for the day which is nearest to the mean of all the days of observation

TRANSIT CIRCLE.—ERRORS OF TABULAR PLACES OF PLANETS. lxi

in the group, an alteration of a unit or more being sometimes made, in order to refer R.A. and N.P.D. to the same day. The errors in longitude and E.P.D. are formed by the use of the numbers P, Q, R, S in the *Appendix* to the *Greenwich Observations*, 1836. For any of the four small planets, Ceres, Pallas, Juno, and Vesta, whose latitude exceeds the limits of the tables, the longitude and E.P.D. are computed, 1st, from the R.A. and N.P.D. of the *Nautical Almanac*; 2nd, from these quantities affected with the errors in R.A. and N.P.D.; and the difference between these results gives the errors in longitude and E.P.D.

For the *Errors in the Tabular Heliocentric Places of the Planets*, the following formulæ are used:—

For the small planets, Ceres, Pallas, Juno, and Vesta.

Let R = radius vector of planet, L = planet's heliocentric longitude,
 Δ = planet's distance from Earth, λ = planet's geocentric longitude,
 r = Earth's radius vector, l = Earth's heliocentric longitude.

Then,

$$\begin{aligned} \text{Error of geoc. long.} &= \frac{R \times \cos \text{hel. lat.} \times \cos (\lambda - L)}{\Delta \times \cos \text{geoc. lat.}} \times \text{error of hel. long.} \\ &- \frac{\sin (\lambda - L)}{\Delta \times \cos \text{geoc. lat.} \times \sin 1''} \times \text{error of projection of planet's radius vector.} \\ \text{Error of hel. E.P.D.} &= \frac{\Delta \times \cos \text{hel. lat.}}{R \times \cos \text{geoc. lat.}} \times \text{error of geocent. E.P.D.} \\ &- \sin \text{hel. lat.} \times \cos \text{hel. lat.} \times \tan (\lambda - L) \times \text{error of geocent. long.} \\ &- \frac{r \times \tan \text{geoc. lat.} \times \cos (L - l)}{R^2 \cos (\lambda - L) \times \sin 1''} \times \text{error of projection of planet's radius vector.} \end{aligned}$$

For the other planets, the following are used:—

$$\begin{aligned} \text{Error of geocent. long.} &= \frac{R \times \cos (\lambda - L)}{\Delta} \times \text{error of planet's heliocentric longitude,} \\ &- \frac{\sin (\lambda - L)}{\Delta \times \sin 1''} \times \text{error of projection of planet's radius vector,} \\ &- \frac{r \times \cos (\lambda - l)}{\Delta} \times \text{error of Earth's heliocentric longitude,} \\ &+ \frac{\sin (\lambda - l)}{\Delta \times \sin 1''} \times \text{error of Earth's radius vector.} \\ \text{Error of hel. E.P.D.} &= \frac{\Delta}{R} \times \text{error of geocent. E.P.D.} \end{aligned}$$

The errors of the Earth's place are retained in the formulæ for the larger planets, as being probably comparable in magnitude with the errors of the places of the planets: but the errors of the tabular places of the small planets being in general large, in their case the Earth may be supposed to move exactly in the orbit assigned by the tables.

lxii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

It is to be remarked that, in the column *Extent of Group*, the day given is, as in past years, the civil day (commencing at midnight) on which the observation was made, and that this occasionally differs by a unit from the astronomical day under which the observations may be found in the sections of *Transits observed* and *Observations of Zenith-Distance*: also in the column *Mean Day* the day given is the civil day corresponding to the time of observation, or to the mean of times of observation.

COMPARISON OF THE TABULAR ERRORS OF R.A. AND N.P.D. OF THE CENTRE
OF THE MOON IN THE YEARS 1905-6-7-8.

In this table are brought together the tabular errors of the centre of the moon, derived from observations of Mösting A and the limb, both with the transit-circle and the altazimuth (see below), and a mean error is deduced. The table starts with the year 1905, that being the first year in which observations of Mösting A were taken.

Corrections representing the Moon's motion in R.A. and N.P.D. in the interval between the observed and tabular times of transit have been applied in forming these Errors of the Moon's tabular place. The Apparent Errors of R.A. and N.P.D. in the section, "Right Ascensions and North Polar Distances of the Centre of the Moon," are formed, as in past years, by simple subtraction of the "R.A. or N.P.D. from Observation" from the "R.A. or N.P.D. in *Nautical Almanac*," without application of the above-mentioned corrections.

MEAN ERRORS OF THE TABULAR PLACES OF THE MOON (1908).

Commencing with year 1908, the mean results from the limb and Mösting A, obtained with the transit-circle and altazimuth (given in the previous table) have been used in this discussion.

The errors of tabular longitude and ecliptic polar distance of the Moon are deduced from the errors of tabular R.A. and N.P.D. by the use of the numbers, P, Q, R, S, for all the meridian observations.

§ 9. *Meridian observations with the Altazimuth.*

The method of reduction of the transits and meridian zenith distances is similar to that for the transit-circle and no further explanation is required for the section of Transits and Meridian Zenith Distances beyond the short notes at

the beginning of these sections. As in the case of the Transit-Circle, the lunar crater Mösting A has been observed with the Altazimuth whenever practicable, and a comparison of the results with those from the moon's limbs is given on pp. {74} and {75}.

In the *Altazimuth Tables* at the end of this *Introduction* are given :—

The values of 1^{rev.} of the micrometer-screws.

The adopted wire intervals.

The adopted division errors.

Further details of the methods by which these were obtained are given in the *Tables* at the end of the *Introduction* for 1901.

A discussion of the R—D discordance and of the colatitude is also given in substantially the same form as for the transit-circle.

A comparison of the North Polar Distances observed in the four positions of the instrument is also given in the *Altazimuth Tables* at the end of this *Introduction*.

§ 10. *Extra-meridian observations with the Altazimuth.*

The method of observation has been already described on page xiv, the essential point being that the means of the times of observation in azimuth and zenith distance should be as nearly as possible the same, so that the azimuth and zenith distance can be readily referred to the same instant of time.

The following explanation refers to the observations and reductions as printed in this section.

The first twelve columns of this section require no explanation. The Apparent Error of Tabular Z.D. in the thirteenth column is the excess above the concluded Zenith Distance of the Tabular Zenith Distance, which is taken from tables constructed for each azimuth with argument N.P.D. according to the formula

$$\begin{aligned} \text{where} \quad \cos (Z + \theta) &= \sec \gamma_0 \cos \Delta \\ \sin \gamma_0 &= \sin \gamma \sin A \\ \tan \theta &= \tan \gamma \cos A \end{aligned}$$

Z being the tabular zenith distance, or the zenith distance on crossing the standard azimuth corresponding to the tabular N.P.D. which is denoted by Δ . The colatitude is denoted by γ and the azimuth reckoned from the south by A .

The tabular places are taken, whenever possible, from the Greenwich Clock Star List

or recent Greenwich Catalogues, or from Newcomb's Fundamental Catalogue, in this order of preference.

The fourteenth column is derived from the thirteenth by converting arc into time at the rate $15'' \sin \gamma \sin A = 15 m$ to a second of time. This is the rate of motion in zenith distance and is constant throughout the same azimuth.

The second column of the right hand page is the mean of the separate taps over the horizontal wires registered on the chronograph reduced to the centre wire by a process analogous to the completion of transits in the meridian. In this case, however, the reduction to the centre wire is simpler, because the transits are usually symmetrical, and also because the motion in zenith distance is constant throughout the azimuth. The observed time of transit over the centre wire includes the correction for curvature of the star's path, when sensible.

The fourth column gives the tabular time T of transit over the standard azimuth, computed with the help of tables from the formulæ :—

$$\begin{aligned} \text{where} \quad \cos (T - \alpha + \phi) &= \tan \gamma_0 \cot \Delta \\ \tan \phi &= \sec \gamma \cot A \end{aligned}$$

α being the tabular right ascension.

The clock error is the excess of this latter quantity over the quantity in the second column corrected by the application of the quantity in the last column of the left hand page. The clock rate is taken from the transit-circle observations, and the clock error reduced to 0^h sidereal with this rate is exhibited in the fifth column. Means are taken in this column for each group observed by the same observer on the same side of the zenith.

The sixth column gives the adopted clock error, which is the mean of the two groups observed, east and west of the zenith.

The observed time of transit over the centre vertical wire given in the seventh column is corrected for curvature of the star's path, when sensible. The errors of collimation and level given in the ninth column are converted into time by the divisors $15 m \cot S$ and $15 m \cot S \sec Z$ respectively, S being the parallactic angle. The seconds corrected are given in the tenth column. The result in the tenth column is corrected for the clock error and rate given in the sixth column, and the excess of the tabular transit T over the quantity thus derived is set down in the eleventh column, under the heading Apparent Error of Tabular Time of Transit. From this apparent error is deduced the Apparent Error of Azimuth, the quantity set down in the twelfth column being the determination of this error from the particular star, derived by multiplying the quantity in eleventh column by $15 m \cot S \operatorname{cosec} Z$.

In the thirteenth column is given the adopted error of azimuth, the weighted mean of the quantities in the preceding column, the weight assigned being the nearest integer to $10 \sin^2 Z$. The Apparent Errors of Tabular Place in Azimuth and Zenith Distance given in the fourteenth and fifteenth columns are derived by multiplying the excess of the azimuth error and clock error as determined from the particular star, over their adopted values by $\sin Z$ and $15 m$ respectively.

The remaining columns require no explanation.

At the foot of the page are given details of the reductions for the Sun, Moon and Planets.

The second column of the left hand page gives the quantity in the seventh column of the right hand page above corrected for clock error, rate, collimation, level, and azimuth. The clock error used for Azimuth observations is derived from stars on both sides of the Zenith; that for Zenith Distance observations from stars on the same side of the Zenith as the object observed. Correction is still required on account of semidiameter, parallax, and orbital motion. This correction is calculated as a fraction $\frac{A}{B}$ whose numerator and denominator are approximately equal to the semidiameter and $15 m \cot S$, a quantity called the collimation divisor. The numerator of this fraction, A , requires to be corrected for parallax and curvature. The former correction is the same fraction of the semidiameter that the perpendicular from the centre of the Earth upon the standard azimuth is of the linear semidiameter of the object observed. The correction for curvature is $\frac{1}{2} \times \frac{(15 t)^2}{206265} \sin \Delta \cos \Delta$ in seconds of arc applicable to N.P.D. Its horizontal component is found by multiplication by $\sin S$. It will be observed that $\sin \Delta \sin S$ is a constant for the azimuth and that $\cos \Delta$ varies between small limits and is never far different from unity. A sufficient approximation to the value of t in the numerator is obtained by dividing the semidiameter by the collimation divisor.

For the denominator B , of the fraction, the collimation divisor requires correction for the contribution of the object's orbital motion to the apparent motion across the azimuth of the instrument.

The contribution arising from the orbital motion in R.A. is the same fraction of the collimation divisor that the orbital motion in R.A. is of the diurnal motion, $15 \sin \Delta$. In the case of the Moon, however, the mean orbital motion is considered as coalescing with the diurnal motion, and the inequalities of its motion are alone attributed to orbital motion. As a consequence, the fraction whose value is exhibited in the ninth column of the printed results is expressed in lunar seconds in the case of

GREENWICH OBSERVATIONS, 1908.

i

the moon and sidereal seconds for the sun and planets. A lunar second denotes the $\frac{1}{86,400}$ part of a lunar day, which is the mean interval between consecutive transits. The contribution to the collimation divisor of a motion in N.P.D. of one second of arc in a second of time is clearly $\sin S$, and for a greater or less motion in N.P.D., the effect is in proportion.

This suffices for the explanation of the third to the ninth columns inclusive. The tenth column is the result of the application of the ninth column (which must be reduced to sidereal seconds when necessary) to the second column.

The eleventh column is taken from the second column of the right hand page above corrected for clock-error and rate. The clock-error employed is derived from the stars on the same side of the zenith only, and is that set down in brackets in the fifth column of the right hand page above.

The difference $T - t_2$, the excess of the tenth column over the eleventh is always small: the twelfth column exhibits this difference converted into arc by the factor $15m$ corrected for orbital motion in R.A. and N.P.D. The method of correction is analogous to the correction of the collimation divisor.

The thirteenth column exhibits the semidiameter, interpolated from the *Nautical Almanac*.

The fourteenth column exhibits the horizontal parallax, reduced to the geocentric distance of the instrument, and multiplied by $\sin(z - \zeta)$ where ζ is the angle of the vertical projected upon the plane of the instrument ($11' 12'' \cos A$), the ellipticity of the Earth being assumed to be $1/300$ and the height of the instrument above sea level being taken as 170 feet.

The fifteenth column exhibits the excess of the true correction for semidiameter and parallax.

$$\sin^{-1} [(\rho \sin(z - \zeta) \pm \mu) \sin p]$$

over the sum of the preceding corrections. In this formula, ρ denotes the geocentric distance of the instrument, μ the linear diameter of the planet, and p its equatorial horizontal parallax.

On the right hand page the second column gives the application of the corrections in the four last columns to the zenith distance in the twelfth column of the left hand page above.

The third and fourth columns contain the N.P.D. and hour angle corresponding to Z , the latter combined with T gives the R.A. in the fifth column: the remaining columns require no further explanation.

The method of reducing lunar observations is described by Mr. Cowell in the *Monthly Notices*, vol. lxii. p. 503, where full formulæ and an example are given.

§ 11. *Ledgers of Mean Right Ascensions and North Polar Distances and Catalogue.*

The star ledgers and the annual star catalogue are printed together. The position of the instrument is indicated in each case, and the Mean N.P.D. is given as obtained from each circle separately. The separate results are corrected for latitude variation, and the means of groups in each position of the instrument are corrected for the R—D discordance.

§ 12. *Results of Observations of Planets.*

The planetary results are similar to those obtained with the transit-circle and require no further explanation.

Attention may be called to the fact that the results of the observations of the Moon with the altazimuth are compared and combined with those obtained with the transit-circle in special tables given at the end of the Planetary Results of the transit-circle observations (see § 8).

§ 13. *Meridian Zenith Distances of Stars observed with the Reflex Zenith Tube.*

It was pointed out by Mr. Chandler in the *Astronomical Journal*, No. 511 (1901 November) that the anomalous results found from observations of γ Draconis with this instrument were to be explained by the variation of latitude and that the instrument was well adapted for the determination of the amount of this variation.

It was decided, therefore, early in 1902, to resume without delay the observations of γ Draconis, which had been intermitted since 1899 May (when all the wires were found broken) and to observe such other stars as passed sufficiently near the zenith and were sufficiently bright. By suitable modification of the illumination it has been found possible to observe stars down to the magnitude 7.0, and, by mounting the eyepiece in a slide, good definition at a distance of 50' or more from the zenith (centre of the field) has been secured.

A working catalogue of suitable stars was prepared and systematic observations began in 1902 June. The daily results are given in substantially the same form as the observations of γ Draconis in volumes up to that for 1881, to which reference may here be made.

An account of the determinations of the mean value of 1 revolution of the screws, of

lxviii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

the errors of the screws, and of the intervals of the wires is given in the *Reflex Zenith Tube Tables* for 1903.

§ 14. *Eclipses, Occultations, and Transits of Jupiter's Satellites, compared with the Nautical Almanac; and Occultations of Stars by the Moon with the Equations deduced from the Occultations.*

The clocks used in these observations are in general compared with the Sidereal Standard or with the clock Hardy near the time of the observation, and the mean solar time is computed in the usual way.

For the computation of the Occultations of stars by the Moon, the star's place is taken from the "Elements for facilitating the Computation of Occultations" in the *Nautical Almanac* when given there. The Moon's geocentric place and horizontal equatorial parallax are interpolated with second differences from the *Nautical Almanac*, and are used without alteration. From the beginning of 1900 a correction has been applied to the *Nautical Almanac* semidiameter, to reduce it from Hansen's bright limb value $15' 34''.09$ to Struve's dark limb value $15' 32''.65$. The correction to be applied to the parallax of the Moon's centre, in order to obtain that for the point of the limb at which the occultation takes place, is derived from a table given in the *Introduction to the Greenwich Observations* of 1899, the principles of the formation of which are given in the volumes of the *Greenwich Observations* from 1843 to 1849.

Let now δ and θ be the N.P.D. and hour-angle of the star, which are the same as the apparent N.P.D. and hour-angle of the point of the Moon's limb; δ' and θ' those of the corresponding point of the limb as seen from the Earth's centre; l the geocentric latitude of the place; and P' the corrected horizontal parallax; then, by an investigation similar to that given in the *Introductions* to the volumes, from 1843 to 1849, it may be shown that

$$\sin (\theta - \theta') = \frac{\text{Radius of parallel for Greenwich}}{\text{Moon's distance from the Earth's axis}} \times \sin \theta.$$

$$\sin (\delta - \delta') = \frac{\sin l \cdot \sin \delta \cdot \sin \theta \cdot \sin P'}{\sin \frac{1}{2} (\theta + \theta') \cdot \cos \frac{1}{2} (\theta - \theta')} - \cot \frac{1}{2} (\theta + \theta') \cdot \tan \frac{1}{2} (\theta - \theta') \cdot \sin (\delta + \delta').$$

For the first step of the computation the following quantities are formed:—

$$F = \log. \sin \text{ star's hour-angle} + \log. \text{ seconds of corrected eq. hor. parallax} + 9.4942268.$$

$$G = \log. \sin \text{ star's N.P.D.} + \log. \sin \text{ star's hour-angle} + \log. \text{ seconds of corrected eq. hor. parallax} + 9.8913966.$$

(The former numerical constant is the logarithm of half the distance of Greenwich from the Earth's axis, and the latter is the logarithm of the distance of Greenwich from the plane of the equator; supposing the Earth's ellipticity = $\frac{1}{300}$).

The preceding formulæ are then adapted to logarithmic computation, and the equations are solved by successive trials, assuming a value for δ' , in the following manner:—

$$\begin{aligned} \log. \frac{1}{2} (\theta - \theta') \text{ in seconds} &= F - \log. \sin \delta' + \log. \frac{\text{sine}}{\text{arc}} \text{ for hor. parallax} \\ &+ \log. \frac{\text{arc}}{\text{sine}} \text{ for } (\theta - \theta'). \end{aligned}$$

$$\begin{aligned} \log. 1^{\text{st}} \text{ number} &= G + \log. \secant \frac{1}{2} (\theta - \theta') + \log. \frac{\text{sine}}{\text{arc}} \text{ for hor. parallax} \\ &+ \log. \frac{\text{arc}}{\text{sine}} \text{ for } (\delta - \delta') - \log. \text{sine } \frac{1}{2} (\theta + \theta'). \end{aligned}$$

$$\begin{aligned} \log. 2^{\text{nd}} \text{ number} &= \log. \frac{1}{2} (\theta - \theta') \text{ in seconds} + \log. \frac{\tan}{\text{arc}} \text{ for } \frac{1}{2} (\theta - \theta') \\ &+ \log. \sin (\delta + \delta' - 180^\circ) + \log. \cot \frac{1}{2} (\theta + \theta') + \log. \frac{\text{arc}}{\text{sine}} \text{ for } (\delta - \delta'). \end{aligned}$$

$$\delta - \delta' = 1^{\text{st}} \text{ number} + 2^{\text{nd}} \text{ number}.$$

In the first trials, the $\log. \frac{\text{arc}}{\text{sine}}$, &c. are omitted. The convergence of these approximations is extremely rapid.

When δ' and θ' are found accurately, the distance of the corresponding point from the Moon's centre is thus found, by means of a subsidiary angle ψ :—

$$\begin{aligned} \text{If } \log. \tan \psi &= \log. \text{diff. R.A. of Moon's centre and corresponding point} + \frac{1}{2} \log. \\ &\sin \text{N.P.D. of Moon's centre} + \frac{1}{2} \log. \sin \delta' - \log. \text{diff. N.P.D. of} \\ &\text{Moon's centre and corresponding point;} \end{aligned}$$

$$\text{Then } \log. \text{dist.} = \log. \text{diff. N.P.D.} - \log. \cos \psi,$$

$$\text{or} = \log. \text{diff. R.A.} + \frac{1}{2} \log. \sin \text{N.P.D. of centre} + \frac{1}{2} \log. \sin \delta' - \log. \sin \psi.$$

The coefficients of small variations of the north polar distance and of the difference of right ascension (in the expression for distance) are computed by the formulæ:—

$$\log. 1^{\text{st}} \text{ number} = 2 \log. \text{diff. R.A.} + \log. \text{sine (sum of N.P.D.)} - \log. \text{dist.} + 4.0835.$$

$$\log. 2^{\text{nd}} \text{ number} = \log. \text{diff. N.P.D.} - \log. \text{distance}.$$

$$\text{Coefficient of variation of greater N.P.D.} = 1^{\text{st}} \text{ number} + 2^{\text{nd}} \text{ number}.$$

$$\text{Coefficient of variation of smaller N.P.D.} = 1^{\text{st}} \text{ number} - 2^{\text{nd}} \text{ number}.$$

$$\text{Log. coefficient of variation of diff. R.A.} =$$

$$\log. \text{diff. R.A.} + \log. \sin \delta' + \log. \sin \text{Moon's N.P.D.} - \log. \text{distance}.$$

The variation of the R.A. of the corresponding point contains the following terms:—

- 1st. The alteration of the R.A. of the star by the quantity e'' will alter the R.A. of the corresponding point by very nearly the same quantity e'' .
- 2nd. The alteration of the horizontal equatorial parallax in the proportion of $1 : 1 + \frac{m}{1000}$ will alter all the deduced parallaxes (in R.A. and in N.P.D.) in

lxx INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

nearly the same proportion: and therefore the R.A. of the corresponding point will be altered by $\frac{m}{1000} \times$ correction for parallax in R.A.

- 3rd. The alteration in the position of the Moon, with regard to the meridian depending on the alteration of time t^s , will introduce an alteration in the correction for parallax. It is computed by the following formula:—

Alteration in correction of R.A. of corresponding point for parallax, depending on the alteration of time =

$$15'' \times t \times \left\{ \begin{array}{l} \sin P \cdot \cos l \cdot \operatorname{cosec} \text{N.P.D.} \cdot \cos \text{hour-angle} \\ + \sin^2 P \cdot \cos^2 l \cdot \operatorname{cosec}^2 \text{N.P.D.} \cdot \cos 2 \text{hour-angle} \end{array} \right\}.$$

The variation of the R.A. of the Moon's centre is $x'' + t \times$ change of R.A. in 1^s .

The variation of the N.P.D. of the corresponding point contains three terms analogous to those for R.A.: namely—

- 1st. The alteration of the star's N.P.D. by the quantity f'' will alter the N.P.D. of the corresponding point by f'' nearly.
- 2nd. The correction for parallax in N.P.D. will be altered by $\frac{m}{1000} \times$ correction for parallax in N.P.D.
- 3rd. The increase of hour-angle (considered positive when the Moon is west of the meridian), depending on the alteration of time t^s , will alter the correction for parallax in N.P.D. by the following quantity:—

$$15'' \times t \times \left\{ \begin{array}{l} (-\sin P \cdot \cos l \cdot \cos \text{N.P.D.} - \sin^2 P \cdot \sin l \cdot \cos l \cdot \cos 2 \text{N.P.D.}) \times \sin \text{hour-angle.} \\ + (-\frac{3}{2} \sin^2 P \cdot \cos^2 l \cdot \cot \text{N.P.D.} + \sin^2 P \cdot \cos^2 l \cdot \cot \text{N.P.D.} \cdot \cos^2 \text{N.P.D.}) \\ \times \sin 2 \text{hour-angle.} \end{array} \right\}$$

The variation of the N.P.D. of the Moon's centre is $y'' + t \times$ change of N.P.D. in 1^s .

The computed distance, with addition of the sum of the products of the preceding variations by their proper coefficients, is made equal to the semidiameter increased by the term, semidiameter $\times \frac{n}{1000}$; and thus the final equation is formed.

For a planet, the R.A. and N.P.D. of the centre are used to find those of the corresponding point of the limb; the planetary parallax is subtracted from that of the Moon; and the distance of the corresponding point is equated to the sum (or difference) of the semidiameters.

§ 15. *Micrometric Observations of Double Stars.*

These observations were made with a position-micrometer on the 28-inch refractor. The method of observation consists in turning the micrometer till its fixed wire is parallel to the line joining the stars (so that they can be simultaneously bisected by it, by moving the telescope with one of the slow-motion rods) and reading the position-circle. The two movable wires, which are perpendicular to the fixed wire, are then moved till they bisect the two stars, and the readings of the micrometer-heads are taken. This may be done in two ways, the wire moved by the right hand screw bisecting the star on the right or left, as the observer wishes, and the wire moved by the left hand screw bisecting the other star in each case. The zero of position-angle is obtained by placing the fixed wire so that the diurnal motion carries a star along it. As the values of 1^{rev.} of the micrometer-screws are equal, and their readings increase in opposite directions, the distance between the stars is obtained by taking the sum of the readings of the micrometer-heads and subtracting the sum of the corresponding readings when the wires coincide; or if the measure has been made with wires reversed, by subtracting the sum of the readings from the reading for coincidence. When the observation has been made both *directly* and with wires reversed, the distance is half the difference of the sum of the micrometer-readings in the two cases.

The value of one revolution of each micrometer-screw is 12".06. It was determined in 1894 November from transits of stars near the pole. The observations are given in the volume for 1894.

The nomenclature of the stars in column 1 refers to the original observer's catalogue. Column 2 gives the name adopted for the star in the Greenwich Catalogues. Columns 3 and 4 contain the approximate R.A. and N.P.D. for 1900. Columns 5 and 6 give the measured position-angle and distance on different nights, and the means. Column 7 gives the number of measures of each element on the separate nights. Column 8 gives the magnitudes derived from the original observers' catalogues. Column 9 gives the epoch of the observations, expressed as the fraction of the year. In column 10 the magnifying power used is given, and in column 11 such remarks as appear necessary.

§ 16. *Micrometric Measures of the Equatorial and Polar Diameters of Jupiter and Satellites in the years 1894, 1895, 1896, 1906, 1907, 1908.*

Full explanations are given in the section itself.

§ 17. *Observations of Comets and Minor Planets from Photographs taken with the 30-inch Reflector of the Thompson Equatorial.*

The Thompson Equatorial, carrying the 26-inch photographic refractor at one end of the declination axis, and the 30-inch reflector at the other, is briefly described on page xix. The focal length of the reflector is approximately 11 feet 5 inches, giving a scale of 1' to the millimetre nearly.

The photographs are taken on plates 16 centimetres square, on which a *réseau* of cross lines 5 millimetres apart is printed in the same manner as for the Photographic Map of the Heavens.

The measurement of the photographs was made with the Astrographic Micrometer, which has a glass diaphragm with cross scales at the focus of the microscope. These scales are divided into 100 parts, and the micrometer is readily adjusted so that 100 divisions of the scale are equal, very nearly, to the distance between two *réseau* lines as viewed with the microscope. The intersection of the scales is made to bisect the object, and the positions of the *réseau* lines on each side are read off on the scales by estimation to the thousandth part of a *réseau* interval. The rectangular co-ordinates of the images are thus obtained in units of 1 *réseau* interval, which for both the 30-inch reflector and the 13-inch refractor is approximately 5'.

The plates are in all cases measured in reversed positions to eliminate personality. Where several images of the same object are obtained on a plate, they are all measured and the mean taken. The magnifying power used is 15.

The determination of Right Ascension and Declination is made by Professor Turner's method (*Monthly Notices, R.A.S.*, vol. liv., p. 11), as follows:—

With an assumed Right Ascension and Declination of the centre of the plate, the "Standard Co-ordinates," ξ and η , of those stars which are used as reference stars are computed by the formula—

$$\xi = \tan (\alpha - A) \cos \phi \sec (\phi - D),$$

$$\eta = \tan (\phi - D)$$

$$\text{where } \tan \phi = \tan \delta \sec (\alpha - A)$$

In this formulæ—

α is the R.A. of the star,

A „ „ centre of the plate

δ is the Dec. of the star

D „ „ centre of the plate

and ϕ is an auxiliary angle.

OBSERVATIONS OF COMETS, MINOR PLANETS, AND SATELLITES OF JUPITER AND SATURN. lxxiii

The "Standard Co-ordinates" of the reference stars are compared with the measured co-ordinates x, y ; and from equations of the form—

$$\begin{aligned}\xi - x &= ax + by + c \\ \eta - y &= dx + ey + f\end{aligned}$$

the constants of each plate a, b, c, d, e, f are deduced. By means of these constants, the "Standard Co-ordinates" of any object on the plate are obtained by applying corrections $ax' + by' + c, dx' + ey' + f$ to the measured co-ordinates x', y' , from which the Right Ascension and Declination are deduced by inversion of the trigonometrical formulæ given above.

It should be noticed that the Right Ascension and Declination are referred to the equator and equinox of 1908·0, and that the position of a comet or minor planet obtained in this way is referred to this epoch and is corrected for the part of the aberration arising from the Earth's motion.

In each case the uncorrected measures of the co-ordinates of the reference stars and the comet are given, and the deduced Right Ascension and Declination for the equator and equinox of 1908·0. For the reference stars the comparison with the assumed Right Ascension and Declination is also given.

The adopted plate constants, and the Apparent Right Ascension and Declination reduced to the equator and equinox of 1908·0 are given in separate tables.

§ 18. *Observations of the Sixth, Seventh and Eighth Satellites of Jupiter from Photographs taken with the 30-inch Reflector, during the opposition 1907–8.*

The photographs were taken and measured as described in the preceding paragraph.

The method of reduction employed, in order to eliminate the optical distortion of the reflector, and the determination of the error of the Tabular Place of the planet are described in the Note at the beginning of the section itself.

§ 19. *Observations of the Ninth Satellite of Saturn (Phæbe) from Photographs taken with the 30-inch Reflector during the opposition 1908.*

The photographs were taken and reduced as in the case of Jupiter's Sixth, Seventh and Eighth satellites.

GREENWICH OBSERVATIONS, 1908.

k

§ 20. *Observations of the Satellite of Neptune from Photographs taken with the 26-inch Refractor of the Thompson Equatorial during the opposition 1907-8.*

In taking the photographs from which these observations were made, an occulting shutter placed immediately in front of the plate was used to screen the planet during the greater part of the long exposure on the satellite, a series of very short exposures being given at intervals of 20 seconds. The shutter is pivoted, and is worked by a small electro-magnet arranged to give exposures automatically at the necessary intervals of time.

The zero of position angle was obtained by stopping the clock and giving a short supplementary exposure. Generally, several short exposures were given, the clock being put out of gear for a short time between successive exposures, so that each photograph contains in addition to the long exposure image of the satellite a series of short exposure images of Neptune.

The photographs were measured in a position micrometer in direct and reversed positions by Mr. Davidson and Mr. Melotte, the means of the measures being taken.

Corrections for refraction have not been applied separately to the distances or angles. These will in all cases be less than $0''.01$. The mean effects of refraction and aberration have, however, been included in the adopted value of the scale.

The value of $1^{\text{rev.}}$ of the screw of the measuring micrometer was determined in terms of a *réseau* interval of one of the *réseaux* used in photographic work with the Thompson and Astrographic Equatorials. For photographs with the 26-inch refractor cleared of the effect of differential refraction and aberration, the *réseau* interval adopted was $150''.92$. The correction for differential refraction on these photographs of Neptune varies from $+ 0''.042$ to $+ 0''.084$, and for aberration from $+ 0''.007$ to $- 0''.014$. A mean total correction of $+ 0''.06$ has been adopted, giving the *réseau* interval = $150''.98$. This value of the scale has been used for all the photographs.

The tabular positions were computed from the data given in the *Connaissance des Temps*, based on the elements given by Dr. Hermann Struve. The eccentricity of the orbit has been neglected, owing to the uncertainty of the present position of the periastron.

W. H. M. CHRISTIE.

*Royal Observatory, Greenwich,
1910 April.*

TRANSIT-CIRCLE TABLES, 1908.

ROYAL OBSERVATORY, GREENWICH.

TRANSIT-CIRCLE TABLES, 1908.

Equatorial Intervals of the Transit Wires.

The following are the intervals from the middle wire, determined from observations of close polar stars, which were in use throughout the year (adopted value of 1^{rev} of the Transit-micrometer = $36''\cdot97$):—

For Wire M = +	$44^{\text{s}}\cdot20$	For Wire T =	$0^{\text{s}}\cdot00$
N = +	$29^{\text{s}}\cdot73$	U = -	$7^{\text{s}}\cdot44$
O = +	$14^{\text{s}}\cdot94$	V = -	$9^{\text{s}}\cdot89$
P = +	$12^{\text{s}}\cdot39$	W = -	$12^{\text{s}}\cdot31$
Q = +	$9^{\text{s}}\cdot92$	X = -	$14^{\text{s}}\cdot80$
R = +	$4^{\text{s}}\cdot88$	Y = -	$29^{\text{s}}\cdot59$
S = +	$2^{\text{s}}\cdot41$	Z = -	$44^{\text{s}}\cdot35$

Apparent Correction to adopted Level-Errors, deduced from a Comparison of Reflexion and Direct Observations of Right Ascension, made in the Year 1908.

Name of Star.	Approx. N.P.D.	Approx. R.A.	Seconds of		R. - D.	Deduced Correction to Level-Error.	Number of Observations.		Weight.
			R.	D.			R.	D.	
ϵ Cephei S.P.....	$-24^{\circ}\cdot16$	$22^{\text{h}}\cdot46^{\text{m}}$	$28^{\text{s}}\cdot150$	$28^{\text{s}}\cdot250$	$-0^{\text{s}}\cdot100$	$+0^{\text{s}}\cdot66$	1	1	1
3 Draconis S.P.....	$-22^{\circ}\cdot45$	$11^{\text{h}}\cdot37^{\text{m}}$	$27^{\text{s}}\cdot450$	$27^{\text{s}}\cdot575$	$-0^{\text{s}}\cdot125$	$+0^{\text{s}}\cdot74$	1	2	1
ϵ Draconis S.P.....	$-19^{\circ}\cdot58$	$19^{\text{h}}\cdot48^{\text{m}}$	$28^{\text{s}}\cdot740$	$29^{\text{s}}\cdot125$	$-0^{\text{s}}\cdot385$	$+1^{\text{s}}\cdot85$	1	2	1
11 Ursæ Minoris S.P.....	$-17^{\circ}\cdot51$	$15^{\text{h}}\cdot17^{\text{m}}$	$9^{\text{s}}\cdot570$	$9^{\text{s}}\cdot370$	$+0^{\text{s}}\cdot200$	$-0^{\text{s}}\cdot82$	1	1	1
Bradley 348 S.P.....	$-17^{\circ}\cdot34$	$2^{\text{h}}\cdot29^{\text{m}}$	$26^{\text{s}}\cdot850$	$26^{\text{s}}\cdot770$	$+0^{\text{s}}\cdot080$	$-0^{\text{s}}\cdot32$	1	1	1
α Vulpeculæ.....	$65^{\circ}\cdot31$	$19^{\text{h}}\cdot25^{\text{m}}$	$57^{\text{s}}\cdot660$	$57^{\text{s}}\cdot630$	$+0^{\text{s}}\cdot030$	$+0^{\text{s}}\cdot23$	1	6	3
γ Geminorum.....	$69^{\circ}\cdot44$	$6^{\text{h}}\cdot23^{\text{m}}$	$37^{\text{s}}\cdot170$	$37^{\text{s}}\cdot200$	$-0^{\text{s}}\cdot030$	$-0^{\text{s}}\cdot24$	1	4	3
Mayer 136.....	$72^{\circ}\cdot56$	$3^{\text{h}}\cdot48^{\text{m}}$	$1^{\text{s}}\cdot120$	$0^{\text{s}}\cdot940$	$+0^{\text{s}}\cdot180$	$+1^{\text{s}}\cdot54$	1	1	2
70 Virginis.....	$75^{\circ}\cdot14$	$13^{\text{h}}\cdot24^{\text{m}}$	$1^{\text{s}}\cdot660$	$1^{\text{s}}\cdot700$	$-0^{\text{s}}\cdot040$	$-0^{\text{s}}\cdot36$	1	1	2
α Ophiuchi.....	$77^{\circ}\cdot22$	$17^{\text{h}}\cdot30^{\text{m}}$	$45^{\text{s}}\cdot360$	$45^{\text{s}}\cdot424$	$-0^{\text{s}}\cdot064$	$-0^{\text{s}}\cdot59$	1	14	2
α Cancri.....	$77^{\circ}\cdot48$	$8^{\text{h}}\cdot53^{\text{m}}$	$33^{\text{s}}\cdot960$	$34^{\text{s}}\cdot000$	$-0^{\text{s}}\cdot040$	$-0^{\text{s}}\cdot37$	1	4	2
β Cancri.....	$80^{\circ}\cdot32$	$8^{\text{h}}\cdot11^{\text{m}}$	$38^{\text{s}}\cdot170$	$38^{\text{s}}\cdot135$	$+0^{\text{s}}\cdot035$	$+0^{\text{s}}\cdot36$	1	2	1
σ Virginis.....	$84^{\circ}\cdot3$	$13^{\text{h}}\cdot13^{\text{m}}$	$3^{\text{s}}\cdot580$	$3^{\text{s}}\cdot640$	$-0^{\text{s}}\cdot060$	$-0^{\text{s}}\cdot63$	1	1	1
β Ophiuchi.....	$85^{\circ}\cdot24$	$17^{\text{h}}\cdot39^{\text{m}}$	$1^{\text{s}}\cdot470$	$1^{\text{s}}\cdot597$	$-0^{\text{s}}\cdot127$	$-1^{\text{s}}\cdot36$	1	7	2
τ Virginis.....	$88^{\circ}\cdot1$	$13^{\text{h}}\cdot57^{\text{m}}$	$3^{\text{s}}\cdot900$	$3^{\text{s}}\cdot912$	$-0^{\text{s}}\cdot012$	$-0^{\text{s}}\cdot14$	1	6	2
θ Aquilæ.....	$91^{\circ}\cdot5$	$20^{\text{h}}\cdot6^{\text{m}}$	$39^{\text{s}}\cdot710$	$39^{\text{s}}\cdot739$	$-0^{\text{s}}\cdot029$	$-0^{\text{s}}\cdot35$	1	10	2
α Hydræ.....	$98^{\circ}\cdot16$	$9^{\text{h}}\cdot23^{\text{m}}$	$10^{\text{s}}\cdot010$	$9^{\text{s}}\cdot883$	$+0^{\text{s}}\cdot127$	$+1^{\text{s}}\cdot89$	1	3	2
Spica.....	$100^{\circ}\cdot41$	$13^{\text{h}}\cdot20^{\text{m}}$	$27^{\text{s}}\cdot070$	$26^{\text{s}}\cdot985$	$+0^{\text{s}}\cdot085$	$+1^{\text{s}}\cdot32$	1	6	1
2 Libræ.....	$101^{\circ}\cdot18$	$14^{\text{h}}\cdot19^{\text{m}}$	$34^{\text{s}}\cdot960$	$34^{\text{s}}\cdot970$	$-0^{\text{s}}\cdot010$	$-0^{\text{s}}\cdot16$	1	1	1

TRANSIT-CIRCLE TABLES.

lxxvii

The weights used have been determined as follows:—

Putting m and n for the number of reflexion and direct observations respectively, the weight to be given to each star, in deducing the mean correction to level-error, is $\frac{4mn}{m+n+\frac{1}{5}mn} \cos Z.D.$, which has been adopted for use to the nearest integer.

Combining these results with the weights attached to them, we have:—

From 14 South Stars, Mean Correction to Level-Error + 0".05, Weight 26.

„ 5 North Stars, „ „ „ + 0".42, „ 5.

Factors for Instrumental-Errors.

N.P.D.	Collimation.	Level.	Azimuth.	N.P.D.	Collimation.	Level.	Azimuth.
	$\frac{1}{15 \sin N.P.D.}$	$\frac{\cos Z.D.}{15 \sin N.P.D.}$	$\frac{\sin Z.D.}{15 \sin N.P.D.}$		$\frac{1}{15 \sin N.P.D.}$	$\frac{\cos Z.D.}{15 \sin N.P.D.}$	$\frac{\sin Z.D.}{15 \sin N.P.D.}$
-45. 0	-0.094	-0.011	+0.094	-17. 30	-0.222	-0.124	+0.184
-44. 0	-0.096	-0.013	+0.095	-17. 0	-0.228	-0.129	+0.188
-43. 0	-0.098	-0.014	+0.096	-16. 30	-0.235	-0.134	+0.192
-42. 0	-0.100	-0.016	+0.098	-16. 0	-0.242	-0.140	+0.197
-41. 0	-0.102	-0.018	+0.100	-15. 30	-0.249	-0.146	+0.202
-40. 0	-0.104	-0.021	+0.102	-15. 0	-0.257	-0.153	+0.207
-39. 0	-0.106	-0.023	+0.104	-14. 30	-0.266	-0.160	+0.213
-38. 0	-0.108	-0.025	+0.105	-14. 0	-0.276	-0.168	+0.219
-37. 0	-0.111	-0.027	+0.107	-13. 30	-0.286	-0.176	+0.225
-36. 0	-0.113	-0.030	+0.109	-13. 0	-0.296	-0.184	+0.232
-35. 0	-0.116	-0.033	+0.111	-12. 30	-0.308	-0.194	+0.240
-34. 0	-0.119	-0.036	+0.114	-12. 0	-0.321	-0.204	+0.248
-33. 0	-0.122	-0.039	+0.116	-11. 30	-0.334	-0.214	+0.256
-32. 0	-0.126	-0.042	+0.119	-11. 0	-0.349	-0.227	+0.266
-31. 0	-0.129	-0.045	+0.121	-10. 30	-0.366	-0.240	+0.276
-30. 0	-0.133	-0.049	+0.124	-10. 0	-0.384	-0.254	+0.288
-29. 0	-0.138	-0.053	+0.127	- 9. 30	-0.404	-0.270	+0.300
-28. 0	-0.142	-0.057	+0.130	- 9. 0	-0.426	-0.288	+0.314
-27. 0	-0.147	-0.061	+0.133	- 8. 30	-0.451	-0.308	+0.330
-26. 0	-0.152	-0.065	+0.137	- 8. 0	-0.479	-0.330	+0.348
-25. 0	-0.158	-0.070	+0.141	- 7. 30	-0.511	-0.355	+0.368
-24. 0	-0.164	-0.076	+0.145	- 7. 0	-0.547	-0.383	+0.390
-23. 0	-0.171	-0.082	+0.150	- 6. 30	-0.589	-0.416	+0.417
-22. 0	-0.178	-0.088	+0.155	- 6. 0	-0.638	-0.455	+0.447
-21. 0	-0.186	-0.094	+0.160				
-20. 0	-0.195	-0.102	+0.166	6. 0	+0.638	+0.538	-0.343
-19. 30	-0.200	-0.106	+0.170	6. 30	+0.589	+0.499	-0.312
-19. 0	-0.205	-0.110	+0.173	7. 0	+0.547	+0.466	-0.286
-18. 30	-0.210	-0.114	+0.176	7. 30	+0.511	+0.438	-0.263
-18. 0	-0.216	-0.119	+0.180	8. 0	+0.479	+0.413	-0.243
-17. 30	-0.222	-0.124	+0.184	8. 30	+0.451	+0.391	-0.226

lxxviii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Factors for Instrumental-Errors—continued.

N.P.D.	Collimation.	Level.	Azimuth.	N.P.D.	Collimation.	Level.	Azimuth.
	$\frac{1}{15 \sin N.P.D.}$	$\frac{\cos Z.D.}{15 \sin N.P.D.}$	$\frac{\sin Z.D.}{15 \sin N.P.D.}$		$\frac{1}{15 \sin N.P.D.}$	$\frac{\cos Z.D.}{15 \sin N.P.D.}$	$\frac{\sin Z.D.}{15 \sin N.P.D.}$
8. 30	+0.451	+0.391	-0.226	44. 0	+0.096	+0.096	+0.009
9. 0	+0.426	+0.371	-0.210	45. 0	+0.094	+0.094	+0.011
9. 30	+0.404	+0.353	-0.196	46. 0	+0.093	+0.092	+0.012
10. 0	+0.384	+0.337	-0.183	47. 0	+0.091	+0.090	+0.013
10. 30	+0.366	+0.323	-0.172	48. 0	+0.090	+0.088	+0.015
11. 0	+0.349	+0.310	-0.161	49. 0	+0.088	+0.087	+0.016
11. 30	+0.334	+0.298	-0.152	50. 0	+0.087	+0.085	+0.017
12. 0	+0.321	+0.287	-0.143	51. 0	+0.086	+0.084	+0.018
12. 30	+0.308	+0.277	-0.135	52. 0	+0.085	+0.082	+0.020
13. 0	+0.296	+0.267	-0.128	53. 0	+0.083	+0.081	+0.021
13. 30	+0.286	+0.259	-0.122	54. 0	+0.082	+0.080	+0.022
14. 0	+0.276	+0.251	-0.116	55. 0	+0.081	+0.078	+0.023
14. 30	+0.266	+0.244	-0.111	56. 0	+0.080	+0.077	+0.024
15. 0	+0.257	+0.236	-0.105	57. 0	+0.080	+0.075	+0.025
15. 30	+0.249	+0.230	-0.099	58. 0	+0.079	+0.074	+0.026
16. 0	+0.242	+0.224	-0.093	59. 0	+0.078	+0.073	+0.027
16. 30	+0.235	+0.218	-0.088	60. 0	+0.077	+0.072	+0.028
17. 0	+0.228	+0.212	-0.084	61. 0	+0.076	+0.070	+0.029
17. 30	+0.222	+0.207	-0.080	62. 0	+0.076	+0.069	+0.030
18. 0	+0.216	+0.202	-0.076	63. 0	+0.075	+0.068	+0.031
18. 30	+0.210	+0.197	-0.072	64. 0	+0.074	+0.067	+0.032
19. 0	+0.205	+0.193	-0.068	65. 0	+0.073	+0.066	+0.033
19. 30	+0.200	+0.189	-0.065	66. 0	+0.073	+0.065	+0.034
20. 0	+0.195	+0.185	-0.062	67. 0	+0.072	+0.064	+0.035
21. 0	+0.186	+0.177	-0.056	68. 0	+0.072	+0.063	+0.035
22. 0	+0.178	+0.171	-0.051	69. 0	+0.071	+0.061	+0.036
23. 0	+0.171	+0.164	-0.046	70. 0	+0.071	+0.060	+0.037
24. 0	+0.164	+0.159	-0.041	71. 0	+0.071	+0.059	+0.038
25. 0	+0.158	+0.153	-0.037	72. 0	+0.070	+0.058	+0.039
26. 0	+0.152	+0.148	-0.033	73. 0	+0.070	+0.057	+0.039
27. 0	+0.147	+0.144	-0.029	74. 0	+0.069	+0.056	+0.040
28. 0	+0.142	+0.140	-0.026	75. 0	+0.069	+0.055	+0.041
29. 0	+0.138	+0.136	-0.023	76. 0	+0.069	+0.054	+0.042
30. 0	+0.133	+0.131	-0.020	77. 0	+0.068	+0.053	+0.043
31. 0	+0.129	+0.128	-0.017	78. 0	+0.068	+0.053	+0.043
32. 0	+0.126	+0.125	-0.014	79. 0	+0.068	+0.052	+0.044
33. 0	+0.122	+0.122	-0.012	80. 0	+0.068	+0.051	+0.045
34. 0	+0.119	+0.119	-0.009	81. 0	+0.067	+0.050	+0.045
35. 0	+0.116	+0.116	-0.007	82. 0	+0.067	+0.049	+0.046
36. 0	+0.113	+0.113	-0.005	83. 0	+0.067	+0.048	+0.047
37. 0	+0.111	+0.111	-0.003	84. 0	+0.067	+0.047	+0.048
38. 0	+0.108	+0.108	-0.001	85. 0	+0.067	+0.046	+0.048
39. 0	+0.106	+0.106	+0.001	86. 0	+0.067	+0.045	+0.049
40. 0	+0.104	+0.104	+0.003	87. 0	+0.067	+0.044	+0.050
41. 0	+0.102	+0.101	+0.004	88. 0	+0.067	+0.043	+0.051
42. 0	+0.100	+0.099	+0.006	89. 0	+0.067	+0.042	+0.052
43. 0	+0.098	+0.097	+0.007	90. 0	+0.067	+0.041	+0.052
44. 0	+0.096	+0.096	+0.009	91. 0	+0.067	+0.040	+0.053

TRANSIT-CIRCLE TABLES.

lxxix

Factors for Instrumental-Errors—continued.

N.P.D.	Collimation.	Level.	Azimuth.	N.P.D.	Collimation.	Level.	Azimuth.
	$\frac{1}{15 \sin N.P.D.}$	$\frac{\cos Z.D.}{15 \sin N.P.D.}$	$\frac{\sin Z.D.}{15 \sin N.P.D.}$		$\frac{1}{15 \sin N.P.D.}$	$\frac{\cos Z.D.}{15 \sin N.P.D.}$	$\frac{\sin Z.D.}{15 \sin N.P.D.}$
91. 0	+0.067	+0.040	+0.053	109. 0	+0.071	+0.024	+0.066
92. 0	+0.067	+0.040	+0.054	110. 0	+0.071	+0.023	+0.067
93. 0	+0.067	+0.039	+0.054	111. 0	+0.072	+0.022	+0.068
94. 0	+0.067	+0.038	+0.055	112. 0	+0.072	+0.021	+0.069
95. 0	+0.067	+0.037	+0.056	113. 0	+0.073	+0.020	+0.070
96. 0	+0.067	+0.036	+0.057	114. 0	+0.073	+0.018	+0.071
97. 0	+0.067	+0.035	+0.057	115. 0	+0.074	+0.017	+0.072
98. 0	+0.067	+0.034	+0.058	116. 0	+0.075	+0.016	+0.072
99. 0	+0.067	+0.033	+0.059	117. 0	+0.075	+0.015	+0.073
100. 0	+0.068	+0.032	+0.060	118. 0	+0.076	+0.014	+0.074
101. 0	+0.068	+0.031	+0.060	119. 0	+0.076	+0.013	+0.075
102. 0	+0.068	+0.030	+0.061	120. 0	+0.077	+0.012	+0.076
103. 0	+0.068	+0.029	+0.062	121. 0	+0.078	+0.011	+0.077
104. 0	+0.069	+0.028	+0.063	122. 0	+0.079	+0.009	+0.078
105. 0	+0.069	+0.027	+0.063	123. 0	+0.080	+0.008	+0.079
106. 0	+0.069	+0.027	+0.064	124. 0	+0.080	+0.007	+0.080
107. 0	+0.070	+0.026	+0.065	125. 0	+0.081	+0.005	+0.081
108. 0	+0.070	+0.025	+0.066	126. 0	+0.082	+0.004	+0.082
109. 0	+0.071	+0.024	+0.066				

*Assumed Mean Right Ascensions of Clock Stars and Circumpolar Stars,
with the Corrections to the R.A. of the Nautical Almanac, for 1908.0.*

Star's Name.	Mag.	Assumed Mean R.A. 1908.0.	Correction to N.A.	Approx. N.P.D.	Star's Name.	Mag.	Assumed Mean R.A. 1908.0.	Correction to N.A.	Approx. N.P.D.
α Andromedæ...	2.1	h m s 0. 3. 37.768	— s 0.014	61. 25	α Piscium.....	4.4	h m s 1. 40. 32.026	— s 0.008	81. 19
γ Pegasi.....	3.0	0. 8. 29.812	— 0.006	75. 20	β Arietis.....	2.8	1. 49. 33.314	+ 0.027	69. 38
ϵ Ceti.....	3.6	0. 14. 44.439	— 0.007	99. 20	α Arietis.....	2.0	2. 1. 59.056	+ 0.007	66. 58
44 Piscium.....	5.8	0. 20. 41.165	— 0.003	88. 34	ξ^1 Ceti.....	4.4	2. 8. 7.304	— 0.016	81. 35
12 Ceti.....	6.2	0. 25. 20.625	— 0.011	94. 28	67 Ceti.....	5.5	2. 12. 23.629	+ 0.010	96. 51
ϵ Andromedæ...	4.6	0. 33. 41.447	— 0.031	61. 11	ξ^2 Ceti.....	4.4	2. 23. 15.945	+ 0.003	81. 57
β Ceti.....	2.1	0. 38. 58.336	+ 0.006	108. 29	ν Ceti.....	4.9	2. 31. 2.685	+ 0.022	84. 48
δ Piscium.....	4.6	0. 43. 54.471	— 0.011	82. 55	δ Ceti.....	4.1	2. 34. 45.929	— 0.022	90. 4
20 Ceti.....	5.0	0. 48. 18.336	+ 0.037	91. 39	γ^2 Ceti.....	3.0	2. 38. 31.941	+ 0.016	87. 9
μ Andromedæ...	3.9	0. 51. 38.532	— 0.045	52. 0	σ Arietis.....	5.5	2. 46. 24.658	— 0.001	75. 18
ϵ Piscium.....	4.5	0. 58. 10.032	+ 0.001	82. 36	ϵ Arietis.....	4.6	2. 53. 56.911	0.000	69. 2
β Andromedæ...	2.2	1. 4. 34.614	+ 0.002	54. 52	α Ceti.....	2.7	2. 57. 28.127	+ 0.001	86. 16
ζ^1 Piscium.....	5.4	1. 8. 55.379	— 0.033	82. 55	δ Arietis.....	4.5	3. 6. 21.934	— 0.023	70. 37
θ Ceti.....	3.8	1. 19. 25.468	+ 0.003	98. 39	τ^1 Arietis.....	5.2	3. 15. 54.784	— 0.014	69. 11
Polaris.....	2.2	1. 26. 1.739	+ 0.147	1. 11	α Tauri.....	3.8	3. 19. 51.635	— 0.002	81. 18
η Piscium.....	3.7	1. 26. 33.482	— 0.004	75. 8	f Tauri.....	4.3	3. 25. 47.494	— 0.032	77. 23
ν Piscium.....	4.7	1. 36. 38.523	— 0.019	84. 59	ϵ Eridani.....	3.7	3. 28. 35.701	— 0.015	99. 46

LXXX. INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Assumed Mean Right Ascensions of Clock Stars and Circumpolar Stars, with the Corrections to the R.A. of the Nautical Almanac, for 1908.0—continued.

Star's Name.	Mag.	Assumed Mean R.A. 1908.0.	Correction to N.A.	Approx. N.P.D.	Star's Name.	Mag.	Assumed Mean R.A. 1908.0.	Correction to N.A.	Approx. N.P.D.
		^h ^m ^s	^s	[°] ['] ^{''}			^h ^m ^s	^s	[°] ['] ^{''}
11 Tauri.....	6.5	3. 35. 16.451	- 0.014	64. 58	ε Hydræ.....	3.8	8. 41. 54.313	- 0.008	83. 15
δ Eridani.....	3.7	3. 38. 50.403	- 0.036	100. 4	α Cancri.....	4.3	8. 53. 27.417	- 0.013	77. 47
η Tauri.....	3.0	3. 42. 0.794	+ 0.002	66. 11	κ Cancri.....	5.0	9. 2. 45.926	- 0.022	78. 58
γ ¹ Eridani.....	3.0	3. 53. 44.184	- 0.021	103. 46	83 Cancri.....	6.6	9. 13. 50.911	- 0.035	71. 54
A ¹ Tauri.....	4.4	3. 59. 15.230	- 0.023	68. 10	α Hydræ.....	2.0	9. 23. 4.014	+ 0.002	98. 16
ω ¹ Tauri.....	5.8	4. 3. 48.223	- 0.051	70. 38	ξ Leonis.....	5.2	9. 26. 59.313	+ 0.001	78. 18
ο ¹ Eridani.....	4.1	4. 7. 22.401	- 0.043	97. 5	ο Leonis.....	3.8	9. 36. 14.515	- 0.008	79. 41
γ Tauri.....	3.9	4. 14. 33.361	- 0.015	74. 36	ε Leonis.....	3.1	9. 40. 37.898	+ 0.008	65. 48
ε Tauri.....	3.7	4. 23. 14.565	- 0.017	71. 1	μ Leonis.....	4.1	9. 47. 32.049	+ 0.052	63. 34
Aldebaran.....	1.0	4. 30. 38.415	+ 0.014	73. 41	π Leonis.....	5.0	9. 55. 21.168	0.000	81. 31
τ Tauri.....	4.4	4. 36. 43.292	- 0.015	67. 13	Regulus.....	1.4	10. 3. 28.455	+ 0.020	77. 35
μ Eridani.....	4.3	4. 40. 54.067	- 0.037	93. 25	γ ¹ Leonis.....	2.0	10. 14. 54.159	+ 0.028	69. 42
ι Aurigæ.....	2.7	4. 51. 0.010	- 0.015	56. 59	μ Hydræ.....	4.1	10. 21. 38.411	- 0.022	106. 22
ε Leporis.....	3.3	5. 1. 33.955	- 0.012	112. 30	ρ Leonis.....	4.0	10. 27. 58.099	- 0.004	80. 13
Rigel.....	0.3	5. 10. 6.962	+ 0.010	98. 18	34 Sextantis.....	6.9	10. 37. 52.485	- 0.013	85. 56
β Tauri.....	1.9	5. 20. 28.513	- 0.003	61. 28	l Leonis.....	5.3	10. 44. 25.365	- 0.011	78. 58
δ Orionis.....	2.4	5. 27. 18.361	+ 0.001	90. 22	d Leonis.....	5.0	10. 55. 48.609	+ 0.022	85. 53
α Leporis.....	2.7	5. 28. 40.295	- 0.051	107. 53	χ Leonis.....	4.7	11. 0. 16.331	- 0.008	82. 10
ε Orionis.....	1.8	5. 31. 32.674	- 0.010	91. 16	δ Leonis.....	2.8	11. 9. 13.049	- 0.015	68. 58
α Columbae.....	2.7	5. 36. 19.027	- 0.030	124. 7	δ Crateris.....	3.9	11. 14. 44.412	+ 0.012	104. 17
κ Orionis.....	2.2	5. 43. 23.569	- 0.014	99. 42	τ Leonis.....	5.1	11. 23. 12.384	0.000	86. 38
α Orionis.....	1.4	5. 50. 11.461	+ 0.008	82. 37	ν Leonis.....	4.5	11. 32. 14.293	- 0.004	90. 19
ι Geminorum.....	4.3	5. 58. 31.664	- 0.015	66. 44	β Leonis.....	2.2	11. 44. 22.087	- 0.003	74. 55
ν Orionis.....	4.4	6. 2. 19.125	- 0.048	75. 13	β Virginis.....	3.7	11. 45. 54.187	+ 0.003	87. 43
η Geminorum.....	3.2	6. 9. 19.455	- 0.039	67. 28	π Virginis.....	4.4	11. 56. 9.538	+ 0.026	82. 52
μ Geminorum.....	3.2	6. 17. 23.692	- 0.018	67. 26	ο Virginis.....	4.3	12. 0. 31.405	+ 0.011	80. 45
β Canis Majoris.....	2.0	6. 18. 38.869	- 0.012	107. 55	ε Corvi.....	3.1	12. 5. 23.477	- 0.002	112. 6
ν Geminorum.....	4.0	6. 23. 30.046	+ 0.008	69. 44	Bradley 1672.....	6.3	12. 14. 25.788	+ 0.460	1. 47
γ Geminorum.....	2.0	6. 32. 23.862	+ 0.002	73. 31	η Virginis.....	4.0	12. 15. 11.930	- 0.014	90. 9
ξ Geminorum.....	3.4	6. 40. 7.566	- 0.014	77. 0	82 Corvi.....	3.0	12. 25. 6.123	- 0.034	106. 0
θ Canis Majoris.....	4.2	6. 49. 54.913	- 0.043	101. 55	β Corvi.....	2.8	12. 29. 33.127	+ 0.027	112. 53
ε Canis Majoris.....	1.5	6. 55. 0.585	- 0.015	118. 51	ρ Virginis.....	5.1	12. 37. 13.698	- 0.015	79. 15
Cephei 51.....	5.3	6. 57. 40.712	+ 0.394	2. 48	35 Virginis.....	6.9	12. 43. 10.338	- 0.004	85. 55
ζ Geminorum.....	3.7	6. 58. 39.173	- 0.028	69. 18	31 Comæ.....	5.0	12. 47. 13.113	+ 0.024	61. 58
γ Canis Majoris.....	4.1	6. 59. 35.756	- 0.031	105. 30	δ Virginis.....	3.7	12. 50. 58.134	+ 0.012	86. 6
51 Geminorum.....	5.4	7. 8. 5.351	- 0.033	73. 41	ε Virginis.....	3.0	12. 57. 35.848	+ 0.013	78. 33
δ Geminorum.....	3.2	7. 14. 37.770	- 0.035	67. 51	θ Virginis.....	4.4	13. 5. 11.107	- 0.001	95. 3
β Canis Minoris.....	3.1	7. 22. 9.740	- 0.009	81. 31	Spica.....	1.2	13. 20. 20.691	+ 0.012	100. 41
Castor.....	2.7	7. 28. 43.921	+ 0.021	57. 55	ζ Virginis.....	3.5	13. 30. 0.263	+ 0.017	90. 8
Procyon.....	0.5	7. 34. 29.190	+ 0.020	84. 32	m Virginis.....	5.3	13. 36. 46.908	+ 0.012	98. 14
Pollux.....	1.1	7. 39. 41.293	+ 0.009	61. 45	τ Boötis.....	4.5	13. 42. 53.435	+ 0.022	72. 5
ξ Argûs.....	3.4	7. 45. 25.477	- 0.026	114. 38	η Boötis.....	2.9	13. 50. 18.253	- 0.004	71. 8
6 Cancri.....	5.0	7. 57. 52.177	- 0.046	61. 57	τ Virginis.....	4.4	13. 56. 57.801	- 0.007	88. 1
ρ Argûs.....	2.9	8. 3. 37.546	+ 0.001	114. 2	94 Virginis.....	6.8	14. 1. 25.364	+ 0.016	98. 27
Groomb. 1119.....	7.1	8. 6. 35.770	- 0.213	1. 5	κ Virginis.....	4.3	14. 7. 59.176	- 0.011	99. 51
β Cancri.....	3.8	8. 11. 31.625	+ 0.012	80. 32	Arceturus.....	0.0	14. 11. 27.894	+ 0.012	70. 20
d ¹ Cancri.....	5.9	8. 18. 5.856	- 0.006	71. 22	f Boötis.....	5.4	14. 22. 10.593	+ 0.001	70. 22
η Cancri.....	5.5	8. 27. 23.428	- 0.009	69. 15	ρ Boötis.....	3.6	14. 27. 51.928	- 0.001	59. 14
γ Cancri.....	4.8	8. 37. 57.854	- 0.006	68. 12	ε ² Boötis.....	3.0	14. 40. 58.170	+ 0.018	62. 32

TRANSIT-CIRCLE TABLES.

lxxxix

Assumed Mean Right Ascensions of Clock Stars and Circumpolar Stars, with the Corrections to the R.A. of the Nautical Almanac, for 1908.0—continued.

Star's Name.	Mag.	Assumed Mean R.A. 1908.0.	Correction to N.A.	Approx. N.P.D.	Star's Name.	Mag.	Assumed Mean R.A. 1908.0.	Correction to N.A.	Approx. N.P.D.
		h m s	s	° '			h m s	s	° '
α Libræ.....	3.0	14. 45. 47.182	- 0.006	105. 40	μ Aquilæ.....	4.7	19. 29. 35.723	- 0.010	82. 49
ξ^2 Libræ.....	5.8	14. 51. 46.445	+ 0.016	101. 2	h^2 Sagittarii.....	4.6	19. 31. 6.619	+ 0.040	115. 5
ψ Boötis.....	4.5	15. 0. 30.212	+ 0.010	62. 42	e^1 Sagittarii.....	5.6	19. 35. 27.219	+ 0.003	106. 30
Groomb. 2283...	7.1	15. 6. 38.397	- 0.075	2. 25	γ Aquilæ.....	2.8	19. 41. 53.168	+ 0.018	79. 37
ι^1 Libræ.....	4.9	15. 6. 58.462	- 0.007	109. 27	α Aquilæ.....	1.0	19. 46. 17.691	+ 0.011	81. 23
β Libræ.....	2.7	15. 12. 3.264	- 0.008	99. 3	β Aquilæ.....	4.0	19. 50. 47.664	+ 0.010	83. 49
σ^2 Libræ.....	6.3	15. 17. 53.773	- 0.001	104. 48	c Sagittarii.....	4.7	19. 57. 0.191	+ 0.024	117. 58
ζ^1 Libræ.....	6.2	15. 23. 3.941	- 0.008	106. 24	θ Aquilæ.....	3.4	20. 6. 33.523	+ 0.016	91. 6
α Coronæ.....	2.4	15. 30. 47.550	+ 0.013	62. 59	α^2 Capricorni.....	3.8	20. 12. 57.076	+ 0.006	102. 50
α Serpentis.....	2.7	15. 39. 44.146	+ 0.020	83. 17	β Capricorni.....	3.4	20. 15. 50.624	- 0.013	105. 4
ϵ Serpentis.....	3.7	15. 46. 13.739	+ 0.008	85. 15	ρ Capricorni.....	5.0	20. 23. 36.878	+ 0.012	108. 7
γ Serpentis.....	4.0	15. 52. 12.172	- 0.009	74. 2	ϵ Delphini.....	4.1	20. 28. 49.071	- 0.008	79. 1
β^1 Scorpii.....	2.0	16. 0. 5.113	+ 0.021	109. 33	α Delphini.....	4.0	20. 35. 21.894	- 0.013	74. 25
δ Ophiuchi.....	2.8	16. 9. 31.377	- 0.008	93. 27	ϵ Aquarii.....	3.8	20. 42. 41.829	+ 0.021	99. 50
γ Herculis.....	3.8	16. 17. 51.672	+ 0.004	70. 38	μ Aquarii.....	4.8	20. 47. 41.573	+ 0.009	99. 20
Antares.....	1.1	16. 23. 45.852	- 0.007	116. 14	32 Vulpeculæ....	5.1	20. 50. 38.335	+ 0.003	62. 18
λ Ophiuchi.....	4.0	16. 26. 16.328	- 0.013	87. 49	θ Capricorni.....	4.3	21. 0. 46.625	+ 0.005	107. 36
ζ Ophiuchi.....	2.8	16. 32. 5.484	- 0.005	100. 23	ζ Cygni.....	3.5	21. 9. 1.222	+ 0.014	60. 9
ζ Herculis.....	3.1	16. 37. 49.092	+ 0.026	58. 14	α Equulei.....	4.1	21. 11. 13.519	+ 0.007	85. 8
κ Ophiuchi.....	3.4	16. 53. 18.780	+ 0.008	80. 29	ϵ Capricorni.....	4.4	21. 17. 7.530	- 0.021	107. 14
ϵ Herculis.....	4.0	16. 56. 46.158	+ 0.004	58. 56	Groomb. 3548...	7.4	21. 18. 2.222	+ 0.259	3. 21
η Ophiuchi.....	2.6	17. 5. 6.032	+ 0.017	105. 37	β Aquarii.....	3.1	21. 26. 43.012	+ 0.010	95. 59
α^1 Herculis.....	3.1	17. 10. 27.120	- 0.001	75. 30	ϵ Aquarii.....	4.8	21. 32. 51.318	- 0.009	98. 16
θ Ophiuchi.....	3.4	17. 16. 21.501	+ 0.015	114. 54	ϵ Pegasi.....	2.4	21. 39. 40.061	+ 0.023	80. 33
σ Ophiuchi.....	4.4	17. 21. 56.974	+ 0.001	85. 47	δ Capricorni.....	3.0	21. 41. 57.878	+ 0.010	106. 33
α Ophiuchi.....	2.2	17. 30. 39.809	+ 0.005	77. 22	16 Pegasi.....	5.0	21. 48. 52.526	- 0.001	64. 30
β Ophiuchi.....	2.9	17. 38. 55.643	- 0.005	85. 24	α Aquarii.....	3.2	22. 1. 3.566	+ 0.010	90. 46
μ Herculis.....	3.5	17. 42. 51.437	- 0.013	62. 14	ϵ Pegasi.....	4.0	22. 2. 43.648	- 0.007	65. 6
89 Herculis.....	5.6	17. 51. 42.484	- 0.041	63. 56	θ Aquarii.....	4.3	22. 11. 58.812	+ 0.024	98. 14
δ Ursæ Minoris..	4.3	18. 1. 57.181	+ 0.396	3. 23	γ Aquarii.....	4.1	22. 16. 54.300	+ 0.008	91. 51
72 Ophiuchi.....	3.8	18. 2. 59.282	+ 0.023	80. 27	σ Aquarii.....	4.8	22. 25. 46.760	- 0.035	101. 9
μ Sagittarii.....	4.1	18. 8. 15.681	+ 0.021	111. 5	η Aquarii.....	4.2	22. 30. 37.760	+ 0.006	90. 36
η Serpentis.....	3.4	18. 16. 32.948	+ 0.025	92. 55	ζ Pegasi.....	3.6	22. 36. 52.395	- 0.008	79. 39
λ Sagittarii.....	3.1	18. 22. 17.568	- 0.019	115. 28	μ Pegasi.....	3.7	22. 45. 33.702	- 0.010	65. 53
α Lyræ.....	0.2	18. 33. 49.393	- 0.019	51. 18	λ Aquarii.....	3.8	22. 47. 48.944	+ 0.011	98. 4
2 Aquilæ.....	4.8	18. 37. 14.207	- 0.031	99. 8	Fomalhaut.....	1.3	22. 52. 34.192	+ 0.036	120. 7
β^1 Lyræ.....	3.4	18. 46. 41.000	+ 0.013	56. 45	α Pegasi.....	2.6	23. 0. 10.638	+ 0.011	75. 17
ϵ Aquilæ.....	4.1	18. 55. 26.822	+ 0.019	75. 3	γ Piscium.....	3.8	23. 12. 23.758	+ 0.011	87. 13
ζ Aquilæ.....	3.1	19. 1. 10.911	+ 0.025	76. 16	κ Piscium.....	5.0	23. 22. 12.969	- 0.013	89. 15
ψ Sagittarii.....	5.2	19. 9. 54.035	+ 0.030	115. 25	Bradley 3147...	5.6	23. 27. 47.255	+ 0.950	3. 12
λ Ursæ Minoris..	6.5	19. 13. 19.027	+ 0.434	1. 0	ι Piscium.....	4.3	23. 35. 13.076	+ 0.010	84. 52
ω Aquilæ.....	5.1	19. 13. 29.904	+ 0.011	78. 34	δ Sculptoris.....	4.6	23. 44. 8.169	+ 0.082	118. 38
δ Aquilæ.....	3.5	19. 20. 51.609	+ 0.015	87. 4	ω Piscium.....	4.2	23. 54. 35.183	0.000	83. 39
α Vulpeculæ.....	4.7	19. 24. 52.640	+ 0.025	65. 31	2 Ceti.....	4.6	23. 59. 1.673	+ 0.013	107. 51

NOTE.—The Right Ascensions are deduced from the "Standard Mean Right Ascensions of Clock Stars for 1900.0 based on Twelve-hour Groups," at the end of the *Introduction to the Second Nine-Year Catalogue for 1900*. The values of the Precession and Proper Motions are taken from Prof. Newcomb's *Catalogue of Fundamental Stars*.

The correction to the R.A. of Procyon for the effect of orbital motion from 1900 (the epoch of the *Nine-Year Catalogue*) to 1908.0 is -0.081. This correction is derived from Prof. Auwers' "Elements" (*Astronomische Nachrichten*, Nos. 1371, 1372, 1373).

The Right Ascensions of the *Nautical Almanac* for 1908.0 are derived from Prof. Newcomb's *Catalogue of Fundamental Stars*.

GREENWICH OBSERVATIONS, 1908.

1

lxxxii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Determination of the Personal Equation of the various Observers using the Chronographic Method of observing Transits.

The observers, in order of seniority, are :—

SE, H, AC, B, HF, W, JS, PM, E, RC, SD, S, BE, HA, GC, G, AW.

When a result depends on a single observation of one of the observers, half weight is given to it. Such cases are denoted by an asterisk.

Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.	Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.
Feb. 3	^h 1	SE AC	^s + 0.06	Apr. 3	^h 3	AC JS	^s - 0.32
June 28	1	AC SE	+ 0.08	May 1	3	JS AC	- 0.31
Apr. 8	1	SE B	+ 0.01	„ 9	2	AC JS	- 0.33
Feb. 3	2	JS SE	- 0.21	Feb. 3	5	AC E	+ 0.03
Feb. 3	6	SE E	+ 0.09	June 28	3	E AC	- 0.03
Apr. 8	4	SE E	+ 0.11	Jan. 18	4	RC AC	+ 0.10
June 28	3	E SE	+ 0.05	Feb. 3	2	RC AC	+ 0.07
Feb. 3	1	RC SE	+ 0.13	Apr. 3	1	AC RC	+ 0.08
June 4	2	SE RC	+ 0.18	May 1	2	RC AC	+ 0.09
Sept. 30	7	RC SE	+ 0.15	„ 9	0	RC AC	+ 0.05
Oct. 15	9	SE RC	+ 0.30	Sept. 2	1	RC AC	+ 0.04
Apr. 8	5	SE HA	- 0.24	„ 6	1	RC AC	+ 0.19
Sept. 30	7	HA SE	- 0.31	May 9	3	HA AC	- 0.29
Sept. 17	9	SE G	+ 0.06	Feb. 8	5	B W	- 0.03
Feb. 23	8	H W	+ 0.18	May 6	2	B W	- 0.05
Feb. 23	1	H JS	- 0.05	July 29	1	W B	- 0.03
Feb. 1	5	B AC	- 0.08	Feb. 1	4	B JS	- 0.27
„ 8	5	B AC	- 0.09	Apr. 8	5	B E	+ 0.10
Feb. 8	0	W AC	- 0.12	May 6	0	B E	+ 0.04
Feb. 1	2	JS AC	- 0.35	July 1	4	B E	+ 0.06
„ 3	3	JS AC	- 0.27	Oct. 7	1	B E	+ 0.12
				„ 14	10	B E	- 0.02*

TRANSIT-CIRCLE TABLES.

lxxxiii

Observations for Personal Equations—Chronographic—continued.

Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.	Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.
Nov. 25	^h 8	B E	^s - 0.02	Feb. 12	^h 6	H A H F	^s - 0.25
May 6	6	B R C	+ 0.05	Apr. 15	4	H A H F	- 0.12
Oct. 8	1	B R C	+ 0.12	June 29	1	H A H F	- 0.20
" 14	0	R C B	+ 0.20	Sept. 29	3	H A H F	- 0.16
" 24	1	R C B	+ 0.26	Oct. 1	2	H A H F	- 0.18
Nov. 25	2	B R C	+ 0.20	Dec. 16	0	G C H F	- 0.13
Oct. 14	1	B S D	+ 0.03	Feb. 12	4	G H F	- 0.08
July 29	2	S B	0.00	Jan. 3	12	W J S	- 0.34
July 1	1	B E B	0.00	Feb. 12	2	J S W	- 0.26
Apr. 8	4	H A B	- 0.25	" 23	7	J S W	- 0.23
May 6	0	H A B	- 0.19	Jan. 3	6	W E	+ 0.08
Oct. 8	1	H A B	- 0.23	Feb. 14	2	E W	+ 0.14
" 24	2	H A B	- 0.14	Mar. 13	1	E W	+ 0.16
Nov. 25	1	H A B	- 0.20	May 6	2	E W	+ 0.09
July 29	1	G C B	- 0.11	" 18	8	W E	+ 0.08
July 1	0	G B	+ 0.09	Sept. 25	2	E W	+ 0.04*
Feb. 12	0	W H F	- 0.06	" 29	3	E W	+ 0.08
Sept. 29	1	H F W	- 0.02	Nov. 8	12	W E	+ 0.09
Oct. 1	1	W H F	- 0.02	Feb. 12	7	W R C	+ 0.13
Feb. 12	2	J S H F	- 0.32	" 28	6	W R C	+ 0.17
June 29	3	E H F	+ 0.06	May 6	4	W R C	+ 0.10
Sept. 29	2	E H F	+ 0.06	Sept. 8	0	W R C	+ 0.20
Feb. 12	7	H F R C	+ 0.07	Oct. 5	9	W R C	+ 0.24
Apr. 15	0	H F R C	+ 0.16	" 21	8	W R C	+ 0.25
Sept. 7	2	S H F	+ 0.09	Dec. 7	10	R C W	+ 0.24
Jan. 25	1	H A H F	- 0.22	Mar. 9	2	S D W	+ 0.01
				Sept. 14	6	S D W	+ 0.04
				" 18	1	W S D	+ 0.02

Observations for Personal Equations—Chronographic—continued.

Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.	Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.
Jan. 3	^h 2	H A W	^s — 0'25	Feb. 13	^h 12	R C J S	^s + 0'29*
Feb. 12	6	H A W	— 0'19	„ 24	4	R C J S	+ 0'35
Apr. 16	3	H A W	— 0'14	Mar. 11	8	J S R C	+ 0'39
May 6	3	H A W	— 0'14	Apr. 3	2	R C J S	+ 0'40
Aug. 7	1	H A W	— 0'10	„ 13	3	J S R C	+ 0'33
„ 11	1	W H A	— 0'12	May 1	0	J S R C	+ 0'40
Sept. 14	10	H A W	— 0'15	„ 6	10	R C J S	+ 0'34
„ 29	4	H A W	— 0'14	„ 9	3	R C J S	+ 0'38
Oct. 1	1	H A W	— 0'16	Jan. 27	11	S D J S	+ 0'22
„ 21	3	H A W	— 0'28	Mar. 23	7	S D J S	+ 0'14
„ 29	1	H A W	— 0'17	Feb. 24	6	S J S	+ 0'41
Dec. 28	2	H A W	— 0'21	Jan. 27	8	H A J S	— 0'03
July 29	1	S W	+ 0'03	Feb. 12	4	H A J S	+ 0'07
„ 31	0	S W	+ 0'11	Mar. 11	1	H A J S	— 0'01
Aug. 11	2	W S	+ 0'13	„ 28	4	H A J S	+ 0'03
„ 14	0	S W	+ 0'11	May 9	5	H A J S	+ 0'04
Sept. 14	10	S W	+ 0'11	Feb. 12	2	G J S	+ 0'24
July 27	3	G C W	— 0'02*	Feb. 24	2	A W J S	+ 0'17
„ 29	1	G C W	— 0'08	June 25	0	P M E	+ 0'03
Aug. 11	1	W G C	— 0'07	June 12	1	P M S	+ 0'14
Feb. 12	4	G W	— 0'02	June 25	2	H A P M	— 0'29
Jan. 3	5	E J S	+ 0'42	Jan. 27	7	E R C	— 0'05*
„ 17	8	E J S	+ 0'42	Feb. 3	7	R C E	+ 0'04
„ 20	9	J S E	+ 0'43	May 6	6	E R C	+ 0'01
Feb. 3	8	J S E	+ 0'30	Oct. 14	10	R C E	+ 0'22*
„ 10	7	E J S	+ 0'35	Nov. 9	8	R C E	+ 0'03
Mar. 23	3	E J S	+ 0'16	„ 25	6	R C E	+ 0'04
Jan. 10	4	R C J S	+ 0'36	Jan. 27	0	S D E	— 0'05
„ 27	4	R C J S	+ 0'22*				
Feb. 3	1	J S R C	+ 0'34				
„ 12	9	J S R C	+ 0'39				

TRANSIT-CIRCLE TABLES.

lxxxv

Observations for Personal Equations—Chronographic—continued.

Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.	Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.
Mar. 23	^h 4	S D E	^s - 0.02	Oct. 2	^h 0	R C S D	^s - 0.18
Aug. 10	1	E S D	+ 0.07	,, 14	2	R C S D	- 0.17
Oct. 14	8	S D E	+ 0.05*	Dec. 10	3	S D R C	- 0.15
Aug. 15	8	S E	+ 0.05	Feb. 24	2	S R C	+ 0.06
Sept. 21	1	E S	+ 0.04	Aug. 27	4	S R C	- 0.10
Oct. 23	1	E S	- 0.03	Oct. 26	1	R C S	- 0.19
July 1	5	B E E	- 0.06	,, 30	3	S R C	- 0.21
Jan. 3	7	H A E	- 0.33	Nov. 23	1	R C S	- 0.10
,, 13	1	H A E	- 0.34	Dec. 7	0	R C S	+ 0.05
Apr. 8	9	H A E	- 0.35	,, 9	3	S R C	- 0.14
May 6	1	H A E	- 0.23	,, 14	7	S R C	- 0.19
June 19	2	H A E	- 0.18	Jan. 4	1	H A R C	- 0.37
,, 25	2	H A E	- 0.32	,, 14	3	H A R C	- 0.32
,, 29	2	E H A	- 0.26	Feb. 22	8	H A R C	- 0.39
Aug. 15	8	H A E	- 0.21	Mar. 11	8	H A R C	- 0.40
Sept. 1	1	H A E	- 0.20	,, 18	1	R C H A	- 0.36
,, 12	6	H A E	- 0.23	Apr. 15	4	H A R C	- 0.28
,, 29	1	H A E	- 0.22	May 6	7	H A R C	- 0.24
Oct. 3	5	H A E	- 0.24	,, 9	3	H A R C	- 0.34
,, 31	0	H A E	- 0.30	June 30	2	H A R C	- 0.35
Nov. 9	10	H A E	- 0.35	Aug. 21	1	R C H A	- 0.27
,, 25	9	H A E	- 0.36	Sept. 4	0	R C H A	- 0.26
,, 27	0	E H A	- 0.31	,, 10	2	H A R C	- 0.34
Mar. 5	1	G C E	- 0.11	,, 30	0	H A R C	- 0.46
June 19	5	E G C	- 0.09	Oct. 8	2	H A R C	- 0.35
Nov. 9	0	G C E	- 0.08	,, 12	2	H A R C	- 0.37
July 1	4	G E	+ 0.03	,, 21	10	H A R C	- 0.53
Jan. 27	7	S D R C	0.00*	,, 24	2	H A R C	- 0.40
July 2	4	S D R C	- 0.15	Nov. 9	2	H A R C	- 0.38
				,, 17	1	R C H A	- 0.44
				,, 25	3	H A R C	- 0.40
				Mar. 21	5	G C R C	- 0.20

lxxxvi INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Observations for Personal Equations—Chronographic—continued.

Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.	Day, 1908.	Interval between Means of Groups of Clock Stars.	Observers.	Excess of Clock Slow by Junior Observer.
May 23	^h 4	G C R C	^s — 0.21	Sept. 14	^h 1	H A S	^s — 0.26
June 14	1	R C G C	— 0.14	„ 15	5	H A S	— 0.30
Aug. 16	12	R C G C	— 0.27	Oct. 10	4	H A S	— 0.30
Nov. 9	7	R C G C	— 0.11	July 29	0	S G C	— 0.11
Feb. 12	11	G R C	— 0.15	Aug. 11	1	G C S	— 0.20
May 20	8	G R C	— 0.09	„ 15	4	G C S	— 0.23
„ 27	5	G R C	— 0.11	Oct. 13	1	S G C	— 0.12
June 17	7	G R C	— 0.09	Nov. 10	0	S G C	— 0.09
Feb. 24	2	R C A W	— 0.18	Feb. 24	4	S A W	— 0.24
June 18	5	S D S	+ 0.09*	July 1	5	B E G	+ 0.09
July 30	2	S S D	— 0.04	June 19	7	H A G C	+ 0.09
Sept. 14	4	S S D	+ 0.07	Aug. 11	1	H A G C	+ 0.05
May 19	3	S D H A	— 0.17	„ 15	4	G C H A	+ 0.03
Sept. 14	4	H A S D	— 0.19	„ 17	3	G C H A	+ 0.04
May 21	4	H A S	— 0.37	Nov. 9	9	H A G C	+ 0.27
Aug. 11	1	H A S	— 0.25	Feb. 12	1	H A G	+ 0.17
„ 15	0	S H A	— 0.26	Apr. 7	5	H A G	+ 0.21
				Sept. 28	1	H A G	+ 0.22

TRANSIT-CIRCLE TABLES.

lxxxvii

From these differences, if we form equations in which each observer's initial shall successively stand first, we obtain the following groups of equations:—

For S E.

$$\begin{aligned}
 4 (S E - A C) &= - 0^s.28 \\
 2 (S E - B) &= - 0^s.02 \\
 2 (S E - J S) &= + 0^s.42 \\
 6 (S E - E) &= - 0^s.50 \\
 8 (S E - R C) &= - 1^s.52 \\
 4 (S E - H A) &= + 1^s.10 \\
 2 (S E - G) &= - 0^s.12
 \end{aligned}$$

For H F.—continued.

$$\begin{aligned}
 4 (H F - E) &= - 0^s.24 \\
 4 (H F - R C) &= - 0^s.46 \\
 2 (H F - S) &= - 0^s.18 \\
 12 (H F - H A) &= + 2^s.26 \\
 2 (H F - G C) &= + 0^s.26 \\
 2 (H F - G) &= + 0^s.16
 \end{aligned}$$

For H.

$$\begin{aligned}
 2 (H - W) &= - 0^s.36 \\
 2 (H - J S) &= + 0^s.10
 \end{aligned}$$

For W.

$$\begin{aligned}
 2 (W - H) &= + 0^s.36 \\
 2 (W - A C) &= - 0^s.24 \\
 6 (W - B) &= - 0^s.22 \\
 6 (W - H F) &= - 0^s.20 \\
 6 (W - J S) &= + 1^s.66 \\
 15 (W - E) &= - 1^s.48 \\
 14 (W - R C) &= - 2^s.66 \\
 6 (W - S D) &= - 0^s.14 \\
 24 (W - H A) &= + 4^s.10 \\
 12 (W - S) &= - 0^s.98 \\
 5 (W - G C) &= + 0^s.32 \\
 2 (W - G) &= + 0^s.04
 \end{aligned}$$

For A C.

$$\begin{aligned}
 4 (A C - S E) &= + 0^s.28 \\
 4 (A C - B) &= + 0^s.34 \\
 2 (A C - W) &= + 0^s.24 \\
 10 (A C - J S) &= + 3^s.16 \\
 4 (A C - E) &= 0^s.00 \\
 14 (A C - R C) &= - 1^s.24 \\
 2 (A C - H A) &= + 0^s.58
 \end{aligned}$$

For J S.

$$\begin{aligned}
 2 (J S - S E) &= - 0^s.42 \\
 2 (J S - H) &= - 0^s.10 \\
 10 (J S - A C) &= - 3^s.16 \\
 2 (J S - B) &= - 0^s.54 \\
 2 (J S - H F) &= - 0^s.64 \\
 6 (J S - W) &= - 1^s.66 \\
 12 (J S - E) &= - 4^s.16 \\
 22 (J S - R C) &= - 7^s.87 \\
 4 (J S - S D) &= - 0^s.72 \\
 2 (J S - S) &= - 0^s.82 \\
 10 (J S - H A) &= - 0^s.20 \\
 2 (J S - G) &= - 0^s.48 \\
 2 (J S - A W) &= - 0^s.34
 \end{aligned}$$

For B.

$$\begin{aligned}
 2 (B - S E) &= + 0^s.02 \\
 4 (B - A C) &= - 0^s.34 \\
 6 (B - W) &= + 0^s.22 \\
 2 (B - J S) &= + 0^s.54 \\
 11 (B - E) &= - 0^s.58 \\
 10 (B - R C) &= - 1^s.66 \\
 2 (B - S D) &= - 0^s.06 \\
 2 (B - S) &= 0^s.00 \\
 2 (B - B E) &= 0^s.00 \\
 10 (B - H A) &= + 2^s.02 \\
 2 (B - G C) &= + 0^s.22 \\
 2 (B - G) &= - 0^s.18
 \end{aligned}$$

For H F.

$$\begin{aligned}
 6 (H F - W) &= + 0^s.20 \\
 2 (H F - J S) &= + 0^s.64
 \end{aligned}$$

For P M.

$$\begin{aligned}
 2 (P M - E) &= - 0^s.06 \\
 2 (P M - S) &= - 0^s.28 \\
 2 (P M - H A) &= + 0^s.58
 \end{aligned}$$

lxxxviii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

For E.

6 (E - SE)	= + 0 ^s .50
4 (E - AC)	= 0 ^s .00
11 (E - B)	= + 0 ^s .58
4 (E - HF)	= + 0 ^s .24
15 (E - W)	= + 1 ^s .48
12 (E - JS)	= + 4 ^s .16
2 (E - PM)	= + 0 ^s .06
10 (E - RC)	= - 0 ^s .41
7 (E - SD)	= - 0 ^s .05
6 (E - S)	= - 0 ^s .12
2 (E - BE)	= + 0 ^s .12
32 (E - HA)	= + 8 ^s .86
6 (E - GC)	= + 0 ^s .56
2 (E - G)	= - 0 ^s .06

For RC.

8 (RC - SE)	= + 1 ^s .52
14 (RC - AC)	= + 1 ^s .24
10 (RC - B)	= + 1 ^s .66
4 (RC - HF)	= + 0 ^s .46
14 (RC - W)	= + 2 ^s .66
22 (RC - JS)	= + 7 ^s .87
10 (RC - E)	= + 0 ^s .41
9 (RC - SD)	= + 1 ^s .30
16 (RC - S)	= + 1 ^s .64
40 (RC - HA)	= + 14 ^s .50
10 (RC - GC)	= + 1 ^s .86
8 (RC - G)	= + 0 ^s .88
2 (RC - AW)	= + 0 ^s .36

For S D.

2 (SD - B)	= + 0 ^s .06
6 (SD - W)	= + 0 ^s .14
4 (SD - JS)	= + 0 ^s .72
7 (SD - E)	= + 0 ^s .05
9 (SD - RC)	= - 1 ^s .30
5 (SD - S)	= - 0 ^s .15
4 (SD - HA)	= + 0 ^s .72

For S.

2 (S - B)	= 0 ^s .00
2 (S - HF)	= + 0 ^s .18
10 (S - W)	= + 0 ^s .98
2 (S - JS)	= + 0 ^s .82

For S—continued.

2 (S - PM)	= + 0 ^s .28
6 (S - E)	= + 0 ^s .12
16 (S - RC)	= - 1 ^s .64
5 (S - SD)	= + 0 ^s .15
12 (S - HA)	= + 3 ^s .48
10 (S - GC)	= + 1 ^s .56
2 (S - AW)	= + 0 ^s .48

For BE.

2 (BE - B)	= 0 ^s .00
2 (BE - E)	= - 0 ^s .12
2 (BE - G)	= - 0 ^s .18

For HA.

4 (HA - SE)	= - 1 ^s .10
2 (HA - AC)	= - 0 ^s .58
10 (HA - B)	= - 2 ^s .02
12 (HA - HF)	= - 2 ^s .26
24 (HA - W)	= - 4 ^s .10
10 (HA - JS)	= + 0 ^s .20
2 (HA - PM)	= - 0 ^s .58
32 (HA - E)	= - 8 ^s .86
40 (HA - RC)	= - 14 ^s .50
4 (HA - SD)	= - 0 ^s .72
12 (HA - S)	= - 3 ^s .48
10 (HA - GC)	= - 0 ^s .96
6 (HA - G)	= - 1 ^s .20

For GC.

2 (GC - B)	= - 0 ^s .22
2 (GC - HF)	= - 0 ^s .26
5 (GC - W)	= - 0 ^s .32
6 (GC - E)	= - 0 ^s .56
10 (GC - RC)	= - 1 ^s .86
10 (GC - S)	= - 1 ^s .50
10 (GC - HA)	= + 0 ^s .96

For G.

2 (G - SE)	= + 0 ^s .12
2 (G - B)	= + 0 ^s .18
2 (G - HF)	= - 0 ^s .16
2 (G - W)	= - 0 ^s .04
2 (G - JS)	= + 0 ^s .48
2 (G - E)	= + 0 ^s .06

TRANSIT-CIRCLE TABLES.

lxxxix

For G.—*continued*.

$$\begin{aligned} 8(G - RC) &= - 0^s.88 \\ 2(G - BE) &= + 0.18 \\ 6(G - HA) &= + 1.20 \end{aligned}$$

For A W.

$$\begin{aligned} 2(AW - JS) &= + 0^s.34 \\ 2(AW - RC) &= - 0.36 \\ 2(AW - S) &= - 0.48 \end{aligned}$$

Now, since one of the quantities denoted by S E, etc., must evidently remain indeterminate, we will refer all the rest to W as a standard; and putting $W = 0$, the initials S E, etc., will represent the value of "clock slow" which would be given by the observations of S E, etc., when that given by $W = 0$.

Then, adding all the equations in each, we get the following equations, in which each observer's personal equation is successively affected by a large coefficient:—

$$\begin{aligned} 28 \text{ S E}-4 \text{ A C}-2 \text{ B}-2 \text{ J S}-6 \text{ E}-8 \text{ R C}-4 \text{ H A}-2 \text{ G} &= - 0^s.92 \\ 4 \text{ H}-2 \text{ W}-2 \text{ J S} &= - 0.26 \\ 40 \text{ A C}-4 \text{ S E}-4 \text{ B}-2 \text{ W}-10 \text{ J S}-4 \text{ E}-14 \text{ R C}-2 \text{ H A} &= + 3.36 \\ 55 \text{ B}-2 \text{ S E}-4 \text{ A C}-6 \text{ W}-2 \text{ J S}-11 \text{ E}-10 \text{ R C}-2 \text{ S D}-2 \text{ S}-2 \text{ B E}-10 \\ \text{H A}-2 \text{ G C}-2 \text{ G} &= + 0.20 \\ 34 \text{ H F}-6 \text{ W}-2 \text{ J S}-4 \text{ E}-4 \text{ R C}-2 \text{ S}-12 \text{ H A}-2 \text{ G C}-2 \text{ G} &= + 2.64 \\ 98 \text{ W}-2 \text{ H}-2 \text{ A C}-6 \text{ B}-6 \text{ H F}-6 \text{ J S}-15 \text{ E}-14 \text{ R C}-6 \text{ S D}-10 \text{ S}-24 \\ \text{H A}-5 \text{ G C}-2 \text{ G} &= + 5.56 \\ 78 \text{ J S}-2 \text{ S E}-2 \text{ H}-10 \text{ A C}-2 \text{ B}-2 \text{ H F}-6 \text{ W}-12 \text{ E}-22 \text{ R C}-4 \text{ S D}-2 \text{ S} \\ -10 \text{ H A}-2 \text{ G}-2 \text{ A W} &= -21.11 \\ 6 \text{ P M}-2 \text{ E}-2 \text{ S}-2 \text{ H A} &= + 0.24 \\ 119 \text{ E}-6 \text{ S E}-4 \text{ A C}-11 \text{ B}-4 \text{ H F}-15 \text{ W}-12 \text{ J S}-2 \text{ P M}-10 \text{ R C}-7 \text{ S D} \\ -6 \text{ S}-2 \text{ B E}-32 \text{ H A}-6 \text{ G C}-2 \text{ G} &= +15.92 \\ 167 \text{ R C}-8 \text{ S E}-14 \text{ A C}-10 \text{ B}-4 \text{ H F}-14 \text{ W}-22 \text{ J S}-10 \text{ E}-9 \text{ S D}-16 \text{ S} \\ -40 \text{ H A}-10 \text{ G C}-8 \text{ G}-2 \text{ A W} &= +36.36 \\ 37 \text{ S D}-2 \text{ B}-6 \text{ W}-4 \text{ J S}-7 \text{ E}-9 \text{ R C}-5 \text{ S}-4 \text{ H A} &= + 0.24 \\ 69 \text{ S}-2 \text{ B}-2 \text{ H F}-10 \text{ W}-2 \text{ J S}-2 \text{ P M}-6 \text{ E}-16 \text{ R C}-5 \text{ S D}-12 \text{ H A} \\ -10 \text{ G C}-2 \text{ A W} &= + 6.35 \\ 6 \text{ B E}-2 \text{ B}-2 \text{ E}-2 \text{ G} &= - 0.30 \\ 168 \text{ H A}-4 \text{ S E}-2 \text{ A C}-10 \text{ B}-12 \text{ H F}-24 \text{ W}-10 \text{ J S}-2 \text{ P M}-32 \text{ E}-40 \\ \text{R C}-4 \text{ S D}-12 \text{ S}-10 \text{ G C}-6 \text{ G} &= -40.16 \\ 45 \text{ G C}-2 \text{ B}-2 \text{ H F}-5 \text{ W}-6 \text{ E}-10 \text{ R C}-10 \text{ S}-10 \text{ H A} &= - 3.76 \\ 28 \text{ G}-2 \text{ S E}-2 \text{ B}-2 \text{ H F}-2 \text{ W}-2 \text{ J S}-2 \text{ E}-8 \text{ R C}-2 \text{ B E}-6 \text{ H A} &= + 1.14 \\ 6 \text{ A W}-2 \text{ J S}-2 \text{ R C}-2 \text{ S} &= - 0.50 \end{aligned}$$

Solving these equations by successive approximations—that is, by first substituting in each equation the approximately known values for all the personal equations except that which has the largest coefficient, thence deducing the value of that one

GREENWICH OBSERVATIONS, 1908.

m

personal equation, then using that deduced value for substitution in the other equations and repeating the process, we get finally the following values:—

$$\begin{array}{llll}
 W - S E = -0.01 & W - H F = -0.04 & W - R C = -0.17 & W - H A = +0.18 \\
 W - H = +0.17 & W - J S = +0.22 & W - S D = -0.04 & W - G C = +0.05 \\
 W - A C = -0.09 & W - P M = -0.04 & W - S = -0.10 & W - G = -0.05 \\
 W - B = -0.02 & W - E = -0.09 & W - B E = 0.00 & W - A W = +0.07
 \end{array}$$

Determination of the Personal Equations of the Various Observers using the Eye-and-Ear Method of observing Transits as referred to the Standard Observer using the Chronographic Method.

From 1885 March 14 it has been the practice for the observer with the transit-circle to observe each night the transits of two clock-stars by the eye-and-ear method as well as with the chronographic method. The following table gives the comparison of the clock-errors deduced from these eye-and-ear transits with those deduced from the chronographic transits by the same observer, these latter being referred to the Standard Observer, Mr. Witchell, by applying the personal equations for chronographic transits found from the observations of 1908:—

Date of Observation, 1908.	Excess of Clock Slow by Eye-and-Ear Observer.	Date of Observation, 1908.	Excess of Clock Slow by Eye-and-Ear Observer.	Date of Observation, 1908.	Excess of Clock Slow by Eye-and-Ear Observer.
Observer W.		Observer J S.		Observer E.	
d h	s	d h	s	d h	s
Feb. 12. 12	+ 0.14	Jan. 2. 10	+ 0.17	Jan. 3. 13	+ 0.02
28. 11	+ 0.03	3. 18	+ 0.17	17. 13	+ 0.05
		12. 7	+ 0.20	27. 10	- 0.01
May 18. 13	+ 0.12	17. 17	+ 0.28	29. 10	+ 0.01
		20. 9	+ 0.07		
		22. 10	+ 0.02	Feb. 3. 16	+ 0.11
July 27. 11	+ 0.07			6. 7	+ 0.02
31. 11	+ 0.02	Feb. 1. 12	+ 0.01	7. 16	+ 0.08
		2. 11	+ 0.03	18. 13	+ 0.23
Aug. 7. 12	- 0.05	23. 10	+ 0.06		
		29. 9	+ 0.12	Mar. 13. 11	+ 0.15
Sept. 18. 10	- 0.12			23. 14	+ 0.06
29. 10	+ 0.09	Mar. 23. 17	- 0.01	Apr. 30. 12	+ 0.17
		28. 12	+ 0.10	May 4. 11	+ 0.16
Mean.....	+ 0.04	Mean.....	+ 0.10		

TRANSIT-CIRCLE TABLES.

xci

Observations for Personal Equations—Eye-and-Ear—continued.

Date of Observation, 1908.	Excess of Clock Slow by Eye-and-Ear Observer.	Date of Observation, 1908.	Excess of Clock Slow by Eye-and-Ear Observer.	Date of Observation, 1908.	Excess of Clock Slow by Eye-and-Ear Observer.
<i>Observer E.—continued.</i>		<i>Observer R C.—continued.</i>		<i>Observer R C.—continued.</i>	
d h	s	d h	s	d h	s
Aug. 10. 10	+ 0.12	Feb. 15. 12	+ 0.31	Sept. 2. 13	+ 0.19
Sept. 29. 11	+ 0.14	27. 12	+ 0.25	30. 13	+ 0.13
Oct. 3. 12	+ 0.18	Mar. 14. 12	+ 0.26	Nov. 17. 12	+ 0.13
Nov. 25. 13	+ 0.13	18. 13	+ 0.21	25. 12	+ 0.28
28. 10	+ 0.21	Apr. 3. 13	+ 0.18	Mean	+ 0.19
Mean	+ 0.10	15. 12	+ 0.23	<i>Observer H A.</i>	
<i>Observer R C.</i>		May 1. 13	+ 0.16		
		9. 12	+ 0.19		
		13. 13	+ 0.24		
<i>Observer R C.</i>		June 2. 13	+ 0.22	Jan. 4. 6	+ 0.06
		10. 13	+ 0.11	Mean	+ 0.06
Jan. 14. 12	+ 0.06	30. 14	+ 0.31		
		Aug. 25. 13	+ 0.03		
		29. 13	+ 0.12		

Collecting these results together, the following are the values of the corrections for personal equation for the observers using the eye-and-ear method referred to the Standard Observer using the Galvanic method :—

$$\begin{array}{lll}
 W_0 - W & = - 0.04^s & W_0 - E & = - 0.10^s & W_0 - H A & = - 0.06^s \\
 W_0 - J S & = - 0.10 & W_0 - R C & = - 0.19 & &
 \end{array}$$

Comparisons of Clock Errors at 0^h Sidereal referred to the Standard Observer, from Observations with the Transit-Circle with Central and with Annular Illumination.

Date.	Observer.	Clock Error.		Central. — Annular.	Date.	Observer.	Clock Error.		Central. — Annular.
		Central Illumina- tion.	Annular Illumina- tion.				Central Illumina- tion.	Annular Illumina- tion.	
1908.		s	s	s	1908.		s	s	s
Apr. 30	E	- 4'95	- 4'92	- 0'03	Sept. 25	E	- 31'24	- 31'25	+ 0'01
May 1	R C	- 5'49	- 5'46	- 0'03	26	R C	- 31'54	- 31'49	- 0'05
2	W	- 6'29	- 6'11	- 0'18	29	E	- 32'83	- 32'75	- 0'08
4	E	- 7'23	- 7'25	+ 0'02	30	R C	- 32'83	- 32'78	- 0'05
6	W	- 8'31	- 8'34	+ 0'03	Oct. 1	W	- 33'00	- 32'99	- 0'01
9	R C	- 9'96	- 9'90	- 0'06	2	S D	- 33'46	- 33'50	+ 0'04
13	R C	- 12'20	- 12'18	- 0'02	4	R C	- 33'52	- 33'53	+ 0'01
18	W	- 15'31	- 15'32	+ 0'01	5	W	- 34'24	- 34'17	- 0'07
21	S	- 17'04	- 16'83	- 0'21	6	G C	- 34'65	- 34'58	- 0'07
23	G C	+ 41'74	+ 41'89	- 0'15	8	R C	- 34'73	- 34'74	+ 0'01
27	G	+ 39'13	+ 39'14	- 0'01	10	H A	- 35'16	- 35'19	+ 0'03
June 2	R C	+ 35'41	+ 35'52	- 0'11	15	S E	- 36'74	- 36'73	- 0'01
4	S E	+ 34'01	+ 34'02	- 0'01	15	R C	- 36'56	- 36'58	+ 0'02
6	R C	+ 32'70	+ 32'72	- 0'02	21	W	- 38'47	- 38'43	- 0'04
10	R C	+ 29'86	+ 29'96	- 0'10	23	E	- 38'89	- 38'90	+ 0'01
25	E	+ 18'18	+ 18'23	- 0'05	26	S	- 39'91	- 39'79	- 0'12
30	R C	+ 14'18	+ 14'21	- 0'03	29	W	- 40'65	- 40'67	+ 0'02
July 1	G	+ 13'44	+ 13'52	- 0'08	Nov. 7	S D	- 44'10	- 43'98	- 0'12
23	S E	- 5'25	- 5'23	- 0'02	9	R C	- 44'75	- 44'73	- 0'02
25	W	- 6'88	- 6'81	- 0'07	17	R C	- 47'49	- 47'47	- 0'02
29	W	- 10'14	- 10'03	- 0'11	23	S	- 49'39	- 49'28	- 0'11
31	S	- 11'91	- 11'86	- 0'05	25	R C	- 49'70	- 49'71	+ 0'01
Aug. 1	H A	- 12'60	- 12'49	- 0'11	26	W	- 50'06	- 50'05	- 0'01
7	W	- 18'31	- 18'30	- 0'01	27	H A	- 50'30	- 50'31	+ 0'01
11	W	- 19'04	- 18'98	- 0'06	28	E	- 50'65	- 50'57	- 0'08
14	W	- 19'73	- 19'70	- 0'03	Dec. 7	R C	- 53'01	- 53'04	+ 0'03
Sept. 6	R C	- 26'24	- 26'15	- 0'09	10	R C	- 54'05	- 54'07	+ 0'02
8	W	- 27'25	- 27'17	- 0'08	11	H A	- 54'61	- 54'59	- 0'02
17	S E	- 29'45	- 29'42	- 0'03	14	R C	- 55'34	- 55'34	0'00
					17	E	- 56'25	- 56'35	+ 0'10

The annular illumination was in use to 1908 April 29. The central illumination with simple condensing lens was in use from April 30 to September 16, and with compound corrected lens from September 17 to end of year.

As the Sun, Mercury and Venus are observed without and the clock stars generally with artificial light, the following corrections have been applied to the observed Right Ascensions of the Sun, Mercury and Venus in the ledgers for the year 1908 to correct to the system of clock error given by the annular illumination :—

Date.	Correction.
January 1 to April 29.	8 ^s .00
April 30 to September 16.	+ 8 ^s .06
September 17 to end of year.	+ 8 ^s .02

TRANSIT-CIRCLE TABLES.

xciii

Errors of Division.

In use from 1897.

The following table contains the adopted result for errors of division for every degree. The errors are always considered in the nature of reading too much. $r = 61''\cdot 4$ nearly.

Degree.	Excess of Reading.	Degree.	Excess of Reading.	Degree.	Excess of Reading.	Degree.	Excess of Reading.	Degree.	Excess of Reading.	Degree.	Excess of Reading.	Degree.	Excess of Reading.	Degree.	Excess of Reading.	Degree.	Excess of Reading.
0	0 ^r 051	36	0 ^r 054	72	0 ^r 053	108	0 ^r 059	144	0 ^r 087	180	0 ^r 067	216	0 ^r 064	252	0 ^r 075	288	0 ^r 073
1	049	37	050	73	049	109	065	145	084	181	069	217	064	253	078	289	078
2	047	38	047	74	048	110	059	146	079	182	069	218	064	254	082	290	077
3	051	39	050	75	053	111	056	147	089	183	067	219	064	255	085	291	080
4	051	40	051	76	056	112	059	148	089	184	067	220	063	256	089	292	081
5	049	41	051	77	052	113	060	149	089	185	061	221	062	257	088	293	079
6	053	42	056	78	053	114	067	150	084	186	056	222	063	258	090	294	088
7	052	43	055	79	054	115	066	151	083	187	059	223	064	259	089	295	092
8	055	44	049	80	060	116	071	152	086	188	063	224	063	260	096	296	087
9	056	45	045	81	063	117	071	153	090	189	063	225	062	261	103	297	085
10	052	46	050	82	067	118	070	154	096	190	053	226	065	262	099	298	081
11	052	47	057	83	068	119	074	155	100	191	049	227	071	263	106	299	085
12	053	48	060	84	070	120	082	156	096	192	056	228	069	264	108	300	096
13	052	49	058	85	069	121	088	157	082	193	061	229	071	265	105	301	097
14	057	50	052	86	070	122	090	158	075	194	058	230	071	266	107	302	093
15	047	51	050	87	073	123	095	159	078	195	057	231	071	267	108	303	096
16	046	52	055	88	072	124	099	160	080	196	058	232	078	268	104	304	101
17	044	53	065	89	069	125	092	161	078	197	054	233	082	269	099	305	097
18	050	54	065	90	068	126	095	162	075	198	057	234	081	270	098	306	102
19	050	55	071	91	067	127	095	163	075	199	064	235	080	271	094	307	102
20	052	56	071	92	072	128	093	164	082	200	060	236	076	272	085	308	099
21	043	57	069	93	069	129	091	165	088	201	061	237	080	273	085	309	093
22	043	58	072	94	071	130	088	166	088	202	062	238	085	274	083	310	092
23	046	59	071	95	069	131	094	167	079	203	064	239	093	275	092	311	099
24	044	60	067	96	067	132	089	168	070	204	063	240	091	276	090	312	098
25	045	61	063	97	064	133	085	169	063	205	063	241	089	277	095	313	097
26	050	62	061	98	071	134	085	170	062	206	062	242	084	278	105	314	097
27	048	63	060	99	077	135	084	171	065	207	063	243	093	279	098	315	090
28	045	64	065	100	074	136	087	172	058	208	061	244	095	280	092	316	090
29	049	65	061	101	071	137	092	173	055	209	063	245	092	281	090	317	090
30	059	66	059	102	071	138	087	174	053	210	072	246	089	282	093	318	091
31	064	67	054	103	070	139	082	175	056	211	075	247	082	283	095	319	083
32	061	68	052	104	069	140	081	176	060	212	070	248	071	284	092	320	087
33	063	69	057	105	062	141	082	177	067	213	067	249	075	285	086	321	079
34	059	70	058	106	063	142	085	178	071	214	063	250	082	286	084	322	078
35	052	71	056	107	058	143	090	179	067	215	060	251	078	287	076	323	087

Division-Correction.

Applicable to Observations made with the six Microscopes.

(Always Additive.)

In use from 1897.

The corrections recur after every 60° of pointer reading.

Pointer Reading.	Corr.	Pointer Reading.	Corr.	Pointer Reading.	Corr.	Pointer Reading.	Corr.	Pointer Reading.	Corr.	Pointer Reading.	Corr.
0° 1'	0".25	10°-11'	0".56	20°-21'	1".06	30°-31'	0".31	40°-41'	0".57	50°-51'	0".50
1- 2	0".28	11-12	0".54	21-22	0".99	31-32	0".32	41-42	0".59	51-52	0".43
2- 3	0".26	12-13	0".57	22-23	0".91	32-33	0".22	42-43	0".59	52-53	0".31
3- 4	0".24	13-14	0".65	23-24	0".82	33-34	0".16	43-44	0".60	53-54	0".29
4- 5	0".28	14-15	0".72	24-25	0".73	34-35	0".31	44-45	0".67	54-55	0".35
5- 6	0".35	15-16	0".73	25-26	0".64	35-36	0".32	45-46	0".61	55-56	0".32
6- 7	0".46	16-17	0".80	26-27	0".56	36-37	0".39	46-47	0".60	56-57	0".29
7- 8	0".49	17-18	0".88	27-28	0".50	37-38	0".47	47-48	0".59	57-58	0".29
8- 9	0".47	18-19	0".95	28-29	0".42	38-39	0".51	48-49	0".56	58-59	0".28
9-10	0".48	19-20	1".10	29-30	0".29	39-40	0".58	49-50	0".47	59-60	0".22

When the Pointer reads 0°.30' the division under Microscope A is 271°; under Microscope C, 211°; under Microscope E, 151°; under Microscope B, 91°; under Microscope D, 31°; and under Microscope F, 331°.

The above table of division-errors is a slight modification of that published and used in the second *Ten Year Catalogue*, 1890, owing to the incorporation at a later date of the results of a special examination of the single degree divisions for five degrees on either side of the pole corresponding to pointer readings 14°-24°.

The corrections for division-errors with the transit-circle have been changed four times since its erection; from 1851 to 1856 the values were derived from the observations made in 1851; from 1857 to 1867 the values were derived from the observations made in 1856; from 1868 to 1879 the values were derived by taking the mean of the observations made in 1851 and 1856; from 1880 to 1896 the last values were still used, but in order to allow for the distance between Microscope A and the pointer being 89°.30' instead of 90°, as it had till then been assumed to be, what had been taken as the correction for 1°—that was, for all pointer readings between 0°.30' and 1°.30'—was considered to be the correction for all pointer readings between 0° and 1°, and so on. An account of the determination of the errors is given in the *Introduction* for 1897, and also tables of the differences of the values now used (resulting from the determinations in 1851, 1856, and 1898) and the values adopted in previous years.

TRANSIT-CIRCLE TABLES.

xcv

Table of Corrections for 5' Divisions.

FOR NADIR.

Pointer Reading ...	179°.30'	35'	40'	45'	50'	55'	180°.0'	5'	10'	15'	20'	25'	30'	No. of Sets.
Division-Correction	0".22	0".30	0".35	0".38	0".40	0".45	0".45	0".41	0".38	0".32	0".29	0".26	0".25	10

FOR COLLIMATORS.

Pointer Reading {	89°.30' or 269°.30'	35'	40'	45'	50'	55'	90°.0' or 270°.0'	5'	10'	15'	20'	25'	30'	No. of Sets.
Division-Correction	0".29	0".28	0".38	0".40	0".41	0".43	0".46	0".50	0".43	0".34	0".26	0".19	0".24	10

FOR POLAR STARS.

Pointer Reading ...	317°.30'	35'	40'	45'	50'	55'	318°.0'	5'	10'	15'	20'	25'	30'	No. of Sets.
Division-Correction	0".88	0".98	1".03	1".04	1".07	1".12	1".14	1".11	1".03	0".94	0".91	0".92	0".95	9

Pointer Reading ...	319°.30'	35'	40'	45'	50'	55'	320°.0'	5'	10'	15'	20'	25'	30'	No. of Sets.
Division-Correction	1".10	1".09	1".14	1".13	1".09	1".05	1".03	1".06	1".09	1".11	1".15	1".15	1".06	14

Pointer Reading ...	321°.30'	35'	40'	45'	50'	55'	322°.0'	5'	10'	15'	20'	25'	30'	No. of Sets.
Division-Correction	0".99	1".06	1".02	0".99	1".01	1".02	1".02	1".03	1".02	0".99	0".94	0".92	0".91	18

Pointer Reading ...	322°.30'	35'	40'	45'	50'	55'	323°.0'	5'	10'	15'	20'	25'	30'	No. of Sets.
Division-Correction	0".91	0".98	0".97	0".90	0".87	0".92	0".98	1".00	1".02	0".99	0".92	0".86	0".82	4

Pointer Reading ...	323°.30'	35'	40'	45'	50'	55'	324°.0'	5'	10'	15'	20'	25'	30'	No. of Sets.
Division-Correction	0".82	0".74	0".77	0".82	0".82	0".81	0".83	0".83	0".77	0".73	0".73	0".76	0".73	9

Personality in the reading of the Microscope-Micrometers of the Transit-Circle.

As an observation of the Sun requires an extra observer to read the microscope-micrometers, any personality in the bisection of the divisions of the circle will introduce a systematic error into the observed declinations.

To determine the amount of this error, readings of the microscopes for the Nadir observation made near to the observation of the Sun have been taken by the microscope-reader as well as by the regular transit-circle observer.

The difference between the means of these microscope readings has been tabulated (Transit-circle observer—Microscope-reader) and a comparison made between each regular observer and each microscope-reader. It is found that, of the four microscope-readers, two exhibit no particular personality, and the other two a well-marked personality in opposite directions.

It was arranged that (from April 15, 1909) the two microscope-readers with marked personality should cease to assist in the observation of the Sun.

T.C. Observer—S D.

Date.	T.C. Observer.	Mean of Microscopes.		Difference.	Date.	T.C. Observer.	Mean of Microscopes.		Difference.		
		T.C. Observer.	S D.				T.C. Observer.	S D.			
1907.					1908.						
Nov.	5	J S	27 ^h 24	27 ^h 65	— 0 ^h 41	May	27	G	18 ^h 02	17 ^h 72	+ 0 ^h 30
	6	E	45 ^h 78	45 ^h 60	+ 0 ^h 18		29	R C	26 ^h 87	26 ^h 83	+ 0 ^h 04
	9	J S	31 ^h 82	31 ^h 90	— 0 ^h 08						
Dec.	2	W	30 ^h 81	30 ^h 52	+ 0 ^h 29	June	22	G	14 ^h 08	14 ^h 59	— 0 ^h 51
	5	R C	7 ^h 00	7 ^h 13	— 0 ^h 13		23	W	43 ^h 73	43 ^h 98	— 0 ^h 25
	6	W	31 ^h 52	31 ^h 36	+ 0 ^h 16		24	S	60 ^h 16	59 ^h 61	+ 0 ^h 55
	7	J S	27 ^h 44	27 ^h 57	— 0 ^h 13		25	E	39 ^h 59	39 ^h 93	— 0 ^h 34
1908.						July	23	S E	8 ^h 00	8 ^h 02	— 0 ^h 02
Jan.	3	W	34 ^h 16	33 ^h 50	+ 0 ^h 66		24	S	34 ^h 97	34 ^h 18	+ 0 ^h 79
	4	R C	25 ^h 94	26 ^h 34	— 0 ^h 40						
	20	J S	29 ^h 51	29 ^h 43	+ 0 ^h 08	Aug.	8	H A	28 ^h 60	28 ^h 68	— 0 ^h 08
	23	J S	34 ^h 16	34 ^h 55	— 0 ^h 39						
Feb.	12	W	19 ^h 41	18 ^h 61	+ 0 ^h 80	Sept.	8	W	33 ^h 35	33 ^h 49	— 0 ^h 14
	14	E	41 ^h 47	41 ^h 63	— 0 ^h 16		10	R C	25 ^h 14	25 ^h 64	— 0 ^h 50
							11	W	31 ^h 27	31 ^h 29	— 0 ^h 02
							12	H A	23 ^h 63	24 ^h 09	— 0 ^h 46
Mar.	2	R C	7 ^h 11	7 ^h 44	— 0 ^h 33						
	3	W	34 ^h 01	34 ^h 42	— 0 ^h 41	Oct.	6	G C	27 ^h 94	28 ^h 39	— 0 ^h 45
	5	E	22 ^h 37	22 ^h 19	+ 0 ^h 18		7	E	36 ^h 99	37 ^h 31	— 0 ^h 32
	31	W	28 ^h 52	28 ^h 61	— 0 ^h 09						
May	7	J S	21 ^h 35	22 ^h 00	— 0 ^h 65	Nov.	2	W	29 ^h 31	29 ^h 22	+ 0 ^h 09
	9	R C	5 ^h 45	5 ^h 51	— 0 ^h 06		4	H A	27 ^h 11	27 ^h 17	— 0 ^h 06
							25	R C	28 ^h 31	28 ^h 30	+ 0 ^h 01
							26	W	33 ^h 83	33 ^h 50	+ 0 ^h 33

TRANSIT-CIRCLE TABLES.

xcvii

T.C. Observer—S.

Date.	T.C. Observer.	Mean of Microscopes.		Difference.	Date.	T.C. Observer.	Mean of Microscopes.		Difference.
		T.C. Observer.	S.				T.C. Observer.	S.	
1907.					1908.				
Nov. 11	R C	29'17	29'44	— 0'27	June 1	E	39'19	39'42	— 0'23
12	W	12'08	11'89	+ 0'19	3	W	38'54	38'41	+ 0'13
16	W	2'00	2'23	— 0'23	5	E	51'38	51'85	— 0'47
					6	R C	27'74	28'25	— 0'51
					29	E	35'98	36'03	— 0'05
					30	R C	25'78	26'45	— 0'67
Dec. 9	R C	9'06	9'43	— 0'37					
10	W	34'05	34'53	— 0'48	July 1	G	23'24	23'60	— 0'36
12	E	36'37	35'76	+ 0'61	2	S D	19'96	20'31	— 0'35
13	R C	5'69	6'42	— 0'73	3	E	35'56	36'19	— 0'63
					22	S D	30'79	31'04	— 0'25
1908.					27	G C	35'02	35'73	— 0'71
Jan. 10	R C	16'25	16'68	— 0'43	29	W	33'82	33'78	+ 0'04
11	W	58'26	57'78	+ 0'48					
28	J S	34'89	35'78	— 0'89	Aug. 1	H A	11'00	10'83	+ 0'17
29	E	17'30	17'06	+ 0'24	17	G C	9'45	10'37	— 0'92
30	R C	26'94	27'71	— 0'77	21	R C	2'72	3'19	— 0'47
31	W	34'75	35'39	— 0'64	22	G C	31'67	32'81	— 1'14
Feb. 1	J S	31'68	32'20	— 0'52	Sept. 1	E	37'91	38'83	— 0'92
16	W	34'06	33'87	+ 0'19	2	R C	31'39	31'38	+ 0'01
18	E	38'95	39'09	— 0'14	5	E	39'46	39'48	— 0'02
					30	R C	13'77	14'33	— 0'56
Mar. 19	W	34'11	34'52	— 0'41					
20	H A	11'16	11'76	— 0'60	Oct. 1	W	15'74	16'34	— 0'60
					29	W	33'58	34'20	— 0'62
Apr. 7	G	40'90	41'06	— 0'16					
8	S E	48'94	49'51	— 0'57	Nov. 10	G C	16'34	17'54	— 1'20
9	R C	34'27	34'60	— 0'33					
May 1	R C	7'14	7'95	— 0'81	Dec. 7	R C	19'53	19'88	— 0'35
23	G C	10'54	11'88	— 1'34	8	W	25'65	26'20	— 0'55

GREENWICH OBSERVATIONS, 1908.

n

T.C. Observer—G C.

Date.	T.C. Observer.	Mean of Microscopes.		Difference.	Date.	T.C. Observer.	Mean of Microscopes.		Difference.
		T.C. Observer.	G C.				T.C. Observer.	G C.	
1907.					1908.				
Nov. 21	H A	24'89	24'62	+ 0'27	July 17	S D	30'21	29'61	+ 0'60
Dec. 16	E	37'15	37'14	+ 0'01	Aug. 11	W	42'40	42'46	- 0'06
17	R C	6'00	5'50	+ 0'50	12	E	21'07	20'58	+ 0'49
20	E	45'20	44'20	+ 1'00	14	W	32'46	32'29	+ 0'17
1908.					15	H A	25'09	24'94	+ 0'15
Mar. 9	S D	22'31	21'63	+ 0'68	Sept. 17	S E	36'49	35'91	+ 0'58
10	R C	16'14	15'84	+ 0'30	Oct. 12	R C	32'01	31'77	+ 0'24
12	W	22'93	22'31	+ 0'62	15	S E	57'53	56'89	+ 0'64
14	R C	36'29	35'84	+ 0'45	16	S	22'28	21'59	+ 0'69
May 18	W	23'51	22'91	+ 0'60	30	S	32'96	31'86	+ 1'10
21	S	19'39	18'61	+ 0'78	Nov. 9	R C	26'03	25'74	+ 0'29
22	H A	17'03	16'58	+ 0'45	13	S D	18'00	16'48	+ 1'52
June 14	R C	59'11	58'46	+ 0'65	Dec. 22	E	29'29	28'65	+ 0'64
15	H A	[25'26]	23'56	[+ 1'70]					
18	S D	59'70	59'39	+ 0'31					

T.C. Observer—H A.

Date.	T.C. Observer.	Mean of Microscopes.		Difference.	Date.	T.C. Observer.	Mean of Microscopes.		Difference.
		T.C. Observer.	H A.				T.C. Observer.	H A.	
1907.					1908.				
Nov. 1	J S	24'31	24'32	- 0'01	May 16	E	41'83	41'27	+ 0'56
28	S E	14'36	14'56	- 0'20	30	W	33'37	33'13	+ 0'24
29	S D	31'55	31'06	+ 0'49	June 11	W	34'84	34'66	+ 0'18
30	E	38'71	38'55	+ 0'16	13	E	41'47	41'43	+ 0'04
1908.					Aug. 4	S D	34'87	35'23	- 0'36
Jan. 13	E	30'08	29'68	+ 0'40	7	W	18'61	18'19	+ 0'42
15	W	35'55	35'38	+ 0'17	25	R C	27'58	27'72	- 0'14
18	R C	9'89	10'01	- 0'12	29	R C	27'97	27'98	- 0'01
Feb. 5	J S	38'57	38'88	- 0'31	Sept. 23	W	22'47	22'42	+ 0'05
8	W	28'96	29'02	- 0'06	26	R C	25'77	25'83	- 0'06
26	E	35'82	35'72	+ 0'10	Oct. 21	W	12'69	12'39	+ 0'30
29	J S	36'38	36'14	+ 0'24	Nov. 7	S D	14'06	14'33	- 0'27
Mar. 24	S E	40'85	40'52	+ 0'33	20	E	30'15	30'59	- 0'44
27	W	22'17	22'35	- 0'18	21	R C	23'29	23'31	- 0'02
Apr. 22	S	19'36	19'02	+ 0'34	Dec. 28	W	20'07	20'10	- 0'03
24	S E	50'76	50'91	- 0'15	30	G C	33'94	34'86	- 0'92
May 12	S E	12'04	12'20	- 0'16					

TRANSIT-CIRCLE TABLES.

xcix

*Mean Differences of Microscope Readings between the Regular Observers
and the Readers of the Microscopes for the Sun.*

	S D.	S.	G C.	H A.
W.	+ '13 ₁₁	- '21 ₁₂	+ '33 ₄	+ '12 ₉
J S.	- '26 ₆	- '71 ₂	...	- '03 ₃
E.	- '09 ₅	- '18 ₉	+ '56 ₄	+ '14 ₆
R C.	- '20 ₇	- '47 ₁₃	+ '41 ₆	- '07 ₅
Mean	- '07 ₂₉	- '32 ₃₆	+ '43 ₁₄	+ '06 ₂₃

Observation to determine the Astronomical Flexure of the Transit-Circle, 1908.

Date.	Observer.	Reader of Micro- scopes.	Mean Concluded Circle-Reading.		Resulting Correction for Astronomical Horizontal Flexure.	Temperature.
			North Collimator.	South Collimator.		
1908. d h August 26. o	W	W	269° 37' 25".77	89° 37' 26".23	- 0".23	...

This observation gives the value of $-0''.23$ for the Astronomical flexure.

The observations for coincidence of north and south collimators were in every case made through the openings in the central cube of the transit-circle.

These observations were made by means of contacts above and below the single horizontal wires of the collimators.

No correction for flexure has been applied since April 1879.

*Excess of Reflexion Results above Direct Results, for Observations of
Zenith-Distances with the Transit-Circle, 1908.*

In this table the reflexion observations are not corrected for inclination of the vertical at the surface
of the mercury to that at the centre of the instrument.

Star's Name.	Approx. R.A.	Approx. N.P.D.	R—D.	No. of Obs. R. and D.	Weight	Star's Name.	Approx. R.A.	Approx. N.P.D.	R—D.	No. of Obs. R. and D.	Weight
<i>θ</i> Cephei S.P.	^{h m} 20. 28	— 27. 3	— 0.83	1	1	Bradley 344 S.P.	^{h m} 2. 35	— 8. 9	— 0.15	1	1
<i>ε</i> Cassiopeia S.P.	1. 48	— 26. 8	— 1.10	1	1	Piazzi IX. 37 S.P.	9. 24	— 8. 3	— 1.01	3	3
76 Ursæ Majoris S.P.	12. 38	— 26. 8	— 1.30	1	1	76 Draconis S.P.	20. 49	— 7. 8	— 0.47	1	1
<i>ι</i> Cephei S.P.	22. 46	— 24. 3	— 0.80	2	2	Piazzi X. 22 S.P.	10. 20	— 7. 0	+ 0.03	1	1
<i>ζ</i> Draconis S.P.	17. 9	— 24. 2	— 1.10	1	1	Bradley 3058 S.P.	22. 55	— 6. 2	— 0.29	1	1
3 Draconis S.P.	11. 37	— 22. 7	— 1.62	1	1	Bradley 1731 S.P.	12. 48	— 6. 1	— 0.18	3	3
<i>σ</i> ² Ursæ Majoris S.P.	9. 2	— 22. 5	— 1.64	1	1	Bradley 1399 S.P.	10. 17	— 5. 3	+ 0.01	2	2
Groombridge 2214 S.P.	15. 14	— 22. 3	— 1.76	1	1	Groombridge 944 S.P.	5. 33	— 4. 9	+ 0.22	1	1
<i>ρ</i> Ursæ Majoris S.P.	8. 54	— 22. 0	— 0.62	1	1	Groombridge 750 S.P.	4. 8	— 4. 7	— 1.06	1	1
<i>γ</i> Draconis S.P.	17. 32	— 21. 8	+ 0.39	1	1	2 Ursæ Minoris S.P.	0. 56	— 4. 2	+ 1.47	2	2
Piazzi VII. 67 S.P.	7. 22	— 21. 3	— 0.10	1	1	8 Ursæ Minoris S.P.	18. 1	— 3. 4	+ 0.26	2	2
<i>ω</i> Draconis S.P.	17. 38	— 21. 2	— 1.72	1	1	Groombridge 3548 S.P.	21. 18	— 3. 4	+ 0.19	1	1
Groombridge 1564 S.P.	9. 35	— 20. 3	— 1.33	1	1	Bradley 3147 S.P.	23. 28	— 3. 2	— 0.19	1	1
<i>ε</i> Draconis S.P.	19. 48	— 20. 0	— 1.41	2	2	Cephei 51 S.P.	6. 59	— 2. 8	— 0.27	2	2
11 Cephei S.P.	21. 41	— 19. 1	— 1.92	1	1	Groombridge 2283 S.P.	15. 6	— 2. 4	+ 0.41	3	3
24 Cephei S.P.	22. 8	— 18. 1	+ 0.10	1	1	Bradley 65 S.P.	0. 58	— 1. 5	+ 0.65	1	1
Bradley 2673 S.P.	20. 30	— 17. 8	— 0.13	1	1	Polaris S.P.	1. 27	— 1. 2	— 0.43	3	3
11 Ursæ Minoris S.P.	15. 17	— 17. 8	— 0.85	1	1	Cephei 24 S.P.	19. 50	— 1. 2	— 0.26	1	1
Bradley 348 S.P.	2. 29	— 17. 6	— 1.09	1	1	Groombridge 1119 S.P.	8. 9	— 1. 1	— 0.45	3	3
16 Cephei S.P.	21. 58	— 17. 3	— 0.28	1	1	λ Ursæ Minoris.	19. 11	— 1. 0	— 0.61	5	4
31 Cephei S.P.	22. 34	— 16. 8	— 0.96	2	2	Cephei 24.	19. 50	— 1. 1	+ 0.58	3	3
Piazzi IX. 187 S.P.	9. 50	— 16. 7	— 0.34	3	3	Polaris.	1. 27	— 1. 2	+ 0.30	2	2
Groombridge 1374 S.P.	7. 49	— 15. 8	+ 2.86	1	1	Bradley 1672.	12. 14	— 1. 8	— 0.41	3	3
Bradley 1147 S.P.	8. 8	— 14. 0	— 1.65	1	1	Groombridge 2283 ...	15. 6	— 2. 4	— 0.08	4	3
5 Ursæ Minoris S.P.	14. 28	— 13. 9	+ 0.85	1	1	Groombridge 1004.	6. 12	— 3. 2	+ 0.48	2	2
Bradley 1146 S.P.	10. 27	— 13. 8	+ 0.29	1	1	Bradley 3147.	23. 28	— 3. 2	— 0.16	2	2
Piazzi VI. 201 S.P.	6. 47	— 12. 9	— 0.99	1	1	Groombridge 3548 ...	21. 18	— 3. 3	— 0.62	5	4
Bradley 431 S.P.	3. 9	— 12. 6	— 0.24	1	1	8 Ursæ Minoris.	18. 1	— 3. 4	+ 0.08	6	4
<i>θ</i> Ursæ Minoris S.P.	15. 34	— 12. 3	— 0.85	3	3	Groombridge 642.	3. 37	— 3. 6	+ 0.60	2	2
77 Draconis S.P.	21. 7	— 12. 2	— 1.08	2	2	2 Ursæ Minoris.	0. 56	— 4. 2	— 0.70	1	1
4 Ursæ Minoris S.P.	14. 9	— 12. 0	— 1.29	1	1	Bradley 1730.	12. 48	— 6. 1	+ 0.07	1	1
<i>ζ</i> Ursæ Minoris S.P.	15. 47	— 11. 9	+ 1.17	1	1	Bradley 1731.	12. 48	— 6. 1	— 0.28	2	2
Bradley 392 S.P.	2. 54	— 11. 0	— 0.73	1	1	76 Draconis.	20. 49	— 7. 8	+ 0.81	1	1
Bradley 117 S.P.	1. 4	— 10. 8	— 1.13	1	1	Piazzi IX., 37.	9. 24	— 8. 3	— 0.95	1	1
Piazzi VII. 75 S.P.	6. 31	— 10. 3	— 0.60	1	1	40 Draconis.	18. 6	— 10. 0	— 1.40	1	1

TRANSIT-CIRCLE TABLES.

ci

*Excess of Reflexion Results above Direct Results, for Observations of
Zenith-Distances with the Transit-Circle, 1908—continued.*

Star's Name.	Approx. R.A.	Approx. N.P.D.	R—D.	No. of Obs. R. and D.	Weight.	Star's Name.	Approx. R.A.	Approx. N.P.D.	R—D.	No. of Obs. R. and D.	Weight.
	h m	°	"				h m	°	"		
Bradley 1508.....	10. 53	11. 7	-0°53	2	2	1 Geminorum.....	5. 59	66. 7	+0°90	1	1
77 Draconis.....	21. 7	12. 2	+0°45	2	2	23 Comæ.....	12. 30	66. 8	+0°16	2	2
θ Ursæ Minoris.....	15. 34	12. 3	-0°09	3	3	α Arietis.....	2. 2	66. 9	+0°70	1	1
Piazzi XVI., 195.....	16. 35	12. 3	-0°60	1	1	A ¹ Tauri.....	3. 59	68. 1	+0°19	1	1
κ Cephei.....	20. 12	12. 5	+1°11	1	1	β Herculis.....	16. 26	68. 3	+0°99	2	2
35 Draconis.....	17. 54	13. 0	+0°09	2	2	20 Comæ.....	12. 25	68. 6	+1°67	1	1
Bradley 6.....	0. 11	13. 5	+0°24	1	1	28 Pegasi.....	22. 6	69. 4	+0°92	1	1
Bradley 1147.....	8. 8	13. 9	-0°08	1	1	110 Herculis.....	18. 42	69. 5	-1°76	1	1
5 Ursæ Minoris.....	14. 28	13. 9	-0°93	2	2	ν Geminorum.....	6. 23	69. 7	-0°63	1	1
19 Ursæ Minoris.....	16. 13	13. 9	-0°15	1	1	γ ¹ Leonis.....	10. 15	69. 7	-0°28	1	1
Groombridge 3813...	22. 27	14. 2	-0°24	1	1	ι Serpentis.....	15. 38	70. 0	+1°15	3	3
						5 Vulpeculæ.....	19. 22	70. 1	+0°69	1	1
β Ursæ Minoris.....	14. 51	15. 5	-1°22	1	1						
79 Draconis.....	21. 52	16. 7	-0°58	1	1	ξ Boötis.....	14. 47	70. 5	+0°61	1	1
16 Cephei.....	21. 58	17. 2	-2°11	2	2	1 Pegasi.....	21. 18	70. 6	0°00	1	1
χ Draconis.....	18. 23	17. 3	+1°45	2	2	γ Sagittæ.....	19. 55	70. 7	+1°27	1	1
40 Cassiopeiæ.....	1. 31	17. 4	-0°71	1	1	15 Arietis.....	2. 6	70. 9	-0°92	1	1
11 Ursæ Minoris.....	15. 17	17. 8	-0°21	2	2	φ Pegasi.....	23. 48	71. 4	+0°27	1	1
γ Ursæ Minoris.....	15. 21	17. 8	+0°25	1	1	Mayer 136.....	3. 48	72. 9	-0°13	2	2
50 Cassiopeiæ.....	1. 56	18. 0	+0°12	2	2	π ¹ Boötis.....	14. 36	73. 2	-0°14	2	2
						β ¹ Sagittæ.....	19. 37	73. 7	+0°14	1	1
ζ Cygni.....	21. 9	60. 1	-0°42	1	1	ε Sagittæ.....	19. 33	73. 7	+0°81	2	2
14 Pegasi.....	21. 46	60. 2	+0°83	3	3	τ ¹ Serpentis.....	15. 22	74. 2	-0°12	2	2
η Pegasi.....	22. 39	60. 2	-0°33	1	1	γ ² Delphini.....	20. 42	74. 2	+0°65	1	1
α Trianguli.....	1. 48	60. 8	+0°68	1	1						
ψ Tauri.....	4. 1	61. 2	+0°82	1	1	β Leonis.....	11. 44	74. 9	+0°60	1	1
B. D. + 28° 1659....	8. 47	61. 4	+1°14	1	1	Piazzi XIV. 221.....	14. 52	75. 2	+0°41	1	1
31 Comæ.....	12. 47	61. 9	-0°45	1	1	δ Delphini.....	20. 39	75. 2	-0°03	1	1
μ Herculis.....	17. 43	62. 2	+0°25	2	2	α Pegasi.....	23. 0	75. 3	+1°22	1	1
						70 Virginis.....	13. 24	75. 7	-0°13	2	2
ε ² Boötis.....	14. 41	62. 5	+0°72	1	1	ω Herculis.....	16. 21	75. 7	+0°57	1	1
ψ Boötis.....	15. 1	62. 7	+1°63	1	1	α Ophiuchi.....	17. 31	77. 3	+1°03	3	3
α Coronæ.....	15. 31	62. 9	+0°77	4	3	f Tauri.....	3. 26	77. 3	+0°60	1	1
34 Boötis.....	14. 38	63. 1	+1°17	1	1	λ Tauri.....	3. 55	77. 7	+0°91	1	1
ν Piscium.....	1. 14	63. 2	+0°29	1	1	α Cancri.....	8. 54	77. 8	+0°99	2	2
Coronæ.....	15. 38	63. 4	+0°20	1	1	β Aquilæ.....	19. 21	78. 2	-1°74	1	1
89 Herculis.....	17. 52	63. 9	-0°14	1	1	π Piscium.....	1. 32	78. 3	+0°45	1	1
10 Vulpeculæ.....	19. 40	64. 4	+0°36	1	1						
ι Pegasi.....	22. 3	65. 1	+1°19	2	2	ι Leonis.....	10. 45	78. 9	-0°01	1	1
ν Cancri.....	8. 57	65. 2	+1°90	1	1	s Tauri.....	3. 25	78. 9	-0°98	1	1
κ Geminorum.....	7. 39	65. 3	-0°76	1	1	ζ Pegasi.....	22. 37	79. 6	-0°51	1	1
α Vulpeculæ.....	19. 25	65. 5	-0°01	1	1	γ Aquilæ.....	19. 41	79. 6	+0°28	2	2
17 Tauri.....	3. 40	66. 1	-0°26	1	1	85 Ceti.....	2. 38	79. 6	+0°59	1	1

*Excess of Reflexion Results above Direct Results, for Observations of
Zenith-Distances with the Transit-Circle, 1908—continued.*

Star's Name.	Approx. R.A.	Approx. N.P.D.	R—D.	No. of Obs. R. and D.	Weight.	Star's Name.	Approx. R.A.	Approx. N.P.D.	R—D.	No. of Obs. R. and D.	Weight.
β Cancri.....	h m 8. 12	80. 5	+0.82	1	1	κ Piscium.....	h m 23. 22	89. 2	+1.09	1	1
ϵ Pegasi.....	21. 40	80. 5	+1.19	4	3	10 Tauri.....	3. 32	89. 8	+2.71	1	1
ω Virginis.....	11. 34	81. 3	+0.37	1	1	η Aquarii.....	22. 31	90. 5	+0.60	1	1
α Aquilæ.....	19. 46	81. 3	+0.48	2	2	θ Aquilæ.....	20. 7	91. 0	-0.16	2	2
σ Piscium.....	1. 40	81. 3	-0.04	1	1	20 Ceti.....	0. 48	91. 6	+1.14	1	1
13 Monocerotis.....	6. 28	82. 5	+0.07	1	1	29 Sextantis.....	10. 25	92. 2	+1.76	1	1
κ Herculis.....	16. 45	82. 6	+0.59	1	1	η Serpentis.....	18. 17	92. 9	+0.24	1	1
10 Leonis.....	9. 32	82. 7	-0.41	1	1	14 Hydræ.....	8. 45	93. 1	+0.83	1	1
μ Aquilæ.....	19. 30	82. 8	+1.46	1	1	δ Ophiuchi.....	16. 10	93. 4	+1.51	1	1
ζ^1 Piscium.....	1. 9	82. 9	+2.88	1	1	6 Sextantis.....	9. 47	93. 8	+1.05	1	1
37 Sextantis.....	10. 41	83. 1	+0.95	2	2	25 Monocerotis.....	7. 33	93. 9	+1.98	1	1
α Serpentis.....	15. 40	83. 2	+0.95	1	1	A Hydræ.....	9. 21	94. 7	+1.02	1	1
β Aquilæ.....	19. 51	83. 8	+1.10	1	1	β Aquarii.....	21. 27	95. 9	+0.49	2	2
σ Virginis.....	13. 13	84. 0	-0.02	2	2	30 Piscium.....	23. 57	96. 5	+0.74	1	1
ν Tauri.....	3. 58	84. 2	+0.09	2	2	σ^2 Eridani.....	4. 11	97. 7	+1.78	1	1
ω Hydræ.....	9. 1	84. 5	+0.44	1	1	α Hydræ.....	9. 23	98. 2	+0.50	1	1
α Equulei.....	21. 11	85. 1	+0.03	1	1	ξ Aquarii.....	21. 33	98. 2	+1.14	3	3
8 Monocerotis.....	6. 19	85. 3	-0.10	1	1	Spica.....	13. 20	100. 6	+0.73	1	1
β Ophiuchi.....	17. 39	85. 4	+0.03	4	3	2 Libræ.....	14. 19	101. 2	+0.75	1	1
35 Virginis.....	12. 43	85. 9	+0.06	1	1	ζ^1 Libræ.....	14. 49	101. 5	+1.60	1	1
67 Ophiuchi.....	17. 56	87. 0	-0.42	1	1	ν Aquarii.....	21. 5	101. 7	+2.68	1	1
γ Piscium.....	23. 12	87. 2	+1.26	2	2	ν^2 Hydræ.....	10. 1	102. 6	+1.11	1	1
τ Virginis.....	13. 57	88. 0	-0.32	1	1	ρ Ceti.....	2. 22	102. 6	+1.44	1	1
ψ Orionis.....	5. 20	88. 2	+0.71	1	1	λ Leporis.....	5. 15	103. 2	+0.62	1	1
26 Ceti.....	0. 59	89. 1	+0.83	1	1	δ Crateris.....	11. 15	104. 2	+0.08	1	1

The weights used have been determined as follows:—Putting m and n for the number of reflexion and direct results respectively, e for the probable error of one observation, e_0 for the probable systematic error affecting all observations of the same star, the weight to be given is proportional to $\frac{mn}{me^2 + ne^2 + 2mne_0^2}$; or assuming $e_0^2 = \frac{1}{10}e^2$, which would make $e_0 = 0''.16$, the weight becomes $\frac{4mn}{m + n + \frac{1}{5}mn}$, which has been adopted. When $m = n$, the weight is proportional to $\frac{m}{10 + n}$, and $\frac{11m}{10 + m}$, the most convenient multiple of this, has been adopted.

TRANSIT-CIRCLE TABLES.

ciii

*Excess of Reflexion Results above Direct Results from Groups of Stars
observed with the Transit-Circle.*

Extent of Group.	Weight.	Mean N.P.D.	Mean Z.D. South.	Mean Value of R—D, cor- rected for Inclination of Verticals.	R—D computed by the Formula $+ 0'' \cdot 130$ $+ 0'' \cdot 876 \times$ $\sin Z.D.$	Apparent Error of Formula.
		°	°	"	"	"
θ Cephei S.P. to γ Ursæ Majoris S.P.	3	-27.0	-65.5	- 1' 23	- 0' 67	+ 0' 56
ϵ Cephei S.P. to f Draconis S.P.	8	-23.0	-61.5	- 1' 13	- 0' 64	+ 0' 49
Piazzi VII. 67 S.P. to Bradley 348 S.P.	10	-19.3	-57.8	- 1' 12	- 0' 61	+ 0' 51
16 Cephei S.P. to Bradley 1146 S.P.	10	-15.9	-54.4	- 0' 22	- 0' 58	- 0' 36
Piazzi VI. 201 S.P. to Piazzi VII. 75 S.P.	12	-11.9	-50.4	- 0' 83	- 0' 55	+ 0' 29
Bradley 344 S.P. to Bradley 1731 S.P.	10	- 7.3	-45.8	- 0' 56	- 0' 50	+ 0' 06
Bradley 1399 S.P. to Bradley 65 S.P.	16	- 3.5	-42.0	+ 0' 14	- 0' 46	- 0' 60
Polaris S.P. to Groombridge 2283.	22	0.6	-37.9	- 0' 30	- 0' 41	- 0' 11
Groombridge 1004 to Bradley 1731.	18	3.8	-34.7	- 0' 18	- 0' 37	- 0' 19
γ Draconis to α Draconis.	3	8.7	-29.8	- 0' 59	- 0' 31	+ 0' 29
Bradley 1508 to Groombridge 3813.	17	12.9	-25.6	- 0' 18	- 0' 25	- 0' 07
β Ursæ Minoris to γ Cassiopeiæ.	12	17.3	-21.2	- 0' 37	- 0' 19	+ 0' 18
ζ Cygni to μ Herculis.	11	61.0	22.5	+ 0' 46	+ 0' 47	+ 0' 01
ϵ^2 Boötis to 17 Tauri.	16	64.0	25.5	+ 0' 68	+ 0' 51	- 0' 17
1 Geminorum to 5 Vulpeculæ.	16	68.7	30.2	+ 0' 59	+ 0' 57	- 0' 02
ξ Boötis to γ^2 Delphini.	15	72.7	34.2	+ 0' 28	+ 0' 62	+ 0' 34
β Leonis to π Piscium.	16	76.7	38.2	+ 0' 59	+ 0' 67	+ 0' 08
ι Leonis to α Piscium.	14	80.2	41.7	+ 0' 49	+ 0' 71	+ 0' 22
13 Monocerotis to 35 Virginis.	20	84.0	45.5	+ 0' 57	+ 0' 76	+ 0' 19
67 Ophiuchi to 10 Tauri.	8	88.2	49.7	+ 1' 01	+ 0' 80	- 0' 21
η Aquarii to 25 Monocerotis.	10	92.3	53.8	+ 1' 01	+ 0' 84	- 0' 17
Δ Hydræ to ξ Aquarii.	9	97.1	58.6	+ 1' 08	+ 0' 88	- 0' 20
Spica to ν Aquarii.	4	101.3	62.8	+ 1' 58	+ 0' 91	- 0' 67
ν^2 Hydræ to δ Crateris.	4	103.2	64.7	+ 0' 95	+ 0' 92	- 0' 03

In forming the mean values of R—D for groups in the last column but two of the above table, a correction of $+ 0'' \cdot 16 \sin Z.D.$ has been applied to the mean of each group for inclination of the vertical at the surface of the mercury used in reflexion observations to that at the centre of the transit-circle, the mercury trough being mounted so as to describe a circle of 8 feet radius.

Assuming that the R—D correction can be represented by

$$x + y \sin Z.D. \text{ south,}$$

it is found by calculation that the following expressions, when tabulated, will give the values which best agree with the errors in the above, namely :—

civ INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

For direct observations,

$$+ 0''.065 + 0''.438 \sin Z.D.$$

For reflexion observations,

$$- 0''.065 - 0''.438 \sin Z.D. + 0''.16 \sin Z.D.$$

or

$$- 0''.065 - 0''.278 \sin Z.D.$$

Beginning with 1906, the corrections for the R—D discordance have been found by using the value of x from the observations made during the year, and by assuming a constant value of $+0''.600$ for y (practically the value adopted for the 1880, 1890, and 1900 Catalogues), giving:—

For direct observations,

$$+ 0''.065 + 0''.600 \sin Z.D.$$

For reflexion observations,

$$- 0''.065 - 0''.440 \sin Z.D.$$

For use, these formulæ have been tabulated as follows, and the corrections have been applied to the mean results for N.P.D. of the Sun, Moon, and Planets in the section of planetary results throughout the year:—

N.P.D.	Corrections to Results of Direct Observations.	Corrections to Results of Reflexion Observations.
— 45°	— 0''.53	+ 0''.37
40	— 0''.52	+ 0''.37
30	— 0''.49	+ 0''.35
20	— 0''.45	+ 0''.31
— 10	— 0''.38	+ 0''.27
0	— 0''.31	+ 0''.21
+ 10	— 0''.22	+ 0''.15
20	— 0''.13	+ 0''.08
30	— 0''.02	0''.00
40	+ 0''.08	— 0''.08
50	+ 0''.18	— 0''.15
60	+ 0''.29	— 0''.23
70	+ 0''.38	— 0''.30
80	+ 0''.46	— 0''.36
90	+ 0''.53	— 0''.41
100	+ 0''.59	— 0''.45
110	+ 0''.63	— 0''.48
120	+ 0''.66	— 0''.50
+ 125	+ 0''.66	— 0''.50

No correction for flexure has been applied since April 1879.

TRANSIT-CIRCLE TABLES.

CV

*Simultaneous Readings at Apparent Noon of the Transit-Circle Exterior
Thermometer and Meteorological Standard Thermometer.*

Date, 1908.	Meteoro- logical Standard Thermom.	Excess of Transit- Circle Exterior Thermometer.	Date, 1908.	Meteoro- logical Standard Thermom.	Excess of Transit- Circle Exterior Thermometer.	Date, 1908.	Meteoro- logical Standard Thermom.	Excess of Transit- Circle Exterior Thermometer.
Jan. d h			Apr. d h			July d h		
3.0	31.6	- 0.8	13.0	46.9	+ 0.8	27.0	71.5	+ 2.0
10.0	33.7	- 0.6	16.0	57.4	+ 0.1	29.0	70.8	+ 1.0
11.0	32.2	- 0.1	17.0	53.8	+ 0.6	30.0	76.5	+ 0.3
13.0	37.8	- 0.5	20.0	42.1	- 1.1	31.0	70.2	- 0.4
15.0	44.2	+ 0.2	22.0	53.6	- 0.7			
20.0	42.1	- 0.8	24.0	36.5	+ 0.2	Aug. 1.0	66.4	+ 1.5
28.0	44.7	+ 0.2				2.0	70.2	- 1.4
29.0	40.0	- 0.2				4.0	76.6	+ 0.6
30.0	38.2	+ 0.3	May 1.0	70.2	- 0.4	7.0	71.6	+ 0.2
31.0	45.7	+ 0.2	7.0	58.4	+ 1.1	8.0	66.7	+ 0.2
			9.0	66.2	- 0.2	11.0	65.5	- 1.0
Feb. 1.0	40.2	- 0.4	12.0	61.6	+ 0.3	14.0	64.4	+ 0.4
2.0	39.4	- 0.9	16.0	61.2	+ 0.6	15.0	69.1	- 1.1
4.0	41.2	- 0.3	18.0	63.6	+ 2.1	16.0	68.9	+ 1.6
5.0	37.4	+ 0.6	21.0	66.7	+ 2.0	17.0	66.1	+ 0.5
8.0	45.6	+ 0.1	22.0	57.1	- 1.1	21.0	71.3	- 0.3
12.0	51.2	- 0.3	27.0	70.9	+ 0.7	22.0	66.0	+ 0.4
14.0	50.6	- 0.7				25.0	66.0	- 0.6
16.0	45.7	- 0.3				29.0	65.2	+ 0.4
26.0	44.7	+ 0.1	June 1.0	78.6	+ 0.4	30.0	64.4	+ 0.4
29.0	39.1	- 0.5	3.0	76.5	+ 0.4			
			4.0	77.9	+ 0.3	Sept. 1.0	57.1	0.0
			5.0	61.7	+ 0.1	2.0	60.1	- 0.1
Mar. 2.0	39.4	- 0.4	6.0	53.6	+ 0.2	7.0	67.5	+ 1.0
3.0	41.7	- 0.5	11.0	65.5	+ 0.9	8.0	72.6	+ 1.2
5.0	42.1	- 0.1	13.0	63.3	+ 0.7	9.0	62.7	+ 0.6
9.0	46.9	+ 0.3	15.0	66.8	+ 1.2	10.0	56.6	+ 0.5
10.0	48.6	- 1.0	18.0	64.6	+ 0.5	11.0	57.5	- 0.4
12.0	43.9	+ 0.6	22.0	62.5	+ 1.2	12.0	57.6	- 0.3
14.0	42.1	+ 0.4	23.0	70.4	+ 0.4	17.0	67.5	+ 1.9
15.0	41.1	+ 0.3	24.0	70.4	+ 1.4	23.0	64.6	- 1.2
19.0	40.0	+ 0.5	25.0	64.8	+ 1.5	26.0	62.6	- 0.2
20.0	39.6	+ 1.2	29.0	73.1	+ 0.8	30.0	76.6	+ 1.9
24.0	58.3	- 0.7	30.0	72.8	+ 0.2			
27.0	50.6	+ 1.2				Oct. 1.0	75.0	+ 1.3
29.0	50.0	+ 1.8	July 1.0	78.6	0.0	3.0	71.6	+ 0.1
30.0	52.6	+ 0.2	2.0	77.1	+ 0.5	6.0	62.4	+ 0.2
31.0	47.8	0.0	3.0	80.4	- 0.2	7.0	66.0	+ 0.9
			17.0	61.1	- 0.1	12.0	65.1	+ 0.4
Apr. 7.0	50.8	0.0	22.0	72.4	+ 2.5	14.0	59.9	- 0.8
8.0	53.6	+ 0.5	23.0	74.0	+ 0.9	15.0	69.6	+ 0.9
9.0	54.5	+ 1.4	24.0	79.4	+ 1.1	21.0	46.3	- 2.2
			26.0	71.4	+ 0.7			

GREENWICH OBSERVATIONS, 1908.

0

Simultaneous Readings at Apparent Noon of the Transit-Circle Exterior Thermometer and Meteorological Standard Thermometer—continued.

Date, 1908.	Meteoro- logical Standard Thermom.	Excess of Transit- Circle Exterior Thermometer.	Date, 1908.	Meteoro- logical Standard Thermom.	Excess of Transit- Circle Exterior Thermometer.	Date, 1908.	Meteoro- logical Standard Thermom.	Excess of Transit- Circle Exterior Thermometer.
Oct. ^{d h} 29. 0	64°·8	+ 0°·2	Nov. ^{d h} 9. 0	45°·0	— 0°·4	Nov. ^{d h} 25. 0	49°·4	+ 0°·4
30. 0	62°·6	+ 0°·1	10. 0	43°·6	— 1°·4	26. 0	52°·6	0°·0
			12. 0	56°·4	+ 0°·2			
			13. 0	56°·6	— 0°·9			
			17. 0	48°·6	+ 0°·2	Dec. 8. 0	48°·8	— 0°·2
Nov. 3. 0	52°·4	+ 0°·4	20. 0	44°·6	+ 0°·1	22. 0	43°·6	— 0°·6
4. 0	52°·6	— 0°·8	21. 0	47°·8	0°·0	28. 0	28°·5	— 1°·0
7. 0	44°·3	— 0°·3	23. 0	49°·8	0°·0	30. 0	19°·1	+ 1°·1

Means of Simultaneous Readings at Apparent Noon of the Meteorological Standard Thermometer and Transit-Circle Exterior Thermometer.

1908.	January.	February.	March.	April.	May.	June.	Mean for Year.
Meteorological Standard..	39°·0 ₁₀	43°·5 ₁₀	45°·6 ₁₅	49°·9 ₉	64°·0 ₉	68°·3 ₁₅	
Excess of Exterior.	— 0°·2	— 0°·3	+ 0°·3	+ 0°·2	+ 0°·6	+ 0°·7	
1908.	July.	August.	September.	October.	November.	December.	Mean for Year.
Meteorological Standard..	73°·6 ₁₂	67°·9 ₁₅	63°·6 ₁₂	64°·3 ₁₀	49°·5 ₁₃	35°·0 ₄	
Excess of Exterior.	+ 0°·7	+ 0°·1	+ 0°·4	+ 0°·1	— 0°·2	— 0°·2	

TRANSIT-CIRCLE TABLES.

cvii

*Correction to Assumed Colatitude, $38^{\circ} 31' 21'' \cdot 80$,
from Observations in the Year 1908.*

(Using Pulkowa Refractions.)

Corrections for Variation of Latitude supplied by Prof. Albrecht have been applied in forming this Table
(see p. {2} of the section "Ledgers of Mean Right Ascensions and North Polar Distances").

Star's Name.	Approximate R.A.	Approximate N.P.D.	Excess of R.A. above Pole.	Excess of N.P.D. above Pole.	No. of Observations.		Weight.
					Above Pole.	Below Pole.	
	h m	$^{\circ} \quad '$	s				
λ Ursæ Minoris.....	19. 11	1. 0	+0.14	+0.27	25	11	22
Groombridge 1119.....	8. 9	1. 6	-0.78	+0.33	15	25	24
Cephei 24.....	19. 50	1. 9		+1.36	6	2	8
Polaris.....	1. 27	1. 10	+0.15	+0.40	24	35	28
Bradley 1672.....	12. 14	1. 48		+0.24	17	11	21
Groombridge 2283.....	15. 6	2. 25		+0.30	16	17	23
Cephei 51.....	6. 59	2. 48	-0.05	+0.11	17	22	24
Bradley 3147.....	23. 28	3. 11		+0.56	22	12	22
Groombridge 3548.....	21. 18	3. 20	-0.45	-0.17	22	7	18
δ Ursæ Minoris.....	18. 1	3. 23	+0.11	+0.45	39	16	25
2 Ursæ Minoris.....	0. 56	4. 14		+0.86	2	4	7
Bradley 1731.....	12. 48	6. 6		-0.32	4	6	11
76 Draconis.....	20. 49	7. 48		+0.45	2	2	6
Piazzi IX. 37.....	9. 24	8. 16		-0.87	2	6	8
77 Draconis.....	21. 7	12. 14		-0.37	4	4	9
θ Ursæ Minoris.....	15. 34	12. 21		+0.75	6	6	13
19 Ursæ Minoris.....	16. 13	13. 54	-0.06	-0.26	3	1	4
5 Ursæ Minoris.....	14. 28	13. 54		+1.26	4	2	6
Bradley 1147.....	8. 8	13. 58		+0.29	2	2	5
16 Cephei.....	21. 58	17. 15		-0.41	4	2	6
11 Ursæ Minoris.....	15. 17	17. 51		+1.30	4	2	6
f Draconis.....	17. 32	21. 48		-0.06	1	2	4
3 Draconis.....	11. 37	22. 45	+0.25	+0.19	2	3	6
4 Camelopardi.....	4. 41	33. 24	-0.01	-1.43	2	1	2
ψ Andromedæ.....	23. 42	44. 5		(+0.53)	1	1	
ρ Cygni.....	21. 31	44. 48	-0.35	(+1.36)	1	1	
ϵ Aurigæ.....	4. 56	46. 19	-0.18	(-2.30)	1	1	

The weights used in the above tables are determined by use of the "Probable Errors of the Greenwich Observations in Zenith-Distance," given by Mr. Stone in the *Monthly Notices of the Royal Astronomical Society* for 1869 June, page 234. Putting n for the number of observations of a star above the pole; e for the probable error of one observation; n_1 and e_1 the similar quantities for the observations below the pole; e_0 the probable systematic error affecting all observations of the same star, and depending on outstanding division-error, uncertainty in the constant of refraction, &c.—the formula

cviii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

employed to determine the weight to be given to that star is $\frac{2nn_1}{n_1e^2 + ne_1^2 + 2nn_1e_0^2}$; or, assuming $e_0^2 = \frac{1}{10} e_1^2$, which would make $e_0 = 0''.16$, the weight becomes $\frac{2nn_1}{n_1e^2 + ne_1^2 + \frac{1}{5} nn_1e^2}$, which has been adopted in the investigation.

If z be the correction to the assumed colatitude $38^\circ 31' 21''.80$, $2z$ + the algebraic sum of the determinations ought to be equal to 0. Combining the whole with the weights above attached to them, $z = - 0''.138$.

The colatitude determined from the observations of 1908 is therefore $38^\circ 31' 21''.66$.

ALTAZIMUTH TABLES, 1908.

ROYAL OBSERVATORY, GREENWICH.

ALTAZIMUTH TABLES, 1908.

Equivalents of the Transit-Micrometer Screw and Equatorial Intervals of the Transit Wires.

The following are the intervals from the middle wire determined from measures with the transit-micrometer and the position micrometer, combined with star transits. A star transits the wires in the order of numeration of the wires in position 0°, d or 180°, r; in the reverse order in position 0°, r or 180°, d. In each case, the order is reversed below the pole.

One revolution of transit-micrometer = 21''·584.

The following are the equivalents (expressed in equatorial time interval) of the revolutions and tenths of a revolution for the portion of the screw that is in general use, taking the equivalent for 12^{rev.} as the Zero.

Parts of Rev.	rev. 12	rev. 13	rev. 14	rev. 15	rev. 16	rev. 17	rev. 18	rev. 19	rev. 20	rev. 21	rev. 22	rev. 23	rev. 24	Parts of Rev.
r	s	s	s	s	s	s	s	s	s	s	s	s	s	r
·0	·000	1'432	2'864	4'298	5'732	7'169	8'606	10'046	11'488	12'933	14'378	15'825	17'271	·0
·1	·143	1'574	3'007	4'441	5'876	7'313	8'750	10'190	11'632	13'077	14'523	15'970	17'416	·1
·2	·286	1'717	3'150	4'584	6'019	7'457	8'894	10'335	11'777	13'222	14'667	16'114	17'560	·2
·3	·430	1'861	3'294	4'728	6'163	7'599	9'038	10'479	11'921	13'366	14'812	16'259	17'705	·3
·4	·573	2'004	3'437	4'871	6'306	7'743	9'182	10'622	12'066	13'510	14'957	16'404	17'850	·4
·5	·716	2'147	3'580	5'015	6'450	7'887	9'326	10'766	12'210	13'655	15'101	16'547	17'994	·5
·6	·859	2'291	3'724	5'158	6'594	8'031	9'470	10'911	12'355	13'800	15'246	16'692	18'139	·6
·7	1'002	2'434	3'867	5'302	6'738	8'175	9'614	11'055	12'499	13'944	15'391	16'837	18'284	·7
·8	1'146	2'578	4'011	5'445	6'882	8'318	9'758	11'199	12'644	14'089	15'536	16'982	18'429	·8
·9	1'289	2'721	4'154	5'589	7'025	8'462	9'902	11'344	12'788	14'233	15'680	17'126	18'573	·9
1'0	1'432	2'864	4'298	5'732	7'169	8'606	10'046	11'488	12'933	14'378	15'825	17'271	18'718	1'0

ALTAZIMUTH TABLES, 1908.

cxi

A number of wires having been broken on Nov. 2, an entirely new set of wires was inserted.

Equatorial Intervals of the Transit Wires.

Wire.	Jan. 1 to Nov. 2.	Nov. 2 to Dec. 31.	Wire.	Jan. 1 to Nov. 2.	Nov. 2 to Dec. 31.
	^s	^s		^s	^s
1	+ 43'48.1	+ 43'31.4	15	- 2'868	- 2'867
2	+ 33'26.1	+ 33'15.3	16	- 4'364	- 4'283
3	+ 30'36.3	+ 30'24.4	17	- 5'798	- 5'728
4	+ 27'43.4	+ 27'44.1	18	- 7'249	- 6'968
5	+ 24'49.9	+ 24'44.0	19	- 8'784	- 8'570
6	+ 21'69.0	+ 21'63.6	20	- 11'539	- 11'702
7	+ 14'49.7	+ 14'49.5	21	- 14'49.5	- 14'40.7
8	+ 11'47.8	+ 11'46.9	22	- 21'64.9	- 21'62.3
9	+ 8'65.1	+ 8'65.6	23	- 24'54.8	- 24'50.8
10	+ 7'21.0	+ 7'21.2	24	- 27'44.5	- 27'47.2
11	+ 5'81.6	+ 5'82.8	25	- 30'36.1	- 30'24.9
12	+ 4'32.8	+ 4'31.9	26	- 33'32.2	- 33'27.1
13	+ 2'80.9	+ 2'81.2	27	- 43'41.7	- 43'37.9
14	0'00.0	0'00.0			

Equivalents of the Zenith Distance Micrometer-Screw and Intervals of the Zenith Distance Wires.

The following are the intervals from the middle wire, determined from measures with the telescope-micrometer and the position-micrometer, combined with star transits.

A star transits the wires in the order of numeration of the wires when the Fixed Circle is south, i.e., in positions 0°, d, to 180°, d, or 180°, r, to 360°, r.

One revolution of telescope-micrometer = 21''635.

The following corrections have been applied for inequality of screw, to the table deduced with the value 21''680.

Rev.	Correction.	Rev.	Correction.	Rev.	Correction.
	"		"		"
0	+ 0'73	7	+ 0'08	13	+ 0'12
1	+ 0'61	8	+ 0'06	14	+ 0'10
2	+ 0'48	9	+ 0'07	15	+ 0'06
3	+ 0'36	10	+ 0'11	16	0'00
4	+ 0'26	11	+ 0'13		
5	+ 0'18	12	+ 0'13		
6	+ 0'12				

Intervals of the Zenith Distance Wires.

Wire.	Jan. 1 to Nov. 2.	Nov. 2 to Dec. 31.	Wire.	Jan. 1 to Nov. 2.	Nov. 2 to Dec. 31.
I.	+ 346 ^h .67	+ 347 ^h .56	XII.	- 43 ^h .04	- 43 ^h .59
II.	+ 325 ^h .13	+ 326 ^h .46	XIII.	- 65 ^h .06	- 65 ^h .21
III.	+ 303 ^h .55	+ 304 ^h .60	XIV.	- 86 ^h .31	- 86 ^h .89
IV.	+ 282 ^h .03	+ 282 ^h .54	XV.	- 108 ^h .07	(- 107 ^h .81)
V.	+ 259 ^h .53	+ 260 ^h .24	XVI.	- 132 ^h .04	- 128 ^h .27
VI.	+ 130 ^h .05	+ 132 ^h .54	XVII.	- 260 ^h .84	- 259 ^h .37
VII.	+ 108 ^h .58	+ 109 ^h .81	XVIII.	- 281 ^h .80	- 280 ^h .74
VIII.	+ 87 ^h .72	+ 88 ^h .95	XIX.	- 304 ^h .60	- 302 ^h .71
IX.	+ 64 ^h .97	+ 67 ^h .20	XX.	- 326 ^h .32	(- 323 ^h .25)
X.	+ 42 ^h .99	+ 43 ^h .69	XXI.	- 346 ^h .37	- 345 ^h .54
XI.	0 ^h .00	0 ^h .00			

Apparent Correction to adopted Level-Errors, deduced from a Comparison of Reflexion and Direct Observations of Right Ascension, made in the Year 1908.

Name of Star.	Approx. N.P.D.	Approx. R.A.	Seconds of R.A.		R. - D.	Deduced Correction to Level-Error.	Number of Observations.		Weight.
			R.	D.			R.	D.	
ψ Cassiopeiæ S.P.....	-22. 21	1. 19	25 ^s .190	25 ^s .200	-0 ^s .010	+0 ^s .06	1	1	1
B. D. + 80° 395.....	9. 34	12. 43	32 ^s .360	32 ^s .140	+0 ^s .220	+0 ^s .31	1	1	2
Lalande F. 2160.....	9. 52	13. 11	17 ^s .850	18 ^s .060	-0 ^s .210	-0 ^s .30	1	1	2
Groombridge 1852.....	12. 35	12. 1	35 ^s .620	35 ^s .580	+0 ^s .040	+0 ^s .07	1	1	2
Bradley 1446.....	13. 49	10. 27	17 ^s .960	18 ^s .140	-0 ^s .180	-0 ^s .35	1	1	2
Groombridge 3834.....	14. 15	22. 31	39 ^s .790	39 ^s .610	+0 ^s .180	+0 ^s .36	1	1	2
Bradley 473.....	14. 34	3. 28	18 ^s .300	18 ^s .300	0 ^s .000	0 ^s .00	1	1	2
Bradley 417.....	15. 57	3. 2	55 ^s .920	56 ^s .610	-0 ^s .690	-1 ^s .51	1	1	2
Groombridge 737.....	16. 12	3. 46	52 ^s .900	52 ^s .630	+0 ^s .270	+0 ^s .60	1	1	2
Bradley 3085.....	16. 16	23. 11	20 ^s .725	20 ^s .805	-0 ^s .080	-0 ^s .18	2	2	3
Groombridge 2337.....	16. 23	16. 16	3 ^s .490	3 ^s .680	-0 ^s .190	-0 ^s .43	1	1	2
Groombridge 1865.....	16. 56	12. 11	22 ^s .820	22 ^s .620	+0 ^s .200	+0 ^s .46	1	1	2
Groombridge 3702.....	18. 4	22. 5	28 ^s .530	28 ^s .000	+0 ^s .530	+1 ^s .31	1	1	2
Lalande 6061.....	18. 7	3. 17	4 ^s .680	4 ^s .770	-0 ^s .090	-0 ^s .22	1	1	2
Bradley 3166.....	22. 42	23. 44	30 ^s .505	30 ^s .495	+0 ^s .010	+0 ^s .03	2	2	3
Bradley 1429.....	22. 58	10. 18	30 ^s .470	30 ^s .660	-0 ^s .190	-0 ^s .57	1	1	2
Groombridge 1962.....	50. 59	13. 5	28 ^s .070	28 ^s .000	+0 ^s .070	+0 ^s .41	1	1	2
τ Coronæ.....	53. 17	16. 6	36 ^s .350	36 ^s .410	-0 ^s .060	-0 ^s .37	1	1	2
14 Canum Venaticum...	53. 43	13. 1	26 ^s .560	26 ^s .390	+0 ^s .170	+1 ^s .04	1	1	2
54 Persei.....	55. 39	4. 14	26 ^s .060	26 ^s .090	-0 ^s .030	-0 ^s .19	1	1	2
σ Boötis.....	59. 51	14. 31	40 ^s .410	40 ^s .505	-0 ^s .095	-0 ^s .66	1	2	2
ψ Tauri.....	61. 15	4. 1	19 ^s .080	19 ^s .080	0 ^s .000	0 ^s .00	1	1	2
β Tauri.....	61. 28	5. 20	28 ^s .520	28 ^s .460	+0 ^s .060	+0 ^s .42	1	2	2
11 Boötis.....	62. 10	13. 57	0 ^s .330	0 ^s .255	+0 ^s .075	+0 ^s .54	1	2	2
38 Lynceis.....	62. 48	9. 13	7 ^s .060	7 ^s .260	-0 ^s .200	-1 ^s .44	1	1	2
ϵ Coronæ.....	62. 51	15. 54	46 ^s .740	46 ^s .780	-0 ^s .040	-0 ^s .29	1	1	2
μ Leonis.....	63. 34	9. 48	32 ^s .020	32 ^s .075	-0 ^s .055	-0 ^s .41	1	4	3
α Arietis.....	66. 58	2. 2	59 ^s .000	59 ^s .020	-0 ^s .020	-0 ^s .16	1	3	3

From 16 North Stars, Mean Correction to Level-Error = -0^h.03.

„ 12 South Stars, „ „ „ = -0^h.10.

ALTAZIMUTH TABLES, 1908.

cxiii

Excess of Reflexion Results above Direct Results for Observations of Zenith-Distance with the Altazimuth, 1908.

Star's Name.	Approximate R.A.	Approximate N.P.D.	R.—D.	No. of Obs. R. and D.	Weight.
Position 180°, r. January, February.					
ζ Draconis S.P.	h m 17. 9	° ' " — 24. 10	— 2°08	1	1
ζ Ursæ Minoris S.P.	15. 47	— 11. 55	+ 0°26	1	1
Bradley 1672 S.P.	12. 14	— 1. 47	— 0°93	1	1
Piazzi VII. 132	7. 41	9. 30	+ 1°10	1	1
Piazzi VII. 187	7. 50	10. 16	+ 1°14	1	1
Groombridge 1030	5. 52	14. 25	+ 1°02	1	1
Bradley 473	3. 28	14. 34	— 2°53	1	1
Groombridge 379	1. 43	14. 52	— 4°49	1	1
Groombridge 948	5. 15	15. 32	— 0°54	1	1
Groombridge 737	3. 46	16. 12	— 0°95	1	1
Bradley 1160	8. 11	17. 18	— 0°13	1	1
Lalande 6061	3. 17	18. 7	— 1°22	1	1
γ Camelopardi	3. 40	18. 57	+ 1°32	1	1
54 Persei	4. 14	55. 39	+ 0°08	1	1
ψ Tauri	4. 1	61. 15	— 0°61	1	1
β Tauri	5. 20	61. 28	— 0°49	1	1
ι Geminorum	7. 20	62. 1	— 1°28	1	1
ω ¹ Cancri	7. 55	64. 21	— 2°23	1	1
ξ Tauri	3. 22	80. 35	— 3°38	1	1
ζ ¹ Piscium	1. 9	82. 55	— 1°40	1	1
μ Piscium	1. 25	84. 20	— 0°88	1	1
δ Orionis	5. 27	90. 22	— 3°35	1	1
ζ Orionis	5. 36	91. 59	+ 0°34	1	1
ε Eridani	3. 29	99. 46	— 0°26	1	1
Position 180°, d. March, April.					
κ Cephei S.P.	20. 12	— 12. 34	— 0°24	1	1
Groombridge 1852	12. 1	12. 35	— 0°33	1	1
Bradley 1446	10. 27	13. 49	+ 0°50	1	1
Groombridge 1865	12. 11	16. 56	— 0°11	1	1
Groombridge 1998	13. 23	17. 15	+ 2°69	1	1
Piazzi X. 126	10. 36	20. 27	— 1°20	1	1
Bradley 1429	10. 17	23. 58	— 0°25	1	1

GREENWICH OBSERVATIONS, 1908.

p

Excess of Reflexion Results above Direct Results for Observations of Zenith-Distance with the Altazimuth, 1908—contd.

Star's Name.	Approximate R.A.	Approximate N.P.D.	R.—D.	No. of Obs. R. and D.	Weight.
Position 180°, d. March, April—continued.					
	h m	° '	"		
Groombridge 1962	13. 5	50. 59	+ 0.65	1	1
α Canum Venaticum	12. 51	51. 11	— 0.58	1	1
38 Lyncis	9. 13	52. 48	0.00	1	1
14 Canum Venaticum	13. 1	53. 43	+ 0.30	1	1
ρ_1 Cancri	8. 46	61. 19	— 0.96	1	1
11 Boötis	13. 57	62. 10	+ 1.20	1	1
μ Leonis	9. 48	63. 34	+ 0.03	1	1
ω^1 Cancri	7. 55	64. 31	— 2.19	1	1
μ Cancri	8. 2	68. 9	— 1.19	1	1
η Leonis	10. 2	72. 47	— 2.35	1	1
Position 0°, d. May, June.					
γ Cassiopeiae S.P.	0. 51	— 29. 50	+ 1.55	1	1
Bradley 82 S.P.	0. 44	— 26. 15	— 1.00	1	1
ψ Cassiopeiae S.P.	1. 19	— 22. 21	— 2.99	1	1
γ Camelopardi S.P.	3. 40	— 18. 57	— 0.23	1	1
Bradley 117 S.P.	1. 4	— 10. 49	(— 3.96)	1	1
B. D. + 80° 395	12. 43	9. 34	+ 2.23	1	1
Lalande F (2) 160	13. 11	9. 52	— 1.20	1	1
ζ Ursæ Minoris	15. 47	11. 55	+ 0.28	1	1
Groombridge 2337	16. 16	16. 23	— 0.53	1	1
τ Coronæ	16. 6	53. 17	— 0.81	1	1
σ Boötis	14. 31	59. 51	+ 0.27	1	1
11 Boötis	13. 57	62. 10	+ 0.12	1	1
14 Comæ	12. 21	62. 13	— 2.14	1	1
ϵ Coronæ	15. 54	62. 51	— 0.55	1	1
ϵ Virginis	12. 58	78. 33	+ 0.73	1	1
δ Virginis	12. 51	86. 6	— 0.28	1	1
β Libræ	15. 12	99. 3	+ 1.56	1	1
Position 0°, r. July, August.					
α Ursæ Majoris S.P.	8. 23	— 28. 58	— 1.01	1	1
43 Camelopardi S.P.	6. 43	— 21. 0	+ 0.58	1	1
Piazzi IX, 187 S.P.	9. 50	— 16. 41	+ 2.35	1	1

ALTAZIMUTH TABLES, 1908.

CXV

Excess of Reflexion Results above Direct Results for Observations of Zenith-Distance with the Altazimuth, 1908—contd.

Star's Name.	Approximate R.A.	Approximate N.P.D.	R.—D.	No. of Obs. R. and D.	Weight.
Position 0°, r. July, August— <i>continued</i> .					
	^h ^m	[°] [']	["]		
Piazzi VI. 75 S.P.	6. 30	— 10. 20	(+ 5'45)	1	1
ζ Cygni	21. 9	60. 9	+ 0'85	1	1
32 Vulpeculæ	20. 51	62. 18	— 1'34	1	1
15 Vulpeculæ	19. 57	62. 30	— 0'49	1	1
c Serpentis	18. 24	92. 3	— 0'71	1	1
η Serpentis	18. 16	92. 55	+ 0'96	1	1
Position 180°, r. September, October.					
σ ² Ursæ Majoris S.P.	9. 2	— 22. 29	— 2'41	1	1
43 Camelopardi S.P.	6. 43	— 21. 0	+ 0'90	1	1
Bradley 1446 S.P.	10. 27	— 13. 49	— 0'06	1	1
Piazzi VI. 75 S.P.	6. 30	— 10. 20	+ 1'46	1	1
Groombridge 3548	21. 18	3. 21	+ 0'61	1	1
Piazzi XXII. 96	22. 17	13. 58	— 0'14	1	1
Groombridge 3834	22. 31	14. 15	— 0'24	1	1
Bradley 3085	23. 11	16. 16	+ 0'45	2	2
16 Cephei	21. 58	17. 15	+ 2'70	1	1
Groombridge 3702	22. 5	18. 5	— 2'66	1	1
Bradley 3166	23. 44	22. 42	+ 0'65	2	2
δ Andromedæ	0. 34	59. 39	— 1'75	1	1
ζ Cygni	21. 9	60. 9	— 2'40	1	1
16 Pegasi	21. 48	64. 31	— 1'65	2	2
κ Pegasi	21. 40	64. 47	(— 4'69)	1	1
α Aquarii	22. 1	90. 46	+ 1'04	1	1
c Serpentis	18. 24	92. 3	— 0'36	1	1
η Serpentis	18. 16	92. 55	+ 0'83	1	1
η Aquarii	20. 52	100. 3	— 2'81	1	1
Position 180°, d. November, December.					
Bradley 417	3. 2	15. 57	— 0'17	1	1
Groombridge 558	2. 46	17. 0	— 0'17	1	1
α Arietis	2. 2	66. 58	— 2'51	1	1
27 Arietis	2. 26	72. 42	— 1'18	1	1

*Excess of Reflexion Results above Direct Results from Groups of Stars
observed with the Altazimuth.*

Extent of Group 1908.	Weight.	Mean N.P.D.	Mean Z.D. South.	Mean Value of R.—D., corrected for Inclination of Verticals.	R.—D. computed by the Formula.	Apparent Error of the Formula.
Position 180°, r. January, February.						
ζ Draconis S.P.....	1	— 24. 10	— 62. 41	— 2'19	+ 0'24	+ 2'43
ξ Ursæ Minoris S.P.....	1	— 11. 55	— 50. 26	+ 0'17	+ 0'19	+ 0'02
Bradley 1672 S.P.....	1	— 1. 47	— 40. 18	— 1'01	+ 0'14	+ 1'15
Piazzi VII. 132 to Piazzi VII. 187.....	2	9. 53	28. 38	+ 1'06	— 0'31	— 1'37
Groombridge 1030 to γ Camelopardi.....	8	16. 15	22. 16	— 0'99	— 0'27	+ 0'72
54 Persei to ω ¹ Cancri.....	5	60. 57	22. 26	— 0'86	— 0'27	+ 0'59
ξ Tauri to μ Piscium.....	3	82. 37	44. 6	— 1'81	— 0'40	+ 1'41
δ Orionis to ε Eridani.....	3	94. 2	55. 31	— 0'99	— 0'45	+ 0'54
Position 180°, d. March, April.						
κ Cephei S.P.....	1	— 12. 34	— 51. 5	— 0'33	+ 0'43	+ 0'76
Groomb. 1852 to Groomb. 1998.....	4	15. 9	— 23. 22	+ 0'64	+ 0'28	— 0'36
Piazzi X. 126 to Bradley 1429.....	2	22. 13	— 16. 18	— 0'75	+ 0'23	+ 0'98
Groomb. 1962 to 14 Canum Venaticum.....	4	52. 10	13. 39	+ 0'12	+ 0'03	— 0'09
ρ ¹ Cancri to ω ¹ Cancri.....	4	62. 54	24. 23	— 0'43	— 0'04	+ 0'39
μ Cancri to η Leonis.....	2	70. 28	31. 57	— 1'71	— 0'09	+ 1'62
Position 0°, d. May, June.						
γ Cassiopeiæ S.P. to γ Camelopardi S.P. B. D. + 80° 395 to Groombridge 2337.	4	— 24. 21	— 62. 52	— 0'78	+ 0'24	+ 1'02
τ Coronæ to ε Coronæ.....	5	60. 4	21. 33	— 0'58	— 0'27	+ 0'31
ε Virginis to δ Virginis.....	2	82. 20	43. 49	+ 0'30	— 0'40	— 0'70
β Libræ.....	1	99. 3	60. 31	+ 1'66	— 0'47	— 2'13
Position 0°, r. July, August.						
ο Ursæ Majoris S.P.....	1	— 28. 58	— 67. 29	— 1'12	+ 0'49	+ 1'61
43 Camelopardi S.P.....	1	— 21. 0	— 59. 31	+ 0'48	+ 0'46	— 0'02
Piazzi IX. 187 S.P.....	1	— 16. 41	— 55. 12	+ 2'25	+ 0'45	— 1'80
Piazzi VI. 75 S.P.....	1	— 10. 20	— 48. 51	(+ 5'36)	+ 0'42	(— 4'94)
ζ Cygni.....	1	60. 9	21. 38	+ 0'89	— 0'03	— 0'92
32 Vulpeculæ.....	1	62. 18	23. 47	— 1'29	— 0'04	+ 1'25
15 Vulpeculæ.....	1	62. 30	23. 59	— 0'44	— 0'04	+ 0'40
ε Serpentis.....	1	92. 3	53. 32	— 0'61	— 0'20	+ 0'41
η Serpentis.....	1	92. 55	54. 24	+ 1'06	— 0'21	— 1'27

*Excess of Reflexion Results above Direct Results from Groups of Stars
observed with the Altazimuth—contd.*

Extent of Group 1908.	Weight.	Mean N.P.D.	Mean Z.D. South.	Mean Value of R.—D. corrected for Inclination of Verticals.	R.—D. computed by the Formula.	Apparent Error of the Formula.
Position 180°, r. September, October.						
σ ² Ursæ Maj. S.P. to Piazzi VI. 75 S.P.	4	— 16. 55	— 55. 26	— 0.13	+ 0.21	+ 0.34
Groombridge 3548	1	3. 21	— 35. 10	+ 0.54	+ 0.12	— 0.42
Piazzi XXII. 96 to Bradley 3166	8	17. 41	— 20. 50	+ 0.19	+ 0.02	— 0.17
δ Andromedæ to κ Pegasi	4	62. 12	23. 41	— 1.81	— 0.29	+ 1.52
α Aquarii to η Aquarii	4	93. 57	55. 26	— 0.23	— 0.45	— 0.22
Position 180°, d. November, December.						
Bradley 417	1	15. 57	— 22. 34	— 0.22	+ 0.27	+ 0.49
Groombridge 558	1	17. 0	— 21. 31	— 0.21	+ 0.27	+ 0.48
α Arietis	1	66. 58	28. 27	— 2.45	— 0.07	+ 2.38
27 Arietis	1	72. 42	34. 11	— 1.11	— 0.10	+ 1.01

In forming the mean values of R—D for groups in the fifth column of the above table, a correction of $+0''.12 \sin Z.D.$ has been applied to the mean of each group, for inclination of the vertical at the surface of the mercury used in reflexion observations to that at the centre of the instrument, the centre of the mercury trough being assumed to describe a circle of 6 feet radius.

Assuming that the R—D discordance can be represented by

$$x + y \sin Z.D. \text{ south,}$$

it is found by calculation that the following expressions, when tabulated, will give the values which best agree with the errors in the above, namely :—

*Observed R—D Discordance for Different Positions of the Altazimuth,
in the Year 1908.*

Position of Instrument.	R—D Discordance.	
180°, r (Jan., Feb.).	—0.90	—0.44 sin Z.
180°, d (Mar., Apr.).	—0.20	—0.72 sin Z.
0°, d (May, June).	—0.18	+0.20 sin Z.
0°, r (July, Aug.).	+0.16	—0.44 sin Z.
180°, r (Sept., Oct.).	—0.40	—1.00 sin Z.
180°, d (Nov., Dec.).	—0.88	—1.74 sin Z.

Assuming the formulæ for the four positions as

$$+ a + b \sin Z.D. \text{ for the positions } 0^\circ, d, \text{ and } 180^\circ, r,$$

and

$$- a + b \sin Z.D. \quad , , \quad , , \quad 0^\circ, r, \text{ and } 180^\circ, d,$$

the correction for the R - D discordance in the four positions of the instrument has been taken as

$$\mp 0''.06 - 0''.20 \sin Z.D.$$

giving a little weight to observations made at the end of 1907 and beginning of 1909.

Comparison of North Polar Distances observed with the Altazimuth in four different positions, in the Year 1908, with those of Newcomb's Fundamental Catalogue.

Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.
R.A.	N.P.D.				R.A.	N.P.D.				R.A.	N.P.D.			
Position 180°, r. January, February.					Position 180°, r. Jan., Feb.—contd.					Position 180°, r. Jan., Feb.—contd.				
h	m	°	"		h	m	°	'		h	m	°	"	
17. 28	— 37.6	+	0.28	I I	2. 7	23.9	— 0.19	I I	4. 53	65.1	+	0.44	I I	
18. 44	— 37.1	+	0.14	I I	5. 21	27.1	+	0.10	I I	3. 44	66.2	+	0.38	I I
					4. 55	29.7	— 0.50	2 2	1. 53	66.9	+	0.02	I I	
20. 43	— 28.5	+	1.34	I I					2. 2	67.1	+	0.42	I I	
20. 10	— 28.2	+	0.02	I I	7. 35	31.1	+	0.20	I I	3. 59	68.2	— 0.22	I I	
18. 13	— 25.6	— 0.29	I I	I I	4. 40	33.4	+	0.70	I I	4. 58	68.5	— 0.60	I I	
18. 36	— 24.6	— 0.56	2 2	2 2	2. 44	34.5	+	0.29	I I	11. 9	69.0	— 1.14	I I	
19. 9	— 24.2	+	2.08	I I					2. 54	69.0	+	0.13	3 3	
17. 9	— 24.2	— 0.81	2 2	2 2	5. 39	40.2	+	0.48	I I	3. 16	69.2	— 0.26	2 2	
19. 13	— 22.5	— 0.37	I I	I I	2. 38	41.2	— 1.51	I I	8. 27	69.3	— 0.31	2 2		
16. 6	— 22.0	+	0.04	I I	3. 24	42.3	+	0.19	I I	3. 10	69.3	+	0.73	I I
					4. 2	42.5	+	0.33	I I	1. 50	69.6	— 0.16	I I	
19. 18	— 16.8	— 0.36	I I	I I	5. 53	45.1	— 0.08	2 2	6. 24	69.7	+	0.92	2 2	
14. 51	— 15.5	— 0.62	I I	I I	4. 27	47.1	— 0.05	I I						
18. 49	— 14.7	— 1.61	I I	I I					4. 23	71.0	— 0.73	I I		
14. 28	— 13.9	+	0.33	I I	5. 45	50.9	— 1.14	I I	4. 46	71.3	— 0.49	I I		
15. 47	— 11.9	+	0.67	I I	3. 53	54.5	+	1.21	I I	8. 18	71.4	+	0.26	I I
					2. 4	55.5	— 0.30	2 2	9. 13	71.9	— 1.07	I I		
18. 3	— 3.4	— 0.46	4 3	3 3	4. 14	55.7	— 0.25	I I	2. 26	72.7	+	0.15	2 2	
18. 5	— 3.0	+	0.90	2 2	7. 29	57.9	— 1.67	4 3	6. 32	73.5	— 0.23	I I		
15. 7	— 2.4	+	0.08	4 3	7. 29	57.9	— 0.89	4 3	7. 8	73.7	— 1.23	I I		
12. 14	— 1.8	— 0.65	I I	I I	3. 39	58.0	+	1.04	I I	4. 14	74.6	— 1.46	I I	
19. 13	— 1.0	+	1.08	I I					4. 59	74.7	— 0.84	I I		
									2. 46	75.3	+	0.44	2 2	
1. 26	1.2	— 0.33	I I	I I	2. 42	61.1	+	0.67	I I	8. 23	75.5	— 1.40	I I	
6. 57	2.8	— 0.15	4 3	3 3	4. 1	61.3	— 0.63	I I	6. 7	75.8	+	0.11	I I	
5. 32	4.9	+	0.79	I I	5. 20	61.5	— 0.30	I I	4. 47	75.9	— 0.42	I I		
					7. 20	62.1	— 0.15	I I	6. 49	76.7	+	3.66	I I	
3. 40	19.0	+	0.25	I I	14. 40	62.5	— 0.10	I I	6. 11	77.7	— 1.05	I I		
					15. 30	63.0	+	0.24	I I	3. 56	77.8	+	0.37	I I
2. 37	22.6	— 0.17	I I	I I	2. 45	63.1	— 0.89	2 2	2. 40	77.9	+	0.06	2 2	
2. 21	23.0	+	0.87	I I	7. 55	64.4	+	0.30	2 2	2. 32	78.0	+	0.26	I I
4. 45	23.8	— 0.46	I I	I I	6. 38	64.8	+	0.09	I I	3. 25	79.0	— 0.29	I I	

Comparison of North Polar Distances observed with the Altazimuth in four different positions, in the Year 1908, with those of Newcomb's Fundamental Catalogue—contd.

Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.
R.A.	N.P.D.				R.A.	N.P.D.				R.A.	N.P.D.			
Position 180°, r. Jan., Feb.— <i>contd.</i>					Position 180°, r. Jan., Feb.— <i>contd.</i>					Position 180°, d. Mar. Apr.— <i>contd.</i>				
h	m	°	"		h	m	°	"		h	m	°		
6. 36		80°0	— 1'04	1 1	6. 41		106°6	— 0'25	2 2	7. 29		57°9	— 0'28	1 1
5. 57		80°4	— 0'22	1 1	5. 29		107°9	— 0'92	1 1	7. 29		57°9	— 0'81	2 2
8. 11		80°5	+ 0'38	1 1	15. 6		109°5	— 0'62	2 2	10. 41		58°8	— 0'12	1 1
3. 22		80°6	+ 0'70	1 1						8. 49		59°1	+ 0'24	1 1
1. 40		81°3	— 0'95	1 1	5. 24		110°8	+ 1'17	2 2	8. 49		59°1	— 1'37	1 1
3. 20		81°3	— 1'12	3 3	5. 47		110°9	— 0'07	2 2					
4. 11		81°3	+ 0'32	1 1	3. 30		111°9	— 0'28	1 1					
7. 22		81°5	— 0'84	2 2	5. 2		112°5	— 1'84	1 1	8. 46		61°3	+ 0'16	1 1
2. 8		81°6	+ 0'17	1 1	7. 45		114°6	+ 0'55	3 3	13. 8		61°7	+ 0'79	1 1
2. 23		82°0	+ 0'21	1 1						7. 40		61°8	— 0'87	1 1
5. 50		82°6	— 1'19	1 1						7. 58		62°0	— 0'40	3 3
11. 56		82°9	— 1'80	1 1						12. 47		62°0	— 0'55	2 2
1. 9		82°9	— 0'80	1 1	Position 180°, d. March, April.									
1. 25		84°3	— 1'43	1 1	21. 43		— 41°1	— 1°40	2 2	9. 13		62°8	+ 0'35	2 2
1. 37		85°0	+ 1'86	1 1	20. 43		— 28°5	— 0'55	1 1	9. 48		63°6	— 0'55	2 2
6. 19		85°4	— 0'33	2 2	22. 46		— 24°3	— 0'28	1 1	17. 27		63°8	— 0'63	1 1
8. 34		86°3	+ 2'09	1 1	20. 58		— 17°3	— 0'87	1 1	17. 51		63°9	— 0'84	2 2
4. 49		87°7	— 1'00	1 1	20. 12		— 12°6	+ 0'19	1 1	7. 55		64°4	— 0'40	1 1
23. 48		88°4	— 1'46	1 1						10. 51		64°8	— 1'13	1 1
										8. 57		65°2	— 0'35	1 1
										10. 12		66°1	— 1'20	1 1
2. 34		90°1	+ 0'01	1 1	21. 18		— 3°4	— 0'43	2 2	12. 30		66°9	— 0'08	1 1
11. 32		90°3	— 1'60	1 1	23. 27		— 3°2	+ 0'02	2 2	7. 15		67°9	— 0'46	1 1
5. 27		90°4	— 1'49	2 2	1. 26		— 1°2	+ 0'46	2 2	8. 2		68°2	+ 0'37	1 1
5. 17		90°5	— 0'18	1 1						10. 15		69°7	— 0'53	1 1
5. 36		92°0	+ 0'16	1 1	8. 3		1°1	+ 0'76	3 3					
5. 34		92°7	— 1'07	2 2	12. 14		1°8	+ 0'08	2 2	14. 22		70°4	+ 0'86	1 1
2. 14		93°4	— 1'81	1 1	18. 3		3°4	+ 0'10	1 1	8. 18		71°4	— 0'81	1 1
16. 9		93°5	+ 0'66	1 1	10. 27		13°8	+ 0'04	1 1	8. 39		71°5	— 0'13	1 1
4. 32		93°5	— 1'37	2 2	10. 36		20°5	+ 0'58	2 2	9. 14		71°9	+ 0'03	2 2
0. 25		94°5	+ 0'05	1 1	10. 18		24°0	— 0'42	1 1	10. 2		72°8	— 0'70	2 2
5. 3		95°2	— 0'63	1 1	11. 49		35°8	+ 1°92	1 1	8. 23		75°5	— 0'10	1 1
5. 5		95°9	— 0'94	1 1	9. 27		37°9	— 0'47	1 1	9. 53		77°1	+ 0'15	1 1
23. 57		96°5	— 0'90	1 1	8. 2		38°2	+ 0'18	1 1	10. 3		77°6	— 1'13	1 1
9. 22		98°3	— 0'98	1 1						10. 44		79°0	— 1'73	1 1
5. 10		98°3	— 0'77	3 3						11. 19		79°0	+ 0'50	1 1
15. 12		99°1	— 1'01	1 1	11. 4		45°0	— 0'55	1 1	9. 36		79°7	— 0'97	1 1
2. 52		99°3	— 0'48	1 1	9. 52		48°5	— 2°11	1 1					
3. 29		99°8	— 0'43	1 1	13. 11		48°7	+ 0°96	1 1	10. 28		80°2	— 0'87	3 3
					13. 16		49°4	— 0°60	1 1	12. 0		80°8	+ 1°48	1 1
										9. 55		81°5	+ 0°08	2 2
16. 31		100°4	+ 1°69	1 1	12. 52		51°2	+ 0°72	1 1	12. 41		81°8	+ 4°11	1 1
2. 21		102°7	— 0°77	1 1	11. 48		51°6	— 0°30	2 2	11. 0		82°2	— 1°71	2 2
5. 15		103°3	— 0°47	2 2	17. 14		52°6	— 0°22	1 1	10. 30		82°6	+ 0°20	1 1
7. 42		104°3	— 0°15	1 1										
15. 55		106°3	+ 1°34	1 1										
15. 23		106°4	+ 0°64	1 1										

ALTAZIMUTH TABLES, 1908.

cxi

Comparison of North Polar Distances observed with the Altazimuth in four different positions, in the Year 1908, with those of Newcomb's Fundamental Catalogue—contd.

Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.
R.A.	N.P.D.				R.A.	N.P.D.				R.A.	N.P.D.			
Position 0°, r. July, August—contd.					Position 0°, r. July, August—contd.					Position 180°, r. Sept., Oct.—contd.				
h	m	°	"		h	m	°	"		h	m	°	"	
19. 13	22.5	-	0.47	1 1	18. 30	98.3	+	0.46	2 2	19. 13	1.0	-	0.20	1 1
19. 9	24.2	+	0.64	1 1	18. 37	99.1	+	0.60	1 1	6. 57	2.8	-	0.44	1 1
16. 23	28.3	-	0.37	1 1	17. 54	99.8	+	2.32	1 1	23. 27	3.2	-	0.33	4 3
20. 43	28.5	-	1.54	1 1						21. 18	3.4	-	0.61	3 3
16. 0	31.2	+	0.28	1 1	21. 5	101.8	+	1.09	1 1	18. 3	3.4	+	0.99	1 1
18. 45	37.1	-	1.51	1 1	17. 5	105.6	-	0.48	1 1	20. 52	9.8	+	0.25	1 1
20. 54	49.2	+	0.73	1 1	19. 35	106.5	-	0.56	1 1					
19. 16	49.8	-	1.84	1 1	23. 59	107.9	+	0.89	1 1	22. 41	14.3	-	0.96	1 1
20. 19	50.0	-	2.62	1 1	20. 23	108.1	-	2.08	1 1	18. 49	14.7	+	1.71	1 1
18. 33	51.3	+	0.10	1 1	23. 5	111.7	-	1.10	1 1	21. 52	16.7	+	0.88	1 1
19. 4	54.1	+	0.57	1 1	17. 16	114.9	+	1.40	1 1	21. 58	17.3	+	0.85	5 4
19. 28	55.8	-	2.58	1 1	19. 31	115.1	+	4.41	1 1	21. 27	19.9	-	2.05	1 1
22. 5	57.3	-	0.41	1 1	16. 16	115.4	+	0.74	1 1					
21. 9	60.2	+	0.60	1 1	18. 22	115.5	+	0.68	2 2	1. 19	22.4	-	0.29	1 1
19. 27	62.2	+	0.34	1 1	15. 53	115.9	+	0.11	1 1	19. 13	22.5	+	0.32	1 1
20. 51	62.3	-	1.34	2 2	18. 40	117.1	+	2.61	1 1	23. 44	22.7	+	0.18	2 2
19. 57	62.5	+	0.04	1 1	19. 57	118.0	-	1.36	1 1	22. 46	24.3	-	0.02	2 2
14. 41	62.5	+	0.65	1 1	22. 53	120.1	+	0.75	1 1	0. 44	26.3	-	0.64	1 1
17. 51	63.9	+	2.46	1 1	18. 0	120.4	+	2.31	2 2	22. 2	27.7	+	2.45	1 1
22. 2	65.1	+	1.68	1 1	Position 180°, r. September, October.									
19. 24	65.5	-	1.41	1 1	8. 57	- 42.5	(- 5.36)	1 1	20. 10	28.2	+	0.46	1 1	
1. 49	69.6	+	0.30	1 1	8. 53	- 41.6	(- 2.68)	1 1	23. 21	28.2	-	0.86	1 1	
15. 42	74.3	+	0.53	1 1	9. 27	- 37.9	+	0.69	3 3	20. 43	28.5	-	0.24	2 2
18. 55	75.1	+	1.33	1 1	11. 49	- 35.8	+	1.42	2 2	0. 51	29.8	-	0.46	1 1
0. 8	75.3	-	0.08	1 1	8. 57	- 35.4	- 3.70	1 1	22. 26	32.1	+	0.91	1 1	
20. 31	75.7	-	1.19	1 1	12. 50	- 33.5	- 1.61	1 1	19. 35	40.0	-	0.05	1 1	
16. 21	75.8	+	1.85	1 1	7. 35	- 31.1	+	0.46	3 3	21. 43	41.1	-	0.49	2 2
17. 31	77.4	+	1.62	1 1	9. 44	- 30.5	- 0.36	1 1	1. 32	41.8	+	0.94	1 1	
19. 46	81.4	-	1.08	1 1	8. 23	- 29.0	- 1.09	1 1	21. 3	42.7	+	0.33	1 1	
19. 29	82.8	-	0.60	2 2	10. 18	- 24.0	+	0.34	2 2	21. 31	44.8	-	0.60	1 1
1. 36	85.0	+	0.76	1 1	9. 2	- 22.5	- 1.49	1 1	20. 38	45.1	+	1.23	1 1	
17. 39	85.4	+	3.10	1 1	6. 43	- 21.0	+	0.16	1 1	19. 42	45.1	+	0.67	2 2
19. 9	87.9	-	1.87	1 1	10. 36	- 20.5	- 0.71	2 2	23. 36	46.2	+	0.40	1 1	
22. 21	89.1	+	0.37	1 1	9. 34	- 20.4	+	0.13	2 2	22. 57	48.2	+	1.09	1 1
23. 22	89.3	-	3.26	1 1	14. 28	- 13.9	+	0.14	1 1	22. 40	48.7	-	1.12	1 1
					10. 27	- 13.8	- 0.66	3 3	20. 54	49.2	+	1.20	1 1	
20. 7	91.1	-	1.52	1 1	12. 1	- 12.6	+	0.78	2 2	20. 19	50.0	-	0.19	1 1
0. 48	91.7	-	0.89	1 1	10. 53	- 11.7	- 0.92	1 1	18. 47	56.8	-	0.20	2 2	
18. 24	92.1	+	0.47	1 1	6. 30	- 10.3	- 0.44	1 1	0. 32	56.8	-	0.35	1 1	
18. 16	92.9	+	0.53	2 2	6. 57	- 2.8	+	0.55	1 1	22. 5	57.3	-	0.11	3 3
18. 42	94.9	-	2.04	1 1	12. 14	- 1.8	+	0.15	2 2	22. 5	57.3	+	0.26	1 1
21. 26	96.0	+	1.06	1 1	8. 3	- 1.1	+	0.53	6 4	0. 34	59.7	-	0.65	1 1
22. 11	98.3	-	0.67	1 1						21. 9	60.2	+	0.48	3 3
										21. 46	60.3	-	0.04	1 1
										20. 51	62.3	+	0.15	1 1

GREENWICH OBSERVATIONS, 1908.

q

xxx INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Comparison of North Polar Distances observed with the Altazimuth in four different positions, in the Year 1908, with those of Newcomb's Fundamental Catalogue—contd.

Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight.
R.A.	N.P.D.				R.A.	N.P.D.				R.A.	N.P.D.			
Position 180°, d. Mar., Apr.—contd.					Position 0°, d. May, June—contd.					Position 0°, d. May, June—contd.				
h	m	°	"		h	m	°	"		h	m	°	"	
9. 32		82.8	— 0.75	1 1	6. 57		— 2.8	— 0.44	1 1	17. 31		77.4	+ 0.40	1 1
11. 56		82.9	+ 0.34	2 2	1. 26		— 1.2	— 0.54	2 2	12. 58		78.6	— 0.11	1 1
11. 41		83.0	+ 0.29	1 1										
10. 41		83.2	— 0.17	1 1	12. 14		1.8	— 1.84	2 2	16. 53		80.5	— 0.27	1 1
13. 13		84.1	+ 1.03	1 1	15. 7		2.4	— 0.33	3 3	21. 40		80.6	+ 0.70	1 1
15. 46		85.3	— 0.43	1 1						12. 1		80.8	+ 0.19	1 1
10. 56		85.9	— 0.29	1 1	15. 47		11.9	+ 0.08	1 1	17. 39		85.4	+ 0.44	1 1
12. 43		85.9	— 0.37	1 1	12. 1		12.6	+ 0.06	1 1	12. 51		86.1	+ 0.34	1 1
10. 38		85.9	— 1.58	1 1	14. 51		15.5	+ 0.58	1 1	14. 42		87.7	+ 0.02	2 2
										13. 57		88.0	— 0.20	3 3
12. 37		91.0	— 0.70	1 1	16. 6		22.0	+ 0.15	1 1	13. 30		90.1	— 1.08	1 1
12. 37		91.0	— 0.46	1 1						11. 32		90.3	— 1.18	1 1
18. 16		92.9	— 0.47	1 1	12. 11		32.5	— 1.44	1 1	22. 1		90.8	+ 1.32	1 1
12. 34		97.5	— 0.27	1 1						12. 32		95.3	+ 0.43	1 1
14. 1		98.5	+ 0.64	1 1						21. 26		96.0	+ 0.16	1 1
					14. 13		43.5	+ 0.71	1 1	15. 12		99.1	+ 0.33	1 1
13. 20		100.7	+ 0.38	1 1	14. 35		45.2	+ 0.98	1 1	14. 8		99.9	+ 0.17	2 2
9. 36		103.9	— 1.25	1 1	13. 11		48.7	+ 1.36	1 1	14. 18		101.3	— 0.14	3 3
11. 15		104.3	+ 1.34	1 1	14. 58		49.3	+ 0.51	1 1	14. 49		101.5	+ 0.14	2 2
7. 41		104.3	— 0.79	1 1						15. 17		104.8	+ 0.89	1 1
15. 30		104.5	— 0.82	1 1	15. 21		52.3	— 0.24	1 1	13. 40		105.7	— 0.43	1 1
15. 17		104.8	+ 0.74	1 1	16. 6		53.3	— 0.19	1 1	15. 23		106.4	+ 0.66	2 2
10. 45		105.7	— 1.74	2 2	13. 1		53.7	+ 0.39	1 1	21. 41		106.6	+ 0.94	1 1
13. 40		105.7	— 0.32	1 1	10. 48		55.3	— 0.58	1 1	21. 0		107.6	+ 0.79	1 1
12. 25		106.0	+ 0.94	1 1	14. 27		59.3	+ 0.19	1 1					
10. 21		106.4	+ 2.12	1 1	14. 31		59.9	— 0.36	2 2	12. 5		112.1	+ 0.34	1 1
10. 55		107.8	— 1.32	2 2						19. 9		115.4	+ 0.16	1 1
11. 40		107.8	+ 0.28	1 1										
					13. 8		61.7	— 0.77	1 1	Position 0°, r. July, August.				
7. 45		114.6	+ 0.04	3 3	13. 57		62.2	— 0.36	1 1	8. 23		— 29.0	— 0.26	1 1
18. 22		115.5	— 1.77	1 1	12. 21		62.2	— 0.50	1 1	6. 43		— 21.0	+ 0.44	1 1
Position 0°, d. May, June.					15. 54		62.9	— 0.77	1 1	9. 50		— 16.7	+ 0.52	1 1
2. 44		— 34.5	— 1.06	1 1	15. 30		63.0	— 0.21	1 1	6. 57		— 2.8	+ 0.22	1 1
0. 4		— 31.4	— 0.12	1 1	15. 39		63.4	+ 0.45	1 1	8. 3		— 1.1	+ 0.04	1 1
0. 51		— 29.8	+ 0.23	1 1	10. 51		64.8	— 0.54	1 1					
0. 44		— 26.3	+ 0.08	1 1	15. 3		64.8	+ 0.67	1 1	19. 13		1.0	— 0.02	1 1
2. 21		— 23.0	— 0.57	1 1						23. 27		3.2	— 1.17	1 1
2. 37		— 22.6	— 0.08	2 2	14. 11		70.3	+ 0.60	2 2	21. 18		3.4	+ 1.44	1 1
1. 19		— 22.4	— 1.07	1 1	14. 22		70.4	+ 0.59	1 1	18. 3		3.4	+ 1.13	3 3
					14. 47		70.5	+ 0.86	1 1	20. 33		15.4	— 0.17	1 1
3. 40		— 19.0	— 0.81	1 1	12. 30		71.1	— 0.66	1 1	23. 50		16.1	— 1.13	1 1
0. 40		— 15.5	— 0.88	1 1	13. 50		71.1	— 0.62	1 1					
1. 4		— 10.8	— 0.49	1 1	18. 55		75.1	+ 0.99	1 1					
					17. 10		75.5	+ 0.49	1 1					
					10. 38		76.3	— 0.54	1 1					

ALTAZIMUTH TABLES, 1908.

cxxi

Comparison of North Polar Distances observed with the Altazimuth in four different positions,
in the Year 1908, with those of Newcomb's Fundamental Catalogue—contd.

Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight	Approximate		Newcomb N.P.D. - Observed N.P.D.	No. of Obs.	Weight
R.A.	N.P.D.				R.A.	N.P.D.				R.A.	N.P.D.			
Position 0°, r. July, August—contd.					Position 0°, r. July, August—contd.					Position 180°, r. Sept., Oct.—contd.				
h	m	°	"		h	m	°	"		h	m	°	"	
19. 13		22.5	- 0.47	1	18. 30		98.3	+ 0.46	2	19. 13		1.0	- 0.20	1
19. 9		24.2	+ 0.64	1	18. 37		99.1	+ 0.60	1	6. 57		2.8	- 0.44	1
16. 23		28.3	- 0.37	1	17. 54		99.8	+ 2.32	1	23. 27		3.2	- 0.33	4
20. 43		28.5	- 1.54	1						21. 18		3.4	- 0.61	3
16. 0		31.2	+ 0.28	1	21. 5		101.8	+ 1.09	1	18. 3		3.4	+ 0.99	1
18. 45		37.1	- 1.51	1	17. 5		105.6	- 0.48	1	20. 52		9.8	+ 0.25	1
20. 54		49.2	+ 0.73	1	19. 35		106.5	- 0.56	1					
19. 16		49.8	- 1.84	1	23. 59		107.9	+ 0.89	1	22. 41		14.3	- 0.96	1
20. 19		50.0	- 2.62	1	20. 23		108.1	- 2.08	1	18. 49		14.7	+ 1.71	1
18. 33		51.3	+ 0.10	1	23. 5		111.7	- 1.10	1	21. 52		16.7	+ 0.88	1
19. 4		54.1	+ 0.57	1	17. 16		114.9	+ 1.40	1	21. 58		17.3	+ 0.85	5
19. 28		55.8	- 2.58	1	19. 31		115.1	+ 4.41	1	21. 27		19.9	- 2.05	1
22. 5		57.3	- 0.41	1	16. 16		115.4	+ 0.74	1					
21. 9		60.2	+ 0.60	1	18. 22		115.5	+ 0.68	2	1. 19		22.4	- 0.29	1
19. 27		62.2	+ 0.34	1	15. 53		115.9	+ 0.11	1	19. 13		22.5	+ 0.32	1
20. 51		62.3	- 1.34	2	18. 40		117.1	+ 2.61	1	23. 44		22.7	+ 0.18	2
19. 57		62.5	+ 0.04	1	19. 57		118.0	- 1.36	1	22. 46		24.3	- 0.02	2
14. 41		62.5	+ 0.65	1	22. 53		120.1	+ 0.75	1	0. 44		26.3	- 0.64	1
17. 51		63.9	+ 2.46	1	18. 0		120.4	+ 2.31	2	22. 2		27.7	+ 2.45	1
22. 2		65.1	+ 1.68	1	Position 180°, r. September, October.					20. 10		28.2	+ 0.46	1
19. 24		65.5	- 1.41	1	8. 57		- 42.5	(- 5.36)	1	23. 21		28.2	- 0.86	1
1. 49		69.6	+ 0.30	1	8. 53		- 41.6	(- 2.68)	1	20. 43		28.5	- 0.24	2
15. 42		74.3	+ 0.53	1	9. 27		- 37.9	+ 0.69	3	0. 51		29.8	- 0.46	1
18. 55		75.1	+ 1.33	1	11. 49		- 35.8	+ 1.42	2	22. 26		32.1	+ 0.91	1
0. 8		75.3	- 0.08	1	8. 57		- 35.4	- 3.70	1	19. 35		40.0	- 0.05	1
20. 31		75.7	- 1.19	1	12. 50		- 33.5	- 1.61	1	21. 43		41.1	- 0.49	2
16. 21		75.8	+ 1.85	1	7. 35		- 31.1	+ 0.46	3	1. 32		41.8	+ 0.94	1
17. 31		77.4	+ 1.62	1	9. 44		- 30.5	- 0.36	1	21. 3		42.7	+ 0.33	1
19. 46		81.4	- 1.08	1	8. 23		- 29.0	- 1.09	1	21. 31		44.8	- 0.60	1
19. 29		82.8	- 0.60	2	10. 18		- 24.0	+ 0.34	2	20. 38		45.1	+ 1.23	1
1. 36		85.0	+ 0.76	1	9. 2		- 22.5	- 1.49	1	19. 42		45.1	+ 0.67	2
17. 39		85.4	+ 3.10	1	6. 43		- 21.0	+ 0.16	1	23. 36		46.2	+ 0.40	1
19. 9		87.9	- 1.87	1	10. 36		- 20.5	- 0.71	2	22. 57		48.2	+ 1.09	1
22. 21		89.1	+ 0.37	1	9. 34		- 20.4	+ 0.13	2	22. 40		48.7	- 1.12	1
23. 22		89.3	- 3.26	1	14. 28		- 13.9	+ 0.14	1	20. 54		49.2	+ 1.20	1
20. 7		191.1	- 1.52	1	10. 27		- 13.8	- 0.66	3	20. 19		50.0	- 0.19	1
0. 48		91.7	- 0.89	1	12. 1		- 12.6	+ 0.78	2	18. 47		56.8	- 0.20	2
18. 24		92.1	+ 0.47	1	10. 53		- 11.7	- 0.92	1	0. 32		56.8	- 0.35	1
18. 16		92.9	+ 0.53	2	6. 30		- 10.3	- 0.44	1	22. 5		57.3	- 0.11	3
18. 42		94.9	- 2.04	1	6. 57		- 2.8	+ 0.55	1	22. 5		57.3	+ 0.26	1
21. 26		96.0	+ 1.06	1	12. 14		- 1.8	+ 0.15	2	0. 34		59.7	- 0.65	1
22. 11		98.3	- 0.67	1	8. 3		- 1.1	+ 0.53	6	21. 9		60.2	+ 0.48	3
									4	21. 46		60.3	- 0.04	1
										20. 51		62.3	+ 0.15	1

GREENWICH OBSERVATIONS, 1908.

q

cxxxii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Comparison of North Polar Distances observed with the Altazimuth in four different positions, in the Year 1908, with those of Newcomb's Fundamental Catalogue—contd.

Approximate		Newcomb N.P.D. —Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. —Observed N.P.D.	No. of Obs.	Weight.	Approximate		Newcomb N.P.D. —Observed N.P.D.	No. of Obs.	Weight.							
R.A.	N.P.D.				R.A.	N.P.D.				R.A.	N.P.D.										
Position 180°, r. Sept., Oct.—contd.																					
h	m	°	"			h	m	°	"			h	m	°	"						
21.	48	64.5	— 0.61	4	3	22.	33	94.7	+ 0.30	1	1	22.	39	60.3	— 0.02	1	1				
21.	40	64.8	— 1.27	1	1	23.	57	96.5	— 0.95	1	1	6.	38	64.8	— 0.54	1	1				
19.	24	65.5	— 0.15	1	1	2.	12	96.8	+ 0.73	1	1	3.	42	66.2	+ 0.06	1	1				
4.	37	67.2	— 0.64	1	1	4.	7	97.1	— 1.66	1	1	2.	2	67.0	— 1.23	1	1				
7.	15	67.9	+ 0.15	1	1	21.	33	98.3	+ 0.80	1	1	4.	58	68.5	— 0.36	1	1				
4.	58	68.5	— 0.09	1	1	22.	15	98.3	+ 0.46	2	2	3.	16	69.2	— 0.07	1	1				
2.	54	69.0	— 0.06	1	1	20.	48	99.3	+ 1.54	1	1	4.	23	71.0	+ 0.45	1	1				
3.	15	69.2	+ 0.61	1	1	20.	43	99.8	— 1.27	3	3	8.	18	71.4	— 1.44	1	1				
Position 180°, r. Sept., Oct.—contd.																					
19.	55	70.8	+ 0.15	1	1	20.	52	100.1	— 0.18	1	1	23.	48	71.4	— 0.30	1	1				
4.	46	71.3	— 0.97	1	1	23.	14	100.1	+ 0.46	1	1	2.	25	72.7	+ 0.24	1	1				
7.	8	73.7	— 1.17	1	1	20.	13	102.8	— 0.40	1	1	7.	13	73.3	+ 0.78	1	1				
19.	33	73.8	+ 0.80	1	1	20.	13	102.8	— 0.24	1	1	4.	14	74.6	+ 0.49	1	1				
4.	15	74.6	— 1.51	1	1	21.	48	104.0	+ 2.36	1	1	2.	46	75.3	+ 0.10	1	1				
20.	39	75.3	+ 0.46	1	1	19.	35	106.5	+ 0.83	1	1	1.	0	75.6	— 0.88	1	1				
1.	0	75.6	— 1.45	1	1	21.	42	106.6	— 0.59	2	2	20.	33	75.7	— 0.56	1	1				
20.	33	75.7	— 1.76	1	1	21.	35	107.1	— 0.87	1	1	3.	22	80.6	— 0.20	1	1				
6.	40	77.0	— 1.70	1	1	21.	1	107.6	+ 2.24	1	1	5.	50	82.6	— 0.16	1	1				
20.	29	79.0	+ 0.58	2	2	21.	38	109.3	— 0.29	1	1	0.	44	82.9	— 1.91	1	1				
1.	46	79.4	— 0.06	1	1	2.	47	111.4	— 1.47	1	1	23.	55	83.7	— 0.92	1	1				
2.	38	79.7	— 0.09	1	1	23.	5	111.7	+ 0.75	1	1	1.	25	84.3	— 1.05	1	1				
22.	36	79.7	— 0.40	1	1	20.	13	112.1	+ 0.58	2	2	7.	34	84.5	+ 0.67	1	1				
Position 180°, d. Nov., Dec.—contd.																					
21.	6	80.2	— 0.93	1	1	21.	23	112.2	+ 1.35	1	1	23.	35	84.9	— 0.39	1	1				
21.	40	80.6	+ 0.74	2	2	2.	8	114.2	+ 0.62	1	1	8.	34	86.3	— 2.37	1	1				
23.	2	81.1	— 0.32	1	1	19.	10	115.4	— 0.25	1	1	1.	13	86.9	— 0.35	1	1				
19.	46	81.4	+ 0.80	2	2	18.	50	116.4	— 0.91	1	1	23.	42	87.0	— 1.02	1	1				
2.	8	81.6	+ 2.09	1	1	22.	53	120.1	+ 0.39	2	2	0.	21	88.4	— 0.57	1	1				
23.	7	81.8	— 0.06	1	1	2.	9	121.2	+ 1.08	1	1	5.	27	90.4	+ 0.32	1	1				
5.	50	82.6	— 0.01	2	2	Position 180°, d. Nov., Dec.										22.	31	90.6	+ 0.74	1	1
19.	30	82.8	+ 0.09	1	1	16.	44	— 33.1	+ 0.15	1	1	0.	48	91.7	— 0.87	1	1				
19.	51	83.8	— 0.46	1	1	16.	0	— 31.2	— 0.64	1	1	5.	36	92.0	— 0.75	1	1				
22.	6	84.3	— 0.09	2	2	16.	6	— 22.0	— 1.89	1	1	5.	20	92.5	— 0.63	1	1				
1.	25	84.3	— 0.67	1	1	18.	3	— 3.4	— 0.89	1	1	0.	30	94.1	— 1.81	1	1				
1.	37	85.0	— 0.32	1	1	15.	7	— 2.4	— 0.08	1	1	4.	7	97.1	— 0.79	1	1				
21.	11	85.1	— 1.09	2	2	8.	4	1.1	— 0.58	1	1	1.	19	98.7	— 1.58	1	1				
22.	59	86.7	— 0.09	1	1	1.	26	1.2	— 0.50	1	1	5.	43	99.7	— 0.06	1	1				
1.	13	86.9	— 0.67	1	1	23.	5	15.1	— 0.12	1	1	3.	29	99.8	— 0.95	1	1				
19.	21	87.1	— 0.34	1	1	3.	34	27.1	+ 0.05	1	1	1.	47	100.8	— 0.72	1	1				
23.	12	87.2	+ 0.09	1	1	3.	52	50.3	— 0.61	1	1	3.	53	103.8	— 0.86	1	1				
21.	56	89.8	+ 0.77	1	1	2.	4	55.5	+ 0.48	1	1	23.	59	107.9	+ 0.30	1	1				
22.	24	90.5	+ 0.67	1	1	2.	12	56.6	— 0.77	1	1	0.	39	108.5	— 1.36	1	1				
22.	31	90.6	— 1.05	1	1	Position 180°, d. Nov., Dec.—contd.										23.	18	110.6	— 0.16	1	1
22.	1	90.8	— 0.08	1	1	3.	52	50.3	— 0.61	1	1	8.	4	114.0	— 0.54	1	1				
18.	24	92.1	+ 0.58	1	1	2.	4	55.5	+ 0.48	1	1	7.	45	114.6	— 2.31	1	1				
18.	17	92.9	— 0.26	2	2	2.	12	56.6	— 0.77	1	1	0.	54	119.9	— 3.04	1	1				

ALTAZIMUTH TABLES, 1908.

cxxxiii

The results obtained in the above tables were divided into groups for each 10° of N.P.D., the divisions of the groups being indicated by the spaces. They were combined according to the following system of weights:—

No. of Observations.	Weight.	No. of Observations.	Weight.
1	1	5 or 6	4
2	2	7, 8 or 9	5
3 or 4	3	10 to 14	6

*Comparison of North Polar Distances observed with the Altazimuth in 1908,
with Newcomb's Fundamental Catalogue.*

Mean N.P.D. of Group.	Mean Tabular —Observed.	Weight.	Mean N.P.D. of Group.	Mean Tabular —Observed.	Weight.	Mean N.P.D. of Group.	Mean Tabular —Observed.	Weight.
Position 180° , r. Jan., Feb.			Position 0° , d. May, June.			Position 180° , r. Sept., Oct.		
°	"		°	"		°	"	
— 37'4	+ 0'21	2	— 32'9	— 0'59	2	— 34'4	— 0'07	11
— 24'8	+ 0'01	10	— 24'4	— 0'25	6	— 23'0	— 0'45	7
— 14'6	— 0'32	5	— 15'1	— 0'73	3	— 14'3	— 0'05	10
— 2'7	+ 0'06	12	— 1'7	— 0'51	3	— 1'4	+ 0'45	9
2'5	— 0'19	5	2'2	— 0'93	5	3'6	— 0'23	11
19'0	+ 0'25	1	13'3	+ 0'24	3	16'9	+ 0'43	9
25'7	— 0'12	7	22'0	+ 0'15	1	25'9	+ 0'06	13
33'0	+ 0'40	3	32'5	— 1'44	1	36'0	+ 0'43	2
43'4	— 0'12	7	46'7	+ 0'89	4	44'9	+ 0'29	11
56'2	— 0'71	14	56'7	— 0'16	7	56'6	— 0'18	9
66'4	— 0'05	30	63'2	— 0'24	8	64'6	— 0'19	16
75'0	— 0'15	22	73'3	+ 0'24	11	76'0	— 0'47	14
82'8	— 0'43	23	85'3	+ 0'08	10	83'9	— 0'08	22
94'6	— 0'79	23	95'3	+ 0'04	8	95'9	— 0'07	17
105'5	— 0'07	12	104'0	+ 0'31	11	104'9	+ 0'25	11
112'4	+ 0'19	9	113'8	+ 0'25	2	115'2	+ 0'28	11
Position 180° , d. March, April.			Position 0° , r. July, August.			Position 180° , d. Nov., Dec.		
— 41'1	— 2'80	1	— 25'0	+ 0'09	2	— 32'1	— 0'25	2
— 26'4	— 0'42	2	— 16'7	+ 0'52	1	— 22'0	— 1'89	1
— 14'9	— 0'34	2	— 1'9	+ 0'13	2	— 2'9	— 0'49	2
— 2'6	+ 0'02	6	3'0	+ 0'61	6	1'1	— 0'54	2
1'7	+ 0'42	6	15'7	— 0'65	2	15'1	— 0'12	1
13'8	+ 0'04	1	25'9	— 0'44	4	27'1	+ 0'05	1
37'3	+ 0'54	3	35'0	— 0'62	2	54'1	— 0'30	3
47'9	— 0'58	4	49'5	— 0'56	2	66'0	— 0'36	6
55'8	— 0'33	10	53'2	— 0'59	4	73'6	— 0'12	9
63'8	— 0'30	24	63'6	+ 0'20	10	84'7	— 0'75	11
75'0	— 0'30	11	75'6	+ 0'68	6	94'6	— 0'64	10
105'4	+ 0'18	14	47'0	— 0'40	8	105'2	— 0'66	4
114'7	— 0'41	4	95'2	+ 0'08	11	114'7	— 1'51	4
			106'0	— 0'23	5			
			116'8	+ 1'17	11			

cxxxiv INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Assuming the differences "Tabular—Observed" to be represented by the formula

$$a + b \sin Z.D.,$$

a solution by the method of least squares gave the results:—

Newcomb — Altazimuth.

Position of Altazimuth.

Position of Altazimuth.

180°, r (Jan., Feb.).	—0'16	—0'23 sin Z.D.	0°, r (July, Aug.).	+0'09	+0'16 sin Z.D.
180°, d (Mar., Apr.).	—0'06	—0'10 sin Z.D.	180°, r (Sept., Oct.).	—0'01	—0'07 sin Z.D.
0°, d (May, June).	—0'17	+0'47 sin Z.D.	180°, d (Nov., Dec.).	—0'52	—0'04 sin Z.D.

Taking the coefficient of $\sin Z.D.$ from the results of the reflexion observations as $-0''\cdot20$, the values found for the α term in the different positions are given below.

Position of Instrument.	Difference of N.P.D. Newcomb — Altazimuth.	Weight.	Position of Instrument.	Difference of N.P.D. Newcomb — Altazimuth.	Weight.
Value of α .			Value of α .		
180°, r (Jan., Feb.).	—0'18	185	0°, r (July, Aug.).	+0'24	76
180°, d (Mar., Apr.).	—0'10	114	180°, r (Sept., Oct.).	+0'01	183
0°, d (May, June).	+0'02	85	180°, d (Nov., Dec.).	—0'44	55

The weighted mean is $\mp 0''\cdot06 - 0''\cdot20 \sin Z.D.$

Adopted Corrections to North Polar Distances of the Altazimuth, 1908.

N.P.D.	Corrections to Results of Direct Observations.		Corrections to Results of Reflexion Observations.	
	0°, d. 180°, r.	0°, r. 180°, d.	0°, d. 180°, r.	0°, r. 180°, d.
— 45	+ 0'14	+ 0'26	— 0'25	— 0'37
— 40	+ 0'14	+ 0'26	— 0'25	— 0'37
— 30	+ 0'13	+ 0'25	— 0'24	— 0'36
— 20	+ 0'11	+ 0'23	— 0'21	— 0'33
— 10	+ 0'09	+ 0'21	— 0'18	— 0'30
0	+ 0'07	+ 0'19	— 0'14	— 0'26
+ 10	+ 0'04	+ 0'16	— 0'09	— 0'21
20	0'00	+ 0'12	— 0'04	— 0'16
30	— 0'03	+ 0'09	+ 0'01	— 0'11
40	— 0'07	+ 0'05	+ 0'07	— 0'05
50	— 0'10	+ 0'02	+ 0'13	+ 0'01
60	— 0'14	— 0'02	+ 0'18	+ 0'06
70	— 0'17	— 0'05	+ 0'23	+ 0'11
80	— 0'20	— 0'08	+ 0'28	+ 0'16
90	— 0'22	— 0'10	+ 0'31	+ 0'19
100	— 0'24	— 0'12	+ 0'34	+ 0'22
110	— 0'25	— 0'13	+ 0'37	+ 0'25
120	— 0'26	— 0'14	+ 0'38	+ 0'26
125	— 0'26	— 0'14	+ 0'38	+ 0'26

ALTAZIMUTH TABLES, 1908.

CXXV

*Altazimuth Colatitude Investigation.*Correction to Assumed Colatitude $38^{\circ} 31' 23''.42$ from observations in the Year 1908.(Using Pulkowa Refractions, computed from the tables given in the Appendix to the *Greenwich Observations for 1898.*)

Corrections for Variation of Latitude supplied by Prof. Albrecht have been applied in forming this table (see p. {20} of the section "Ledgers of Mean Right Ascensions and North Polar Distances of Stars observed with the Altazimuth"):

Star's Name.	Approximate R.A.	Approximate N.P.D.	Excess of N.P.D. above Pole.	No. of Observations		Weight	Star's Name.	Approximate R.A.	Approximate N.P.D.	Excess of N.P.D. above Pole.	No. of Observations		Weight
				Above Pole.	Below Pole.						Above Pole.	Below Pole.	
λ Ursæ Minoris..	h m 19 13	° ' ° 1 0	- 0".77	2	1	4	16 Cephei.....	h m 21 58	° ' ° 17 15	+ 0".66	6	1	4
Groomb. 1119...	8 4	1 5	- 0".60	4	7	12	Lalande 6061...	3 7	18 7	+ 0".05	2	1	3
Polaris.....	1 26	1 11	+ 0".72	2	4	7	γ Camelopardi..	3 41	18 57	+ 1".03	2	2	5
Bradley 1672....	12 14	1 47	+ 1".03	4	4	10	Piazzi X. 126...	10 36	20 27	+ 0".01	2	2	5
Groomb. 2283...	15 8	2 25	+ 0".44	3	5	10	Groomb. 2320...	16 6	21 57	+ 0".94	1	2	4
Cephei 51.....	6 58	2 48	+ 0".32	5	3	10	ψ Cassiopeiæ....	1 19	22 21	- 0".19	1	2	4
Bradley 3147...	23 28	3 12	+ 0".36	5	2	8	δ Draconis.....	19 13	22 30	+ 0".61	2	1	3
Groomb. 3548...	21 18	3 21	+ 1".06	5	2	8	Bradley 366.....	2 37	22 34	+ 0".36	1	2	4
δ Ursæ Minoris..	18 4	3 23	- 0".09	5	5	12	ϵ Cassiopeiæ....	2 21	23 1	- 0".19	1	1	2
Piazzi VII. 132.	7 41	9 30	- 0".25	2	1	4	Bradley 1429...	10 18	23 58	+ 0".06	2	2	5
B. D. + 80° 395	12 43	9 34	+ 0".68	2	2	6	ζ Draconis.....	19 9	24 11	- 2".49	1	1	2
Lalande F.(2) 160	13 11	9 52	- 0".45	2	2	6	ϵ Cephei.....	22 46	24 17	+ 0".05	2	1	3
Piazzi VII. 187.	7 50	10 16	- 0".76	2	1	3	Bradley 82.....	0 44	26 15	- 0".01	1	2	3
ζ Ursæ Minoris..	15 47	11 55	- 0".55	2	2	5	68 Draconis.....	20 10	28 12	- 0".37	1	1	2
Groomb. 1852...	12 1	12 35	- 0".93	2	2	5	η Cephei.....	20 43	28 31	+ 0".48	3	2	5
Bradley 1446....	10 27	13 49	+ 0".81	2	4	7	γ Cassiopeiæ....	0 51	29 47	+ 0".92	1	2	3
Bradley 473.....	3 28	14 34	+ 0".16	2	2	5	ϵ Lyncis.....	7 35	31 4	- 0".56	1	3	4
50 Draconis.....	18 49	14 40	+ 0".02	1	1	3	θ Draconis.....	16 0	31 11	+ 0".70	1	1	2
Groomb. 379....	1 44	14 52	+ 0".62	2	1	3	η Persei.....	2 44	34 29	+ 0".85	1	1	2
β Ursæ Minoris.	14 51	15 28	+ 0".16	1	1	3	γ Ursæ Majoris.	11 49	35 48	- 3".14	1	2	3
21 Cassiopeiæ....	0 40	15 31	+ 0".61	1	1	3	Groomb. 2671...	18 45	37 7	+ 1".56	1	1	2
Groomb. 1865...	12 11	16 56	+ 0".29	2	2	5	θ Ursæ Majoris.	9 27	37 54	- 0".02	1	3	4
Groomb. 1998...	13 23	17 15	+ 1".48	2	1	3	π^2 Cygni.....	21 43	41 7	(+ 2".08)	2	2	3

The weights used in the above tables are the same as those used in the Transit-Circle investigation.

The assumed colatitude $38^{\circ} 31' 23''.42$ was found by applying to the adopted colatitude of the Transit-Circle, with Pulkowa Refractions $38^{\circ} 31' 21''.75$, a correction of $+1''.67$ for geodetic difference of latitude as measured on the plan of the Observatory.

The resulting correction to the assumed colatitude $38^{\circ} 31' 23''.42$ from these results is $-0''.178$. The colatitude determined from the observations of 1908 is, therefore, $38^{\circ} 31' 23''.24$.

cxxxvi INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 70°.									N.P.D.
	61°.	62°.	63°.	64°.	65°.	66°.	67°.	68°.	69°.	
0	9°7490	9°9165	10°0773	10°2318	10°3799	10°5216	10°6572	10°7867	10°9102	0
1	7519	9192	0799	2343	3823	5239	6594	7888	9122	1
2	7547	9220	0825	2368	3847	5262	6616	7909	9142	2
3	7575	9247	0851	2393	3871	5285	6638	7930	9162	3
4	7603	9274	0878	2418	3895	5308	6660	7951	9182	4
5	7631	9301	0904	2443	3919	5331	6682	7972	9202	5
6	7659	9328	0930	2469	3943	5354	6704	7993	9222	6
7	7688	9355	0956	2494	3967	5377	6726	8014	9242	7
8	7717	9382	0983	2519	3991	5400	6748	8035	9262	8
9	7745	9409	1009	2544	4015	5423	6770	8056	9282	9
10	7773	9436	1035	2569	4039	5446	6792	8077	9302	10
11	7801	9463	1061	2594	4063	5469	6814	8098	9322	11
12	7830	9491	1087	2619	4087	5492	6836	8119	9342	12
13	7858	9519	1113	2644	4111	5515	6857	8139	9361	13
14	7886	9547	1139	2669	4135	5538	6879	8160	9381	14
15	7914	9574	1165	2694	4159	5561	6901	8181	9401	15
16	7943	9601	1191	2719	4183	5584	6923	8202	9421	16
17	7971	9628	1217	2744	4207	5607	6945	8223	9441	17
18	7999	9655	1243	2769	4230	5629	6966	8243	9460	18
19	8027	9682	1269	2794	4254	5652	6988	8264	9480	19
20	8055	9709	1295	2818	4278	5675	7010	8285	9500	20
21	8083	9736	1321	2843	4302	5698	7032	8306	9520	21
22	8111	9763	1347	2868	4326	5721	7054	8327	9539	22
23	8139	9790	1373	2893	4350	5743	7075	8347	9559	23
24	8167	9816	1399	2918	4373	5766	7097	8368	9578	24
25	8195	9843	1424	2943	4397	5789	7119	8389	9598	25
26	8223	9870	1450	2968	4421	5812	7141	8410	9618	26
27	8251	9897	1475	2993	4445	5834	7162	8430	9637	27
28	8279	9924	1501	3017	4468	5857	7184	8451	9657	28
29	8307	9951	1527	3042	4492	5879	7205	8471	9676	29
30	8335	9978	1553	3067	4516	5902	7227	8492	9696	30
31	8363	10°0005	1579	3092	4540	5925	7249	8513	9716	31
32	8391	0031	1605	3116	4563	5947	7270	8533	9735	32
33	8419	0058	1630	3141	4587	5970	7292	8554	9755	33
34	8447	0085	1656	3165	4610	5992	7313	8574	9774	34
35	8474	0112	1682	3189	4634	6015	7335	8595	9794	35
36	8502	0138	1708	3214	4657	6037	7356	8615	9813	36
37	8529	0164	1734	3239	4681	6060	7378	8636	9833	37
38	8557	0191	1759	3264	4704	6082	7399	8656	9852	38
39	8585	0217	1784	3289	4728	6105	7421	8677	9872	39
40	8613	0244	1810	3313	4751	6127	7442	8697	9891	40
41	8641	0270	1836	3338	4775	6149	7463	8717	9910	41
42	8669	0297	1862	3362	4798	6172	7485	8738	9930	42
43	8696	0324	1888	3386	4822	6194	7506	8758	9949	43
44	8724	0351	1914	3410	4845	6217	7528	8779	9969	44
45	8751	0377	1939	3435	4869	6239	7549	8799	10°9988	45
46	9°8779	10°0404	10°1965	10°3459	10°4892	10°6261	10°7570	10°8819	11°0007	46

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second—continued.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 70°.									N.P.D.
	61°.	62°.	63°.	64°.	65°.	66°.	67°.	68°.	69°.	
47	9"8807	10"0430	10"1990	10"3484	10"4916	10"6283	10"7591	10"8839	11"0026	47
48	8835	0457	2014	3508	4938	6306	7613	8860	0046	48
49	8862	0483	2039	3533	4961	6328	7634	8880	0065	49
50	8890	0510	2065	3557	4984	6350	7655	8900	0084	50
51	8917	0536	2090	3581	5008	6372	7676	8920	0103	51
52	8945	0563	2116	3605	5031	6394	7697	8940	0122	52
53	8972	0589	2141	3630	5054	6417	7719	8961	0142	53
54	9000	0615	2167	3654	5077	6439	7740	8981	0161	54
55	9027	0642	2192	3679	5101	6461	7761	9001	0180	55
56	9055	0668	2217	3703	5124	6483	7782	9021	0199	56
57	9082	0694	2243	3727	5147	6505	7803	9041	0218	57
58	9110	0720	2268	3751	5170	6528	7825	9062	0238	58
59	9137	0746	2293	3775	5193	6550	7846	9082	0257	59
60	9165	10"0773	10"2318	10"3799	10"5216	10"6572	10"7867	10"9102	11"0276	60

N.P.D.	Azimuth 70°.									N.P.D.
	70°.	71°.	72°.	73°.	74°.	75°.	76°.	77°.	78°.	
0	11"0276	11"1391	11"2447	11"3444	11"4382	11"5263	11"6086	11"6851	11"7559	0
1	0295	1409	2464	3460	4397	5277	6099	6863	7570	1
2	0314	1427	2481	3476	4412	5291	6112	6875	7581	2
3	0333	1445	2498	3492	4428	5306	6126	6888	7593	3
4	0352	1463	2515	3508	4443	5320	6139	6900	7604	4
5	0371	1481	2532	3524	4458	5334	6152	6912	7615	5
6	0390	1499	2549	3540	4473	5348	6165	6924	7626	6
7	0409	1517	2566	3556	4488	5362	6178	6936	7637	7
8	0428	1535	2583	3572	4503	5376	6192	6949	7649	8
9	0447	1553	2600	3588	4518	5390	6205	6961	7660	9
10	0466	1571	2617	3604	4533	5404	6218	6973	7671	10
11	0485	1589	2634	3620	4548	5418	6231	6985	7682	11
12	0504	1607	2651	3636	4563	5432	6244	6997	7693	12
13	0522	1624	2668	3652	4578	5446	6257	7009	7705	13
14	0541	1642	2685	3668	4593	5460	6270	7021	7716	14
15	0560	1660	2702	3684	4608	5474	6283	7033	7727	15
16	0579	1678	2719	3700	4623	5488	6296	7045	7738	16
17	0598	1696	2736	3716	4638	5502	6309	7057	7749	17
18	0616	1713	2752	3731	4652	5516	6322	7069	7760	18
19	0635	1731	2769	3747	4667	5530	6335	7081	7771	19
20	0654	1749	2786	3763	4682	5544	6348	7093	7782	20
21	0673	1767	2803	3779	4697	5558	6361	7105	7793	21
22	0692	1785	2820	3795	4712	5572	6374	7117	7804	22
23	0710	1802	2836	3810	4726	5585	6386	7129	7815	23
24	11"0729	11"1820	11"2853	11"3826	11"4741	11"5599	11"6399	11"7141	11"7826	24

cxxxviii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second—continued.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 70°.									N.P.D.
	70°.	71°.	72°.	73°.	74°.	75°.	76°.	77°.	78°.	
25	11°0748	11°1838	11°2870	11°3842	11°4756	11°5613	11°6412	11°7153	11°7837	25
26	0767	1856	2887	3858	4771	5627	6425	7165	7848	26
27	0785	1873	2903	3873	4786	5641	6438	7177	7859	27
28	0804	1891	2920	3889	4800	5654	6450	7188	7870	28
29	0822	1908	2936	3904	4815	5668	6463	7200	7881	29
30	0841	1926	2953	3920	4830	5682	6476	7212	7892	30
31	0860	1944	2970	3936	4845	5696	6489	7224	7903	31
32	0878	1961	2986	3951	4859	5709	6501	7236	7914	32
33	0897	1979	3003	3967	4874	5723	6514	7247	7924	33
34	0915	1996	3019	3982	4888	5736	6526	7259	7935	34
35	0934	2014	3036	3998	4903	5750	6539	7271	7946	35
36	0952	2031	3052	4014	4918	5764	6552	7283	7957	36
37	0971	2049	3069	4029	4932	5777	6564	7294	7968	37
38	0989	2066	3085	4045	4947	5791	6577	7306	7978	38
39	1008	2084	3102	4060	4961	5804	6589	7317	7989	39
40	1026	2101	3118	4076	4976	5818	6602	7329	8000	40
41	1044	2118	3134	4091	4990	5832	6615	7341	8011	41
42	1063	2136	3151	4107	5005	5845	6627	7352	8021	42
43	1081	2153	3167	4122	5019	5859	6640	7364	8032	43
44	1100	2171	3184	4138	5034	5872	6652	7375	8042	44
45	1118	2188	3200	4153	5048	5886	6665	7387	8053	45
46	1136	2205	3216	4168	5062	5899	6677	7399	8064	46
47	1154	2223	3233	4184	5077	5913	6690	7410	8074	47
48	1173	2240	3249	4199	5091	5926	6702	7422	8085	48
49	1191	2258	3266	4215	5106	5940	6715	7433	8095	49
50	1209	2275	3282	4230	5120	5953	6727	7445	8106	50
51	1227	2292	3298	4245	5134	5966	6739	7456	8116	51
52	1245	2309	3314	4260	5149	5980	6752	7468	8127	52
53	1264	2327	3331	4276	5163	5993	6764	7479	8137	53
54	1282	2344	3347	4291	5178	6007	6777	7491	8148	54
55	1300	2361	3363	4306	5192	6020	6789	7502	8158	55
56	1318	2378	3379	4321	5206	6033	6801	7513	8168	56
57	1336	2395	3395	4336	5220	6046	6814	7525	8179	57
58	1355	2413	3412	4352	5235	6060	6826	7536	8189	58
59	1373	2430	3428	4367	5249	6073	6839	7548	8200	59
60	11°1391	11°2447	11°3444	11°4382	11°5263	11°6086	11°6851	11°7559	11°8210	60

ALTAZIMUTH TABLES, 1908.

cxxxix

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second—continued.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 70°.											N.P.D.
	79°.	80°.	81°.	82°.	83°.	84°.	85°.	86°.	87°.	88°.	89°.	
0	11°8210	11°8804	11°9341	11°9821	12°0244	12°0610	12°0920	12°1175	12°1372	12°1513	12°1598	0
1	8220	8813	9349	9828	0251	0616	0925	1179	1375	1515	1599	1
2	8231	8823	9358	9836	0257	0621	0929	1182	1378	1517	1600	2
3	8241	8832	9366	9843	0264	0627	0934	1186	1380	1518	1600	3
4	8252	8842	9375	9851	0270	0632	0938	1189	1383	1520	1601	4
5	8262	8851	9383	9858	0277	0638	0943	1193	1386	1522	1602	5
6	8272	8860	9391	9865	0283	0644	0948	1197	1389	1524	1603	6
7	8282	8870	9400	9873	0290	0649	0952	1200	1392	1526	1604	7
8	8293	8879	9408	9880	0296	0655	0957	1204	1394	1527	1604	8
9	8303	8889	9417	9888	0303	0660	0961	1207	1397	1529	1605	9
10	8313	8898	9425	9895	0309	0666	0966	1211	1400	1531	1606	10
11	8323	8907	9433	9902	0315	0671	0971	1215	1403	1533	1607	11
12	8333	8916	9441	9910	0322	0677	0975	1218	1405	1534	1608	12
13	8344	8926	9450	9917	0328	0682	0980	1222	1408	1536	1608	13
14	8354	8935	9458	9925	0335	0688	0984	1225	1410	1537	1609	14
15	8364	8944	9466	9932	0341	0693	0989	1229	1413	1539	1610	15
16	8374	8953	9474	9939	0347	0698	0993	1232	1416	1541	1611	16
17	8384	8962	9482	9946	0353	0704	0998	1236	1418	1542	1611	17
18	8394	8972	9491	9954	0360	0709	1002	1239	1421	1544	1612	18
19	8404	8981	9499	9961	0366	0715	1007	1243	1423	1545	1612	19
20	8414	8990	9507	9968	0372	0720	1011	1246	1426	1547	1613	20
21	8424	8999	9515	9975	0378	0725	1015	1249	1428	1549	1614	21
22	8434	9008	9523	9982	0384	0730	1020	1253	1431	1550	1614	22
23	8444	9017	9532	9990	0391	0736	1024	1256	1433	1552	1615	23
24	8454	9026	9540	11°9997	0397	0741	1029	1260	1436	1553	1615	24
25	8464	9035	9548	12°0004	0403	0746	1033	1263	1438	1555	1616	25
26	8474	9044	9556	0011	0409	0751	1037	1266	1440	1556	1617	26
27	8484	9053	9564	0018	0415	0756	1041	1270	1443	1558	1617	27
28	8494	9062	9572	0026	0422	0762	1046	1273	1445	1559	1618	28
29	8504	9071	9580	0033	0428	0767	1050	1277	1448	1561	1618	29
30	8514	9080	9588	0040	0434	0772	1054	1280	1450	1562	1619	30
31	8524	9089	9596	0047	0440	0777	1058	1283	1452	1563	1619	31
32	8534	9098	9604	0054	0446	0782	1062	1286	1454	1565	1620	32
33	8543	9106	9612	0061	0452	0788	1067	1290	1457	1566	1620	33
34	8553	9115	9620	0068	0458	0793	1071	1293	1459	1568	1621	34
35	8563	9124	9628	0075	0464	0798	1075	1296	1461	1569	1621	35
36	8573	9133	9636	0082	0470	0803	1079	1299	1463	1570	1621	36
37	8583	9142	9644	0089	0476	0808	1083	1302	1465	1572	1622	37
38	8592	9150	9651	0096	0482	0813	1088	1306	1468	1573	1622	38
39	8602	9159	9659	0103	0488	0818	1092	1309	1470	1575	1623	39
40	8612	9168	9667	0110	0494	0823	1096	1312	1472	1576	1623	40
41	8622	9177	9675	0117	0500	0828	1100	1315	1474	1577	1623	41
42	8632	9186	9683	0124	0506	0833	1104	1318	1476	1578	1623	42
43	8641	9194	9690	0130	0512	0838	1108	1322	1479	1580	1624	43
44	8651	9203	9698	0137	0518	0843	1112	1325	1481	1581	1624	44
45	8661	9212	9706	0144	0524	0848	1116	1328	1483	1582	1624	45
46	11°8671	11°9221	11°9714	12°0151	12°0530	12°0853	12°1120	12°1331	12°1485	12°1583	12°1624	46

GREENWICH OBSERVATIONS, 1908.

r

cxix INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second—continued.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 70°.											N.P.D.
	79°.	80°.	81°.	82°.	83°.	84°.	85°.	86°.	87°.	88°.	89°.	
47	11"8680	11"9229	11"9722	12"0158	12"0536	12"0858	12"1124	12"1334	12"1487	12"1584	12"1624	47
48	'8690	'9238	'9729	'0164	'0541	'0862	'1128	'1337	'1489	'1586	'1625	48
49	'8699	'9246	'9737	'0171	'0547	'0867	'1132	'1340	'1491	'1587	'1625	49
50	'8709	'9255	'9745	'0178	'0553	'0872	'1136	'1343	'1493	'1588	'1625	50
51	'8719	'9264	'9753	'0185	'0559	'0877	'1140	'1346	'1495	'1589	'1625	51
52	'8728	'9272	'9760	'0191	'0565	'0882	'1144	'1349	'1497	'1590	'1625	52
53	'8738	'9281	'9768	'0198	'0570	'0886	'1148	'1352	'1499	'1591	'1626	53
54	'8747	'9289	'9775	'0204	'0576	'0891	'1152	'1356	'1501	'1592	'1626	54
55	'8757	'9298	'9783	'0211	'0582	'0896	'1156	'1358	'1503	'1593	'1626	55
56	'8766	'9307	'9791	'0218	'0588	'0901	'1160	'1361	'1505	'1594	'1626	56
57	'8776	'9315	'9798	'0224	'0593	'0906	'1164	'1364	'1507	'1595	'1626	57
58	'8785	'9324	'9806	'0231	'0599	'0910	'1167	'1366	'1509	'1596	'1626	58
59	'8795	'9332	'9813	'0237	'0604	'0915	'1171	'1369	'1511	'1597	'1626	59
60	11"8804	11"9341	11"9821	12"0244	12"0610	12"0920	12"1175	12"1372	12"1513	12"1598	12"1626	60

ALTAZIMUTH TABLES, 1908.

cxxxix

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second—continued.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 60°.									N.P.D.
	61°.	62°.	63°.	64°.	65°.	66°.	67°.	68°.	69°.	
0	10°3708	10°5284	10°6800	10°8259	10°9660	11°1002	11°2289	11°3519	11°4692	0
1	3735	5310	6825	8283	9683	1024	2310	3539	4711	1
2	3761	5336	6850	8307	9706	1046	2331	3559	4730	2
3	3788	5361	6875	8331	9728	1067	2352	3579	4749	3
4	3814	5387	6899	8354	9751	1089	2373	3599	4768	4
5	3841	5412	6924	8378	9774	1111	2394	3619	4787	5
6	3867	5438	6949	8402	9796	1133	2415	3639	4806	6
7	3894	5463	6974	8426	9819	1155	2436	3659	4825	7
8	3920	5488	6998	8449	9842	1176	2456	3678	4844	8
9	3947	5514	7022	8473	9864	1198	2477	3698	4863	9
10	3973	5540	7047	8497	9887	1220	2498	3718	4882	10
11	4000	5566	7071	8521	9910	1242	2519	3738	4901	11
12	4027	5591	7096	8544	9932	1264	2540	3758	4920	12
13	4054	5617	7121	8567	9955	1285	2560	3777	4938	13
14	4080	5643	7146	8591	10°9978	1307	2581	3797	4957	14
15	4107	5668	7171	8614	11°0001	1329	2602	3817	4976	15
16	4134	5694	7196	8638	0023	1351	2623	3837	4995	16
17	4160	5719	7220	8661	0046	1372	2643	3857	5014	17
18	4186	5745	7245	8685	0069	1394	2664	3876	5032	18
19	4213	5770	7269	8708	0091	1415	2684	3896	5051	19
20	4239	5796	7294	8732	0113	1437	2705	3916	5070	20
21	4265	5821	7318	8755	0135	1459	2726	3936	5089	21
22	4292	5846	7343	8779	0158	1480	2746	3955	5108	22
23	4318	5872	7367	8802	0180	1502	2767	3975	5126	23
24	4345	5897	7391	8826	0203	1523	2787	3994	5145	24
25	4371	5922	7415	8849	0225	1545	2808	4014	5164	25
26	4397	5948	7439	8873	0248	1567	2829	4034	5183	26
27	4424	5973	7464	8896	0270	1588	2849	4053	5201	27
28	4450	5998	7488	8920	0293	1610	2870	4073	5220	28
29	4476	6024	7513	8943	0315	1631	2890	4092	5238	29
30	4503	6049	7537	8966	0338	1653	2911	4112	5257	30
31	4529	6074	7561	8989	0360	1674	2931	4132	5276	31
32	4556	6100	7586	9013	0383	1696	2952	4151	5294	32
33	4582	6125	7610	9036	0405	1717	2972	4171	5313	33
34	4608	6150	7634	9059	0427	1739	2993	4190	5331	34
35	4634	6176	7659	9082	0449	1760	3013	4210	5350	35
36	4660	6201	7683	9106	0471	1781	3033	4229	5369	36
37	4686	6226	7707	9129	0494	1803	3054	4249	5387	37
38	4712	6251	7732	9152	0516	1824	3074	4268	5406	38
39	4738	6276	7756	9175	0538	1846	3095	4288	5424	39
40	4765	6301	7780	9199	0560	1867	3115	4307	5443	40
41	4791	6326	7804	9222	0582	1888	3135	4326	5461	41
42	4817	6351	7828	9245	0605	1909	3156	4346	5480	42
43	4843	6376	7852	9268	0627	1931	3176	4365	5498	43
44	4869	6401	7876	9292	0649	1952	3197	4384	5517	44
45	4895	6426	7900	9315	0671	1973	3217	4404	5535	45
46	10°4921	10°6451	10°7924	10°9338	11°0693	11°1994	11°3237	11°4423	11°5553	46

cxxxii INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second—continued.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 60°.									N.P.D.
	61°.	62°.	63°.	64°.	65°.	66°.	67°.	68°.	69°.	
47	10"4947	10"6476	10"7948	10"9361	11"0716	11"2015	11"3257	11"4442	11"5572	47
48	4974	6501	7972	9384	0738	2037	3278	4462	5590	48
49	5000	6526	7996	9407	0760	2058	3298	4481	5609	49
50	5026	6551	8020	9430	0782	2079	3318	4500	5627	50
51	5052	6576	8044	9453	0804	2100	3338	4519	5645	51
52	5078	6602	8068	9476	0826	2121	3358	4538	5663	52
53	5103	6627	8092	9499	0848	2142	3379	4558	5682	53
54	5129	6652	8116	9522	0870	2163	3399	4577	5700	54
55	5155	6677	8140	9545	0892	2184	3419	4596	5718	55
56	5181	6701	8164	9568	0914	2205	3439	4615	5736	56
57	5207	6726	8188	9591	0936	2226	3459	4634	5754	57
58	5232	6751	8212	9614	0958	2247	3479	4654	5773	58
59	5258	6776	8236	9637	0980	2268	3499	4673	5791	59
60	10"5284	10"6800	10"8259	10"9660	11"1002	11"2289	11"3519	11"4692	11"5809	60

N.P.D.	Azimuth 60°.									N.P.D.
	70°.	71°.	72°.	73°.	74°.	75°.	76°.	77°.	78°.	
0	11"5809	11"6872	11"7878	11"8829	11"9726	12"0568	12"1355	12"2087	12"2765	0
1	5827	6889	7894	8844	9740	0582	1368	2099	2776	1
2	5845	6906	7910	8860	9755	0595	1380	2111	2787	2
3	5864	6924	7927	8875	9769	0609	1393	2122	2797	3
4	5882	6941	7943	8891	9784	0622	1405	2134	2808	4
5	5900	6958	7959	8906	9798	0636	1418	2146	2819	5
6	5918	6975	7975	8921	9812	0649	1431	2158	2830	6
7	5936	6992	7991	8936	9827	0663	1443	2169	2841	7
8	5954	7010	8008	8952	9841	0676	1456	2181	2851	8
9	5972	7027	8024	8967	9856	0690	1468	2192	2862	9
10	5990	7044	8040	8982	9870	0703	1481	2204	2873	10
11	6008	7061	8056	8997	9884	0716	1493	2216	2884	11
12	6026	7078	8072	9012	9899	0730	1506	2227	2894	12
13	6044	7095	8089	9028	9913	0743	1518	2239	2905	13
14	6062	7112	8105	9043	9928	0757	1531	2250	2915	14
15	6080	7129	8121	9058	9942	0770	1543	2262	2926	15
16	6098	7146	8137	9073	9956	0783	1555	2273	2937	16
17	6116	7163	8153	9088	9970	0796	1568	2285	2947	17
18	6133	7180	8169	9104	9985	0810	1580	2296	2958	18
19	6151	7197	8185	9119	11"9999	0823	1593	2308	2968	19
20	6169	7214	8201	9134	12"0013	0836	1605	2319	2979	20
21	6187	7231	8217	9149	0027	0849	1617	2330	2989	21
22	6205	7248	8233	9164	0041	0862	1630	2342	3000	22
23	6222	7264	8249	9179	0056	0876	1642	2353	3010	23
24	11"6240	11"7281	11"8265	11"9194	12"0070	12"0889	12"1655	12"2365	12"3021	24

ALTAZIMUTH TABLES, 1908.

cxxxiii

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second—continued.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 60°.									N.P.D.
	70°.	71°.	72°.	73°.	74°.	75°.	76°.	77°.	78°.	
25	11°6258	11°7298	11°8281	11°9209	12°0084	12°0902	12°1667	12°2376	12°3031	25
26	6276	7315	8297	9224	0098	0915	1679	2387	3041	26
27	6294	7332	8313	9239	0112	0928	1691	2399	3052	27
28	6311	7348	8328	9254	0126	0942	1704	2410	3062	28
29	6329	7365	8344	9269	0140	0955	1716	2422	3073	29
30	6347	7382	8360	9284	0154	0968	1728	2433	3083	30
31	6365	7399	8376	9299	0168	0981	1740	2444	3093	31
32	6382	7416	8392	9314	0182	0994	1752	2455	3104	32
33	6400	7432	8407	9329	0196	1007	1765	2467	3114	33
34	6417	7449	8423	9344	0210	1020	1777	2478	3125	34
35	6435	7466	8439	9359	0224	1033	1789	2489	3135	35
36	6453	7483	8455	9374	0238	1046	1801	2500	3145	36
37	6470	7499	8471	9389	0252	1059	1813	2511	3155	37
38	6488	7516	8486	9403	0266	1072	1825	2523	3166	38
39	6505	7532	8502	9418	0280	1085	1837	2534	3176	39
40	6523	7549	8518	9433	0294	1098	1849	2545	3186	40
41	6541	7566	8534	9448	0308	1111	1861	2556	3196	41
42	6558	7582	8549	9463	0322	1124	1873	2567	3206	42
43	6576	7599	8565	9477	0335	1137	1885	2579	3217	43
44	6593	7615	8580	9492	0349	1150	1897	2590	3227	44
45	6611	7632	8596	9507	0363	1163	1909	2601	3237	45
46	6628	7648	8612	9522	0377	1176	1921	2612	3247	46
47	6646	7665	8627	9536	0391	1189	1933	2623	3257	47
48	6663	7681	8643	9551	0404	1201	1945	2634	3268	48
49	6681	7698	8658	9565	0418	1214	1957	2645	3278	49
50	6698	7714	8674	9580	0432	1227	1969	2656	3288	50
51	6715	7730	8690	9595	0446	1240	1981	2667	3298	51
52	6733	7747	8705	9609	0459	1253	1993	2678	3308	52
53	6750	7763	8721	9624	0473	1265	2004	2689	3318	53
54	6768	7780	8736	9638	0486	1278	2016	2700	3328	54
55	6785	7796	8752	9653	0500	1291	2028	2711	3338	55
56	6802	7812	8767	9668	0514	1304	2040	2722	3348	56
57	6820	7829	8783	9682	0527	1317	2052	2733	3358	57
58	6837	7845	8798	9697	0541	1329	2063	2743	3368	58
59	6855	7862	8814	9711	0554	1342	2075	2754	3378	59
60	11°6872	11°7878	11°8829	11°9726	12°0568	12°1355	12°2087	12°2765	12°3388	60

cxixiv INTRODUCTION TO GREENWICH ASTRONOMICAL OBSERVATIONS, 1908.

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second—continued.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 60°.											N.P.D.
	79°.	80°.	81°.	82°.	83°.	84°.	85°.	86°.	87°.	88°.	89°.	
0	12°3388	12°3957	12°4472	12°4932	12°5339	12°5690	12°5988	12°6231	12°6421	12°6556	12°6637	0
1	3398	3966	4480	4939	5345	5695	5992	6235	6424	6558	6638	1
2	3408	3975	4488	4946	5351	5700	5997	6238	6426	6560	6639	2
3	3417	3984	4496	4954	5358	5707	6001	6242	6429	6561	6639	3
4	3427	3993	4504	4961	5364	5712	6006	6245	6431	6563	6640	4
5	3437	4002	4512	4968	5370	5717	6010	6249	6434	6565	6641	5
6	3447	4011	4520	4975	5376	5722	6014	6253	6437	6567	6642	6
7	3457	4020	4528	4982	5382	5728	6019	6256	6439	6569	6643	7
8	3466	4029	4536	4990	5389	5733	6023	6260	6442	6570	6643	8
9	3476	4038	4544	4997	5395	5739	6028	6263	6444	6572	6644	9
10	3486	4047	4552	5004	5401	5744	6032	6267	6447	6574	6645	10
11	3496	4056	4560	5011	5407	5749	6036	6270	6450	6576	6646	11
12	3506	4065	4568	5018	5413	5754	6041	6274	6452	6577	6647	12
13	3515	4073	4576	5025	5420	5760	6045	6277	6455	6579	6647	13
14	3525	4082	4584	5032	5426	5765	6050	6281	6457	6580	6648	14
15	3535	4091	4592	5039	5432	5770	6054	6284	6460	6582	6649	15
16	3545	4100	4600	5046	5438	5775	6058	6287	6462	6584	6650	16
17	3554	4109	4608	5053	5444	5780	6062	6291	6465	6585	6650	17
18	3564	4117	4615	5060	5450	5786	6067	6294	6467	6587	6651	18
19	3573	4126	4623	5067	5456	5791	6071	6298	6470	6588	6651	19
20	3583	4135	4631	5074	5462	5796	6075	6301	6472	6590	6652	20
21	3593	4144	4639	5081	5468	5801	6079	6304	6474	6591	6653	21
22	3602	4153	4647	5088	5474	5806	6083	6307	6477	6593	6653	22
23	3612	4160	4654	5094	5480	5811	6088	6311	6479	6594	6654	23
24	3621	4169	4662	5101	5486	5816	6092	6314	6482	6596	6654	24
25	3631	4178	4670	5108	5492	5821	6096	6317	6484	6597	6655	25
26	3641	4187	4678	5115	5498	5826	6100	6320	6486	6598	6655	26
27	3650	4195	4686	5122	5504	5831	6104	6323	6488	6600	6656	27
28	3660	4204	4693	5128	5509	5836	6108	6327	6491	6601	6656	28
29	3669	4212	4701	5135	5515	5841	6112	6330	6493	6603	6657	29
30	3679	4221	4709	5142	5521	5846	6116	6333	6495	6604	6657	30
31	3688	4230	4717	5149	5527	5851	6120	6336	6497	6605	6657	31
32	3698	4238	4724	5156	5533	5856	6124	6339	6499	6606	6658	32
33	3707	4247	4732	5162	5538	5861	6128	6343	6502	6608	6658	33
34	3717	4255	4739	5169	5544	5866	6132	6346	6504	6609	6659	34
35	3726	4264	4747	5176	5550	5871	6136	6349	6506	6610	6659	35
36	3735	4272	4755	5183	5556	5876	6140	6352	6508	6611	6659	36
37	3745	4281	4762	5189	5562	5881	6144	6355	6510	6612	6660	37
38	3754	4289	4770	5196	5567	5886	6148	6358	6513	6614	6660	38
39	3764	4298	4777	5202	5573	5891	6152	6361	6515	6615	6661	39
40	3773	4306	4785	5209	5579	5895	6156	6364	6517	6616	6661	40
41	3782	4314	4792	5216	5585	5900	6160	6367	6519	6617	6661	41
42	3792	4323	4800	5222	5590	5905	6164	6370	6521	6618	6661	42
43	3801	4331	4807	5229	5596	5909	6167	6373	6523	6620	6662	43
44	3811	4339	4815	5235	5601	5914	6171	6376	6525	6621	6662	44
45	3820	4348	4822	5242	5607	5919	6175	6379	6527	6622	6662	45
46	12°3829	12°4356	12°4829	12°5249	12°5613	12°5924	12°6179	12°6382	12°6529	12°6623	12°6662	46

ALTAZIMUTH TABLES, 1908.

CXXXV

Azimuthal Motion in Great Circle in One Sidereal Second, or of a Moon moving uniformly parallel to the Equator in One Lunar Second—concluded.

This term is referred to as "Collimation Divisor" in the Extra-Meridian Observations.

N.P.D.	Azimuth 60°.										N.P.D.	
	79°.	80°.	81°.	82°.	83°.	84°.	85°.	86°.	87°.	88°.		89°.
47	12"3838	12"4365	12"4837	12"5255	12"5618	12"5928	12"6183	12"6385	12"6531	12"6624	12"6662	47
48	3848	4373	4844	5262	5624	5933	6186	6387	6533	6625	6663	48
49	3857	4382	4852	5268	5629	5937	6190	6390	6535	6626	6663	49
50	3866	4390	4859	5275	5635	5942	6194	6393	6537	6627	6663	50
51	3875	4398	4866	5281	5641	5947	6198	6396	6539	6628	6663	51
52	3884	4406	4874	5288	5646	5951	6202	6399	6541	6629	6663	52
53	3894	4415	4881	5294	5652	5956	6205	6401	6543	6630	6664	53
54	3903	4423	4889	5301	5657	5960	6209	6404	6545	6631	6664	54
55	3912	4431	4896	5307	5663	5965	6213	6407	6547	6632	6664	55
56	3921	4439	4903	5313	5668	5970	6217	6410	6549	6633	6664	56
57	3930	4447	4910	5320	5674	5975	6220	6413	6551	6634	6664	57
58	3939	4456	4918	5326	5679	5978	6224	6415	6552	6635	6664	58
59	3948	4464	4925	5333	5685	5983	6227	6418	6554	6636	6664	59
60	12"3957	12"4472	12"4932	12"5339	12"5690	12"5988	12"6231	12"6421	12"6556	12"6637	12"6664	60