

FORMATION OF SATURN'S E-RING

K. E. Herkenhoff and D. J. Stevenson

The observed geometry of the E-ring has not been adequately explained, in particular, the vertical thickening with increasing distance from Saturn is not understood. Dynamical evolution requires more than the short lifetime of the ring to produce the present configuration: at the present grain density of $4 \times 10^{-9} \text{ cm}^{-3}$, the mean time between collisions is at least 200 years. NH_3 particles nucleated from a 173°K melt will leave Enceladus at $7.3 \times 10^4 \text{ cm s}^{-1}$ and form a lens-shaped ring extending from 3.15 to $5.01 R_s$ with a vertical thickness of about $3 \times 10^4 \text{ km}$ [Fig. 2]. To achieve the present shape of the ring, some other mechanism(s) must be invoked. Gravitational scattering by the larger satellites Tethys and Dione may be important to the evolution of the ring. The magnetospheric plasma probably imparts an electrostatic potential on the grains, causing orbital perturbations due to Lorentz forcing. If a 1μ ammonia grain is charged to a voltage similar to the plasma electron energy (over 20 eV), the Lorentz force will be over 1% of the gravitational force. Radiation pressure will make the orbits gradually decay, and direct particle drag from impinging ions and distant Coulomb drag spread the ring at a rate dependent on the grain charge and plasma parameters. A numerical model including these effects would contribute much toward the understanding of the E-ring's structure.




Figure 2: Comparison of initial distribution of ammonia ice grains formed from a 173°K melt (top) and the three-dimensional distribution of E-ring particles found by Baum et al. [1]. M, E, T, D and R mark the orbits of Mimas, Enceladus, Tethys, Dione and Rhea, respectively. Filled circles represent the attenuation per kilometer (left-hand ordinate) in the equatorial plane, while open circles represent the true physical thickness (right-hand ordinate) measured perpendicular to the equatorial plane.

REFERENCES

- [1] Baum, W. A. et al. "Saturn's E-Ring" (pre-print, 1983)
- [2] Baum, W. A. et al. *Icarus* 47, 84 (1980)
- [3] Pang, K. D. et al. *Lunar and Planetary Science abstracts* 14, 592 (1983)
- [4] McKinnon, W. B. *Lunar and Planetary Science abstracts* 14, 487 (1983)
- [5] Smith, B. A. et al. *Science* 215, 504 (1982)
- [6] Squyres, S. W. et al. *Icarus* 53, 319 (1983)
- [7] Stevenson, D. J. *Nature* 298, 142 (1982)
- [8] Raizer, Y. P. *Soviet Physics JETP* 37, 1229 (1960)
- [9] Haff, P. K. et al. "Ring and Plasma: The Enigma of Enceladus" (preprint, 1983)
- [10] Morfill, G. E. et al. *J. Geophys. Res.* 88, 5573 (1983)