Post-outburst phase of McNeil's nebula (V1647 Orionis)

D. K. Ojha,^{1*} S. K. Ghosh,¹ A. Tej,¹ R. P. Verma,¹ S. Vig,¹ G. C. Anupama,² D. K. Sahu,² P. Parihar,² B. C. Bhatt,² T. P. Prabhu,² G. Maheswar,² H. C. Bhatt,² B. G. Anandarao³ and V. Venkataraman³

¹Tata Institute of Fundamental Research, Mumbai (Bombay) 400 005, India ²Indian Institute of Astrophysics, Koramangala, Bangalore 560 034, India ³Physical Research Laboratory, Ahmedabad 390 009, India

Accepted 2006 February 3. Received 2006 February 2; in original form 2005 December 13

ABSTRACT

We present a detailed study of the post-outburst phase of McNeil's nebula (V1647 Orionis) using optical B, V, R, I and near-infrared (NIR) J, H, K photometric and low-resolution optical spectroscopic observations. The observations were carried out with the Himalaya Faint Object Spectrograph Camera (HFOSC), NIR camera (NIRCAM), the Tata Institute of Fundamental Research (TIFR) Near-Infrared Camera (TIRCAM) and NICMOS cameras on the 2-m Himalayan Chandra Telescope (HCT) and 1.2-m Physical Research Laboratory (PRL) telescopes during the period 2004 February-2005 December. The optical and NIR observations show a general decline in the brightness of the exciting source of McNeil's nebula (V1647 Ori). Our recent optical images show that V1647 Ori has faded by more than 3 mag since February 2004. McNeil's nebula has also faded considerably. The optical/NIR photometric data also show a significant variation in the magnitudes ($\Delta V = 0.78$, $\Delta R = 0.44$, $\Delta I = 0.21$, $\Delta J = 0.24$ and $\Delta H = 0.20$ mag) of V1647 Ori within a period of one month, which is possibly undergoing a phase similar to eruptive variables, like EXors or FUors. The optical spectra show a few features such as strong H α emission with blue-shifted absorption and the Ca II IR triplet (8498, 8542 and 8662 Å) in emission. As compared to the period just after outburst, there is a decrease in the depth and extent of the blue-shifted absorption component, indicating a weakening in the powerful stellar wind. The presence of the Ca II IR triplet in emission confirms that V1647 Ori is a pre-main-sequence star. The long-term, post-outburst photometric observations of V1647 Ori suggest an EXor rather than an FUor event. An optical/IR comparison of the region surrounding McNeil's nebula shows that the optical nebula is more widely and predominantly extended to the north, whereas the IR nebula is relatively confined (diameter \sim 60 arcsec), but definitely extended, to the south, too. The large colour gradient from north to south and the sudden absence of an optical nebula to the south are suggestive of a large-scale disc-like structure (or envelope) surrounding the central source that hides the southern nebula.

Key words: stars: formation – stars: pre-main-sequence – stars: variables: other – ISM: individual: (McNeil's nebula) – Reflection nebulae.

1 INTRODUCTION

On the night of 2004 January 23, Jay McNeil discovered a new reflection nebula in the L1630 cloud in Orion (McNeil 2004). The nebula, which now bears his name, is recognized as a reflection nebula surrounding a young stellar object (YSO). It appears to be a faint optical counterpart of the infrared (IR) source catalogued as

IRAS 05436–0007 that had gone into outburst, producing a large reflection nebulosity (Ábrahám et al. 2004; Andrews, Rothberg & Simon 2004; Briceño et al. 2004; Kun et al. 2004; Reipurth & Aspin 2004; Vacca, Cushing & Simon 2004; Walter et al. 2004; Ojha et al. 2005). Later, the McNeil's object was designated as V1647 Orionis (Samus 2004). Reipurth & Aspin (2004) reported that NIR images taken with the Gemini 8-m telescope show that the object has brightened by about 3 mag in *JHK*, relative to the 1998 2-micron All-Sky Survey (2MASS) measurements. Gemini spectra reveal strong features of CO and hydrogen Br γ in emission in the IR. In the optical,

^{*}E-mail: ojha@tifr.res.in

 $H\alpha$ appears in emission with a P Cygni profile (Reipurth & Aspin 2004).

McGehee et al. (2004) described the change in optical and NIR colours of V1647 Ori from its 'quiescent' state to its outburst. They concluded that the colour changes associated with brightening suggest an EXor-like outburst rather than a simple dust clearing event. Kun et al. (2004) analysed optical and NIR measurements of V1647 Ori and suggested that the outburst was due to an intrinsic increase in the luminosity of the central object (rather than a decrease in extinction), and identified the object as an FUor, rather than an EXor. Vacca et al. (2004) presented NIR spectroscopy of the object and suggested that V1647 Ori is a heavily embedded low-mass Class I protostar, surrounded by a disc, whose brightening is due to a recent accretion event. Ábrahám et al. (2004) described IR and sub-mm properties of V1647 Ori and concluded that the object has a disc of about 0.5 M_{\odot}, a pre-outburst bolometric luminosity of 5.6 L_{\odot}, and is probably a Class II protostar of age of about 0.4 Myr. Andrews et al. (2004) presented the measurements of V1647 Ori in the mid-IR and sub-mm, and concluded that the 12-µm flux of this source has increased by a factor of 25 after the outburst, whereas the sub-mm continuum remains at its pre-outburst level. The IR slope of the spectral energy distribution (SED) characterizes the illuminating source as a flat-spectrum protostar, in its both active and quiescent states. Muzerolle et al. (2005) report on Spitzer NIR photometry of V1647 Ori in its early post-outburst phase (2004 March). They found that the brightness of the star increased by roughly the same factor of 15-20 across the optical and NIR. Their results indicated that the outburst may be intermediate between FUor- and EXor-type events. Aspin & Reipurth (2005) reported on observations made at the Gemini telescope and showed that V1647 Ori has faded by approximately 1 mag in r' band since 2005 January 8. Kóspál et al. (2005) reported the most recent VRI measurements of V1647 Ori obtained between 2005 October 4 and November 19 which show that V1647 Ori has started a very rapid fading ($\sim 1 \text{ mag month}^{-1}$). They concluded that V1647 Ori is a unique (somewhat intermediate) object among FUors and EXors.

The McNeil's source at the apex of the nebula was also observed to brighten dramatically in X-ray wavelength. The X-ray data from Chandra give strong evidence that the probable cause of the outburst is the sudden infall of gas on to the surface of the star from an orbiting disc of gas (Kastner et al. 2004). Grosso et al. (2005) describe X-ray spectra of V1647 Ori taken with XMM-Newton on 2004 April 4. They find variations in both the brightness and the hardness of the source. There is a very exciting hint of periodicity in the X-ray brightness, which might provide information on the physical nature of the X-ray luminosity. Vig et al. (2004, 2005) presented post-outburst radio continuum observations of the region around McNeil's nebula and report no radio continuum emission from the nebula. They explored various scenarios to explain the radio upper limit of the nebula and suggested that the homogeneous H II region scenario is consistent with the radio, H α as well as X-ray measurements.

In this paper, we present the optical and NIR morphologies of Mc-Neil's nebula and report on the optical/NIR photometric and optical spectroscopic monitoring of V1647 Ori beginning with the epoch of the recent outburst. Our results provide variability measurements of V1647 Ori for a duration of about 21 months and allow us to check if the central star has changed significantly since its emergence in early 2004. Our optical/NIR observations also show significant variations in the source brightness within periods of one week and one month. In Section 2, we present the details of observations and datareduction procedures; Section 3 deals with the results and discussion concerning the short and long time variability of V1647 Ori that illuminates McNeil's nebula, and describes the optical/IR nebula in detail. We also discuss spectroscopic results in this section. We then summarize our conclusions in Section 4.

2 OBSERVATIONS AND DATA REDUCTION

2.1 Optical photometry

Imaging observations of McNeil's nebula region at optical wavelengths (standard Bessel BVRIH α filters) were obtained on 19 nights during 2004 February-2005 November with the 2-m Himalayan Chandra Telescope (HCT) situated at the Indian Astronomical Observatory (IAO), Hanle, India. The Himalaya Faint Object Spectrograph Camera (HFOSC) instrument equipped with a SITe $2K \times 4K$ pixel² CCD was used. The central $2K \times 2K$ region, used for imaging, corresponds to a field of view (FOV) of $\sim 10 \times 10$ arcmin², with a pixel scale of 0.296 arcsec. Further details on the telescope and the instrument can be found at http://www.iiap.res.in/iao/hfosc.html. The observations were carried out under good photometric sky conditions. The average seeing [full width at half-maximum (FWHM)] in all the bands was ~ 1.6 arcsec during the observations. Bias and flat frames were obtained at the beginning and at the end of each observing night. Atmospheric extinction and transformation coefficients were obtained from the observations of photometric standard stars around 98-556, 98-618, 98-961, PG 1525-071 and PG 0231+051 regions (Landolt 1992). The details of the observations are given in Table 1. The Landolt standard regions were not observed on five nights, hence the magnitudes of V1647 Ori were obtained by using a reference star as a secondary standard in the field (see Table 1). During five nights, observations were carried out for the target in R filter only, and a reference star (see Section 3) in the field was used to obtain the differential magnitudes.

Data reduction was done using the National Optical Astronomy Observatories' (NOAO) IRAF ¹ package tasks. Flat-fielding with a median flat frame was applied. Identification and photometry of point sources were performed by using the DAOFIND and DAOPHOT packages in IRAF, respectively. Because of nebulosity around V1647 Ori, photometry was obtained using the point-spread function (PSF) algorithm ALLSTAR in the DAOPHOT package (Stetson 1987). The residuals to the photometric solution are ≤ 0.05 mag. The absolute optical photometry of V1647 Ori is presented in Table 2.

2.2 NIR photometry

The *JHK* NIR observations were carried out during 2004 February 12, October 18, November 19, 2005 November 28, 29 and 30 with the HCT NIR camera (NIRCAM), equipped with 512 × 512 Mercury Cadmium Telluride (HgCdTe) array of 18-µm pixel size. For our observations, the NIRCAM was used in the *Camera-B* mode which has a FOV of 3.6×3.6 arcmin². Further details on NIRCAM can be found at http://www.iiap.res.in/iao/nir.html. We obtained several dithered exposures of the target centred at $(\alpha, \delta)_{2000} = (05^{h}46^{m}14^{\circ}.0, -00^{\circ}0'40''.1)$ in each of the NIR bands. Typical integration times per frame were 20, 15 and 5 s in the *J*, *H* and *K* bands, respectively. The images were co-added to obtain the final image in each band. We also obtained several dithered sky

¹ IRAF is distributed by the NOAO, which are operated by the Association of Universities for Research in Astronomy, Inc. under contract with the National Science Foundation.

Date (UT)	JD	FWHM (arcsec) ^a	Filter(s) ^b	Exposure time (s)	Notes
2004 Feb 12	245 3048.1	1.2	R, I	480, 240	1
2004 Feb 13	245 3049.1	2.3	R	90	2
2004 Feb 14	245 3050.2	1.5	V, R, I	200, 270, 90, 900	
2004 Feb 22	245 3057.9	1.4	V, R, I	300, 150, 150	
2004 Feb 23	245 3058.9	1.7	B, R	1200, 150	1
2004 Oct 18	245 3297.4	2.3	B, V, R, I	300, 300, 300, 300, 600	
2004 Nov 17	245 3327.4	1.4	B, V, R, I	600, 900, 540, 360, 900	
2004 Dec 7	245 3347.3	1.8	B, V, R, I	600, 720, 600, 480, 1200	
2005 Jan 8	245 3379.1	1.5	R	120	2
2005 Jan 9	245 3380.1	2.1	R	60	2
2005 Mar 28	245 3458.1	1.7	R	900	2
2005 Sept 26	245 3639.6	1.5	V, R	1200, 600	
2005 Sept 29	245 3643.5	1.4	B, V, R	900, 450, 450	
2005 Nov 9	245 3684.3	1.3	V, R, I	300, 440, 420	1
2005 Nov 20	245 3695.2	2.0	V, R, I	600, 450, 450	1
2005 Nov 21	245 3696.4	1.3	V, R, I	420, 420, 420	1
2005 Nov 28	245 3703.2	1.5	V, R, I	600, 1800, 720	
2005 Nov 29	245 3704.2	1.7	R	1200	2
2005 Nov 30	245 3705.2	1.6	V, R, I	600, 1200, 720	

Table 1. Observing log of optical photometric observations.

^aMeasured average FWHM. This is a measure of the seeing.

^bObservations were made in Bessel B, V, R, I filters.

¹Magnitudes of V1647 Ori were obtained by using a reference star as a secondary standard in the field.

²Only differential photometry was done.

Date (UT)	V1647 Ori ^a			McNeil's nebula ^b				
	В	V	R	Ι	В	V	R	Ι
2004 Feb 12 ^c			16.91±0.02	14.82 ± 0.02		13.18±0.02	12.13±0.02	
2004 Feb 14		$18.68 {\pm} 0.04$	$16.83 {\pm} 0.02$	$14.86 {\pm} 0.02$		$13.98 {\pm} 0.04$	$13.19 {\pm} 0.02$	12.19 ± 0.02
2004 Feb 22		$18.85 {\pm} 0.01$	16.92 ± 0.02	$14.81 {\pm} 0.01$	$14.54 {\pm} 0.01$	13.62 ± 0.01	$12.98 {\pm} 0.02$	$11.94{\pm}0.01$
2004 Feb 23 ^c	20.44 ± 0.01		$16.74 {\pm} 0.01$		$14.44 {\pm} 0.01$		$13.00 {\pm} 0.01$	
2004 Oct 18 ^d		18.02 ± 0.05	16.51 ± 0.05	$14.76 {\pm} 0.05$	$14.77 {\pm} 0.05$	$13.87 {\pm} 0.05$	$13.16 {\pm} 0.05$	12.05 ± 0.05
2004 Nov 17	20.64 ± 0.04	$18.80 {\pm} 0.02$	$16.95 {\pm} 0.01$	$14.97 {\pm} 0.02$	$14.85 {\pm} 0.04$	$13.87 {\pm} 0.02$	13.19 ± 0.01	12.09 ± 0.02
2004 Dec 7	20.72 ± 0.05	$19.08 {\pm} 0.01$	17.15 ± 0.02	$15.13 {\pm} 0.02$	$14.82 {\pm} 0.05$	$13.90 {\pm} 0.01$	13.26 ± 0.02	12.05 ± 0.02
2005 Sept 26 ^d		$19.82 {\pm} 0.02$	17.94 ± 0.02			14.92 ± 0.02	14.12 ± 0.02	
2005 Sept 29 ^d		$19.82 {\pm} 0.01$	18.09 ± 0.01		$15.88 {\pm} 0.01$	15.01 ± 0.01	14.11 ± 0.01	
2005 Nov 9 ^c		21.45 ± 0.05	$19.46 {\pm} 0.04$	17.34 ± 0.05		16.02 ± 0.05	$14.94{\pm}0.04$	$13.57 {\pm} 0.05$
2005 Nov 20 ^c		>21.45 ^e	$19.76 {\pm} 0.04$	17.61 ± 0.02		$16.66 {\pm} 0.05$	15.24 ± 0.04	13.72 ± 0.02
2005 Nov 21 ^c		>21.45 ^e	$19.83 {\pm} 0.02$	17.56 ± 0.02		16.96 ± 0.04	15.19 ± 0.02	$13.58 {\pm} 0.02$
2005 Nov 28		22.11 ± 0.05	20.02 ± 0.03	$17.93 {\pm} 0.05$		$16.90 {\pm} 0.05$	$15.14 {\pm} 0.03$	$13.60 {\pm} 0.05$
2005 Nov 30		>22.11 ^e	$20.10 {\pm} 0.05$	17.95±0.02		$16.62 {\pm} 0.05$	$15.18 {\pm} 0.05$	13.78±0.02

^aPSF magnitudes.

^bMagnitudes in 80-arcsec aperture centred on V1647 Ori.

^cMagnitudes are derived using star A (see text) as a secondary calibrator.

 d Too faint to be detected in *B* band.

^eSource is not detected in V band with exposure time of 600 s.

frames close to the target position in each NIR band. Photometric calibration was done from the observations made on the same nights on the standard stars around AS13 region (Hunt et al. 1998) at airmasses close to that of the target observations. We used an aperture radius of 18 pixels (\sim 7 arcsec) for photometry of V1647 Ori. The local sky was evaluated in an annulus with an inner radius of 128 pixel and a width of 12 pixel.

The *JHK* NIR observations were also carried out on 2004 October 12, using TIFR Near-Infrared Camera (TIRCAM) at the f/13 Cassegrain focus of the Mount Abu 1.2-m telescope belonging to Physical Research Laboratory (PRL, India). TIRCAM is based on the Santa Barbara Research Center (SBRC) InSb Focal Plane Array ($58 \times 62 \text{ pixel}^2$) sensitive between 1 and 5 µm. The details of the TIRCAM system are described in Ojha et al. (2002). Individual object frames were of 2-s integration in *J*, *H* and *K* band. Photometric calibration was obtained by observing the standard star HR1852 from the list of the UK Infrared Telescope's (UKIRT's) bright standards.²

We also made NIR *JHK'* photometric observations by using PRL NICMOS camera on Mount Abu 1.2-m telescope. At the core of the

² http://www.jach.hawaii.edu/UKIRT/astronomy/calib/phot_cal/bright_stds.html

Date (UT)	J	Н	Κ	FWHM (arcsec)	Telescope
1998 Oct 7	14.74±0.03	12.16±0.03	10.27 ± 0.02		2MASS
2004 Feb 3	11.1 ± 0.1	9.0 ± 0.1	7.4 ± 0.1	0.5	Gemini ^a
2004 Feb 11	10.79 ± 0.01	$8.83 {\pm} 0.01$	7.72 ± 0.01		$USNO^b$
2004 Feb 12	$10.68 {\pm} 0.04$	$8.78 {\pm} 0.04$		1.2	HCT
2004 Feb 23	$10.74{\pm}0.05$	$8.93 {\pm} 0.01$	$7.40 {\pm} 0.02$	1.6	IRSF ^c
2004 Feb 25	$10.79 {\pm} 0.05$	$8.95 {\pm} 0.01$	7.47 ± 0.01	0.9	IRSF ^c
2004 Feb 29	$10.90 {\pm} 0.06$	9.11 ± 0.01	$7.62 {\pm} 0.02$	1.4	IRSF ^c
2004 Mar 20	$10.85 {\pm} 0.07$	$8.83 {\pm} 0.02$	$7.71 {\pm} 0.05$	1.7	PRL/NICMOS
2004 Mar 21	$10.76 {\pm} 0.02$	$8.91 {\pm} 0.01$	$7.59 {\pm} 0.03$	1.8	PRL/NICMOS
2004 Apr 11	11.09 ± 0.03	$9.01 {\pm} 0.02$	$7.49 {\pm} 0.03$	2.5	PRL/NICMOS
2004 Apr 12	$10.94{\pm}0.02$	$9.06 {\pm} 0.02$	$7.59 {\pm} 0.02$		$USNO^b$
2004 Oct 12	$10.76 {\pm} 0.06$	$8.87 {\pm} 0.06$	$7.34{\pm}0.06$	2.1	TIRCAM
2004 Oct 16	11.15 ± 0.01	$9.24{\pm}0.02$	$7.89 {\pm} 0.01$	1.7	PRL/NICMOS
2004 Oct 17	11.18 ± 0.02	9.23 ± 0.01	$7.87 {\pm} 0.02$	1.7	PRL/NICMOS
2004 Oct 18	$10.89 {\pm} 0.04$	9.13 ± 0.04	$7.48 {\pm} 0.04$	2.3	HCT
2004 Nov 19	11.13 ± 0.04	$9.33 {\pm} 0.04$	<8.03	1.0	HCT^d
2004 Dec 21	$11.04{\pm}0.03$	$9.20 {\pm} 0.01$	$7.83 {\pm} 0.03$	2.3	PRL/NICMOS
2004 Dec 22	11.03 ± 0.02	$8.98 {\pm} 0.01$	7.76 ± 0.03	2.1	PRL/NICMOS
2005 Nov 25		$10.02 {\pm} 0.07$		2.5	PRL/NICMOS
2005 Nov 26	13.28 ± 0.15	$10.69 {\pm} 0.09$		2.5	PRL/NICMOS
2005 Nov 28		$10.80 {\pm} 0.02$	$8.95 {\pm} 0.03$	1.5	HCT
2005 Nov 29	13.05 ± 0.05	$10.70 {\pm} 0.05$	$8.86 {\pm} 0.04$	1.7	HCT
2005 Nov 30	13.06 ± 0.02	$10.70 {\pm} 0.03$	$8.93 {\pm} 0.02$	1.6	HCT
2005 Dec 3	13.45 ± 0.01	$10.87 {\pm} 0.01$		1.9	PRL/NICMOS

^aReipurth & Aspin (2004): photometry is obtained in an aperture with radius 0.9 arcsec.

^bMcGehee et al. (2004).

^cOjha et al. (2005): IRSF photometry is obtained in an aperture with radius 7.2 arcsec.

^dUpper limit in K band is given (source is saturated in K band).

NICMOS camera is a 256×256 HgCdTe array with pixel scale of 1.0 arcsec and a FOV of 4×4 arcmin². Photometric observations were made in *JHK'* bands on 10 occasions, i.e. on 2004 March 20 and 21, April 11, October 16 and 17, December 21 and 22, 2005 November 25 and 26 and December 3 under photometric conditions. Single frame integration times were 10/20 s for *J*, 5/10 s for *H* and 2 s for *K'* band. Overall integration times were 300-360 s for *J*, 200–300 s for *H* and 200–320 s for *K'* bands. A sufficient number of dithered sky frames were also taken along with dark frames. Photometric calibration was done by using the standard stars around AS13 region (Hunt et al. 1998) and HD 40335.

The data reduction was done using IRAF software tasks. All the NIR images went through the standard pipeline reduction procedures like dark/sky-subtraction and flat-fielding. Photometric magnitudes were extracted by using the IRAF DAOPHOT/PHOT and AP-PHOT/PHOT tasks. NIR photometry of V1647 Ori obtained on 17 nights (2004 February–2005 December) with NIRCAM, TIRCAM and NICMOS cameras is listed in Table 3 along with other published results in the literature.

2.3 Optical spectroscopy

Spectroscopic observations of V1647 Ori were carried out with the HFOSC instrument at the 2-m HCT on 10 occasions beginning on 2004 February 22. The log of observations is given in Table 4. To avoid the contamination from nebular emission around McNeil's nebula, the slit was positioned on V1647 Ori in east–west direction. The spectra were obtained at resolutions of ~15 and ~7 Å in the wavelength range 5000–9100 Å. These resolutions are sufficient to resolve the H α line detected in the spectrum of V1647 Ori. The spectra were reduced using standard IRAF routines. The

Date (UT)	JD	Resolution (Å)	Exposure (s)
2004 Feb 22	245 3057.9	7	900
2004 Feb 23	245 3059.2	7	900
2004 Nov 16	245 3326.5	15	900
2004 Nov 19	245 3329.4	15	900
2005 Jan 8	245 3379.2	7	900
2005 Jan 9	245 3380.2	7	900
2005 Jan 16	245 3387.1	7	900
2005 Mar 28	245 3458.2	7	900
2005 Sept 8	245 3621.7	7	4500
2005 Sept 28	245 3642.5	7	1800

one-dimensional spectra were extracted from the bias-subtracted and flat-field corrected images using the optimal extraction method. Wavelength calibration of the spectra was done using the FeAr and FeNe lamp sources. Subsequently, the spectra were corrected for instrumental response using spectrophotometric standards (Hiltner 600, Feige 34 and G191B2B) observed on the same night and brought to a flux scale. For the nights of 2005 September 8 and 28, the multiple flux calibrated spectra were combined and scaled to a weighted mean.

3 RESULTS AND DISCUSSION

3.1 Optical and IR morphology of McNeil's Nebula

A colour composite image was constructed from the HFOSC V, $H\alpha$ and *I*-band images (V represented in blue, $H\alpha$ in green and *I* in red) obtained on 2004 February 12 and is shown in Fig. 1. The

Table 4.	Spectrosco	ic observations of	V1647 Ori at HCT.
----------	------------	--------------------	-------------------



Figure 1. $V H\alpha I$ three-colour composite image of McNeil's nebula (V: blue, H α : green and I: red) obtained with HFOSC mounted on the 2-m HCT telescope on 2004 February 12. FOV is ~10 × 10 arcmin². North is up and east is to the left-hand side. The location of V1647 Ori is marked at the centre of the image. The well-known HH objects, such as the HH 24 complex, are seen in green colour because of the H α emission. The diffuse H α emission due to HH 23, which may be driven by V1647 Ori, is marked by the box at top centre. The star marked with A is a reference star (see the text).

location of V1647 Ori is marked at the centre of the image. The wellknown Herbig–Haro (HH) objects, such as the HH 24 complex, are seen in green colour because of the H α emission. The diffuse H α emission due to HH 23 is highlighted by the box at top centre. The star associated with McNeil's nebula was proposed as the exciting source of HH 23 (Eisloffel & Mundt 1997; Kastner et al. 2004). A bright knot HH 22 is marked by an arrow, which is seen as a 'smoky cloud' covering a part of the new object.

The enlarged view of McNeil's nebula region is shown in Fig. 2. Fig. 2(a) shows the central region of nebula in optical wavelengths. The right-hand panel (Fig. 2b) shows the *JHK* composite colour image (*J*: blue, *H*: green and *K*: red) of McNeil's nebula observed on 2004 October 18. The cometary IR nebula is clearly seen extending towards north from the NIR source (V1647 Ori) and has a cavity structure with possibly two rims extending towards north-east and north-west (see also Ojha et al. 2005). The north-east rim is brighter and sharp, while the north-west rim is diffuse and is not seen clearly. The north-east rim can be traced out to ~30 arcsec or ~12 000 au at a surface brightness of $K \sim 18.0$ mag arcsec⁻² (where we assume that the distance to McNeil's nebula is 400 pc; Anthony-Twarog 1982). Such kind of morphology can probably be seen when the eruption has cleared out some of the dust and gas in the envelope surrounding the young star, allowing light to escape and to illuminate a coneshaped cavity carved out by previous eruptions (Reipurth & Aspin 2004).

In Fig. 3(a), we show a contour diagram of the region around V1647 Ori, as seen in our *K*-band image obtained on 2004 October 18. The star is surrounded by a compact reflection nebulosity, and there is a southwards continuation of the extended nebula in NIR, which is demonstrated in the PSF-subtracted *K*-band image of Fig. 3(b). This was confirmed by comparing the PSF of the NIR central source (V1647 Ori) with the PSF of the star located about 30 arcsec south-west of V1647 Ori (Fig. 3b). We find that the compact IR nebula that encloses all of the nebular structure, including the two rims, has a diameter of about 60 arcsec.

An optical/IR comparison of the region surrounding McNeil's nebula (see Figs 2 and 4) shows that the position of the bright IR source is coincident with that of an optical source at the apex of the nebula that is illuminating the fan-shaped cloud of gas, or nebula (Reipurth & Aspin 2004). The optical nebula is more widely and predominantly extended to the north, whereas the IR nebula is relatively confined, but definitely extended to the south too. The



Figure 2. (a) Enlarged view of $V H \alpha I$ three-colour composite image of McNeil's nebula (see Fig. 1). (b) JHK three-colour composite image of McNeil's nebula (J: blue, H: green and K: red) obtained with NIRCAM mounted on the 2-m HCT telescope on 2004 October 18. North is up and east is to the left.



Figure 3. (a) Compact nebula seen around the illuminating star of McNeil's nebula in *K*-band image obtained on 2004 October 18. The lowest contour and the contour interval are 18.0 and 0.5 mag arcsec⁻², respectively. (b) PSF-subtracted image of McNeil's nebula region in the *K* band displayed on a logarithmic scale. North is up and east is to the left-hand side. The locations of the NIR central source (V1647 Ori) and the star \sim 30 arcsec south-west of V1647 Ori are indicated by filled triangles.

greenest portion of the optical nebula is not seen in IR including the knot HH 22 to the north-east and the green patch to the north (see Figs 1, 2 and 4). There is a clear colour gradient from north to south (H - K colour varies from 0.53 to 1.33 from north to south), as well as from inside to outside of the cavity. The gradient is very steep near the central star, and the southern extension of the nebula is seen only at IR. This large colour gradient and the absence of optical nebula to the south are suggestive of a large-scale disc-like structure (or envelope) surrounding the central source that hides the southern nebula in optical. This is often the case for YSOs associated with a cometary nebula (e.g. Tamura et al. 1991).

3.2 Optical/IR variability of V1647 Ori

Optical and NIR photometric results of V1647 Ori obtained on 27 nights (2004 February–2005 December) are listed in Tables 2 and 3.

The optical photometry shows a general decline of brightness in V1647 Ori both in the beginning and at later part of our observations with an intermediate brightening phase. In our data set, V1647 Ori is brightest on 2004 October 18 (see Table 2). McNeil's nebula itself has now faded considerably (see Table 2). A significant variation in the source brightness ($\Delta V = 3.43$, $\Delta R = 3.36$ and $\Delta I = 3.14$ mag) can be seen during a period of about 21 months (2004



Figure 4. NIR and optical images of the region surrounding McNeil's nebula. HCT K-band image (grey colour) is overlaid with contours from the optical H α image obtained on 2004 October 18. North is up and east is to the left-hand side.

February–2005 November; Table 2). We can also see a considerable variation in the source brightness ($\Delta V = 0.78$, $\Delta R = 0.44$ and $\Delta I = 0.21$ mag) during a period of one month (2004 October–November; Table 2). The difference in magnitudes is much larger than the photo-

metric errors (≤ 0.05 mag). For comparison, the *BVRI* magnitudes of the reference star A [$(\alpha, \delta)_{2000} = 05^{h}46^{m}22^{s}4, -00^{\circ}03'37'';$ marked in Fig. 1] are listed in Table 5 along with other results from the literature, which match quite well with our measurements. As

 Table 5. BVRI photometry of reference star A at HCT.

Reference star A	В	V	R	Ι	Date (UT)
USNO-A2.0 0825-01669040		15.69 ± 0.01	14.93±0.02	14.24 ± 0.02	2004 Feb 14 ^a
	$16.89 {\pm} 0.01$	$15.65 {\pm} 0.01$	$14.88 {\pm} 0.02$	14.23 ± 0.01	2004 Feb 22 ^a
	$16.93 {\pm} 0.04$	$15.67 {\pm} 0.03$	$14.88 {\pm} 0.03$	$14.20 {\pm} 0.03$	2004 Oct 18 ^a
	$16.93 {\pm} 0.02$	$15.63 {\pm} 0.02$	$14.88 {\pm} 0.01$	14.23 ± 0.02	2004 Nov 17 ^a
	$16.92 {\pm} 0.02$	$15.68 {\pm} 0.01$	$14.86 {\pm} 0.02$	$14.23 {\pm} 0.01$	2004 Dec 7 ^a
		$15.65 {\pm} 0.01$	$14.89 {\pm} 0.01$		2005 Sept 26 ^a
	$16.94 {\pm} 0.01$	$15.66 {\pm} 0.01$	$14.89 {\pm} 0.01$		2005 Sept 29 ^a
		$15.67 {\pm} 0.01$	$14.91 {\pm} 0.01$	$14.23 {\pm} 0.01$	2005 Nov 28 ^a
		$15.69 {\pm} 0.01$	$14.92 {\pm} 0.01$	$14.25 {\pm} 0.01$	2005 Nov 30 ^a
	$16.92 {\pm} 0.01$	$15.66 {\pm} 0.01$	$14.87 {\pm} 0.01$	14.21 ± 0.01	2004 Feb ^b
		15.65	14.91	14.26	2004 Apr 24 ^c
		$15.66 {\pm} 0.02$	$14.89 {\pm} 0.02$	$14.23 {\pm} 0.05$	2004 Aug–Oct ^d

^aFrom HCT.

 $^b{\rm From}$ Henden's (2004) table of photometric measurements around McNeil's nebula.

^cFrom Kun et al. (2004).

^dFrom Semkov (2004).



Figure 5. The optical light curve of V1647 Ori in *R* band. The differential magnitudes are relative to a single comparison star (see the text), which appears constant to within 0.05 mag. The filled circles show our measurements (2004 February–2005 November). The cross symbols show the photometric measurements from McGehee et al. (2004). The filled triangles are the *R*-band photometric observations by Semkov (2004) during the period 2004 August–October.

can be seen in Table 5, we do not see any significant variation (>0.05 mag, which is the estimated photometric error) in the magnitudes of the reference star A. Hence, the brightness variation seen in V1647 Ori during a period of one month (2004 October–November) is real. It is also seen that the variability is larger at shorter wavelengths compared to the longer ones (Table 2).

The differential light curve (2004 February–2005 November) of V1647 Ori in *R* band is shown in Fig. 5. Semkov (2004) reported the *VRI* photometric observations of V1647 Ori and found that the source varied with amplitude of 0.5 mag during the period 2004 August 18–October 3. In Fig. 5, we have also plotted the *R*-band light curve from Semkov (2004). The optical light curve shows a variable nature of V1647 Ori. After a significant increase in the brightness around 2004 October 18, a general decline in the source brightness can be seen in recent observations. Over the one-and-ahalf year that we followed the star (2004 February–2005 November), the brightness has decreased by more than 3 mag in *R* band. The shapes of the light curves are similar in other optical bands.

A comparison of the *JHK* magnitudes (see Table 3) of V1647 Ori shows that the source has varied over the time of observation between 2MASS (1998 October) and HCT/PRL (2005 November/December). We also see a significant variation in the source brightness ($\Delta J = 0.13$, $\Delta H = 0.26$ and $\Delta K = 0.14$ mag) during a period of one week (2004 October 12–18). A similar kind of weekly variation in NIR magnitudes of V1647 Ori during the early phase of outburst (2004 February 23–29) was reported by Ojha, Kusakabe & Tamura (2004) and Ojha et al. (2005). The short time variability in optical and NIR wavelengths in V1647 Ori was also noted by Walter et al. (2004). They found a general decline in brightness of about 0.3–0.4 mag during 87 d (2004 February 11–May 7).

Fig. 6 shows a J - H/H - K colour–colour (CC) diagram. We note changes of the IR colours of V1647 Ori from before eruption (1998) to after the eruption (2004–2005). A general movement along a reddening vector can be seen during 1998–2005. The source had moved towards the tip of the locus of classical T Tauri stars (Meyer, Calvet & Hillenbrand 1997) during 2004. This suggests that V1647 Ori has NIR colour similar to a dereddened T Tauri star. The J - Hand H - K colours indicate that the circumstellar matter of $A_V \sim 8$ mag was probably cleared in the eruption in less than 6 yr since the 2MASS observations. This scenario is consistent with the present IR morphology of the McNeil's nebula region (see Section 3.1). There are colour changes between the most recent measurements (2005 November) and those close to peak brightness (2004). The



Figure 6. J - H/H - K CC diagram showing the location of V1647 Ori as observed with 2MASS (filled circle) on 1998 October 7; with IRSF (filled squares) on 2004 February 23–29; with Gemini (filled triangle) on 2004 February 3; with USNO (asterisks) on 2004 February 11 and April 12; with TIRCAM (open circle) on 2004 October 12; with HCT (plus symbols) on 2004 October–2005 November and with NICMOS (open triangles) on 2004 March–December. The left small arrow shows the limit on H - K colour on 2004 November 19 (HCT) due to the saturation in K band. The sequences of field dwarfs (solid curve) and giants (thick dashed curve) are from Bessell & Brett (1988). The dotted line represents the locus of T Tauri stars (Meyer et al. 1997). The dashed straight lines represent the reddening vectors (Rieke & Lebofsky 1985). The crosses on the dashed lines are separated by $A_V = 5$ mag.

source has now become redder and fainter and moved back along a reddening vector in the CC diagram.

The brightness increase in all the optical and NIR bands near the outburst peak (early 2004) tells that the accretion disc luminosity that depends directly on the accretion rate must have gone up. As the enhanced accretion has decreased over a period of about 2 yr, the disc luminosity too has decreased. So, it may be predominantly the stellar plus the original disc excess as seen in the IR. The position in the CC diagram is therefore a combination of extinction and thermal dust emission. As the object is embedded in dust and gas, there will also be some reflection effects.

3.3 Spectral features

The optical spectra of V1647 Ori obtained with HCT are shown in Fig. 7. The spectra show a very red continuum with a number of emission features. The H α and Ca II IR triplet lines are in emission. The presence of the Ca II IR triplet in emission confirms that V1647 Ori is a pre-main-sequence star. We also identified a few lines of the Paschen series of hydrogen at low signal-to-noise ratio level (e.g. Pa 23, 20, 19, 18, 17, 12, 11 and 10) in our deep spectra of 2005 September 8 and 28 in the wavelength range 8340–9020 Å. In our spectra, the OI λ 7773 line is seen in absorption. The detection of OI λ 7773 line indicates that the optical light probably originates in the low-gravity, hot photosphere of the inner disc (Walter et al. 2004). Since our spectra are not corrected for telluric features, we see the



Figure 7. Optical spectra of V1647 Ori obtained with HFOSC at the 2-m HCT telescope in the wavelength range 5500–9100 Å.

absorptions at about 7600 and 8230 Å due to atmospheric O2-A and H_2Oz bands, respectively. In the spectrum taken on 2005 September 8, we detected the Mg I line at 8807 Å and Fe I at 8824 Å rather marginally. Table 6 shows the equivalent width (EW) measurements of H α , O I λ 7773 and Ca II IR triplet from HCT spectra. We estimate the measurement uncertainties in the EW to be of the order of \sim 2 per cent. The EW of the Ca II triplet is nearly same, which is in contrast with the results of Walter et al. (2004), where the ratio of the triplet EWs was shown to be 2:2:1. Interestingly, Walter et al. had used a resolution similar to that of ours. It must be noted here that in our data, the Ca II triplet lines have not been de-blended for the H I Pa 13, 15 and 16 lines. Since the other Paschen lines are not seen as prominently as the towering Ca II triplet, we do not expect significant contamination in Ca II triplet by Pa 13, 15 and 16 lines.

Due to the somewhat low/medium resolution, we cannot resolve the actual FWHMs and peak heights of the components of the triplet which would have probably thrown some light on the possible optical depth variations within the Ca II emission regions. In general, the Ca II triplet arises from optically thick regions of electron densities of about 10^9-10^{10} cm⁻³, with temperatures between about 4000 and 10 000 K (Ferland & Persson 1989). Following the arguments of Hamann & Persson (1992) for T Tauri stars, we infer from the nearly equal EWs of Ca II triplet lines that the collisional decay is a more dominant process than radiative decay. This in turn implies that the product of optical depth and electron density needs to be more than or equal to about 10^{13} cm⁻³ (see Hamann & Persson 1992). Coupling this argument with the absence of the [Ca II] lines at ~7300 Å in the HCT spectra, we may conclude that the density of the Ca II region is very large and is optically very thick.

On one occasion, i.e. on 2005 September 8, the HCT spectra clearly show the Ca II forbidden doublet at 8912/8927 Å, while the other forbidden doublet at 7291/7323 Å is still absent. These forbidden lines arise in optically thin regions in comparison to the Ca II permitted lines, otherwise they will be de-excited by collisions. The doublet 8912/8927 Å is about 8.4 eV from the ground state, while the doublet 7291/7323 Å is only 1.7 eV (e.g. Lang 1978). This is probably indicative of the high temperature of the chromosphere of the T Tauri type star that we are monitoring. This region must also have a lower density than the triplet region in order not to get de-excited by collisions. Hamann & Persson (1992) noted that [Ca II] lines occur rarely in T Tauri stars, and in some cases where they did, they showed blue-shifted profiles. Since the HCT spectrum is rather spiky in the region near the doublet, it is difficult to say if the line peak is Doppler-shifted.

The HCT spectra show prominent broad H α line on all occasions with varying EW (see Fig. 8). The line seems to arise from regions having large-scale turbulent velocities in comparison to the narrow lines of Ca II. The FWHMs of the H α emission lines (after



Figure 8. The H α profiles of V1647 Ori seen in the spectra obtained with HFOSC between 2004 February 22 and 2005 September 28. The H α line shows a pronounced blue-shifted P Cygni profile in the spectra taken in the early phase of outburst (2004 February 22 and 23).

Table 6. H α (λ 6563), OI (λ 7773) and Ca II IR triplet ($\lambda\lambda\lambda$ 8498, 8542 and 8662) measurements from HCT spectra.

Date (UT)	W_{λ} (6563)	$W_{\lambda}~(O{\scriptscriptstyle I}~7773)$	W_{λ} (8498)	W_{λ} (8542)	W_{λ} (8662)
2004 Feb 22	-34.59	3.82	-8.67	-9.56	-7.72
2004 Feb 23	-33.62	2.63	-7.67	-7.84	-6.58
2004 Nov 16	-27.25	5.41	-12.91	-10.14	-12.39
2004 Nov 19	-41.73	4.02	-15.62	-14.79	-12.16
2005 Jan 8	-28.81	2.81	-9.70	-9.08	-7.20
2005 Jan 9	-27.72	2.74	-9.34	-9.09	-7.17
2005 Jan 16	-32.54	2.69	-9.66	-10.17	-8.68
2005 Mar 28	-20.34	2.98	-10.66	-9.62	-11.23
2005 Sept 8	-26.16	2.58	-16.73	-19.10	-18.43
2005 Sept 28 ^a	-40.75		-18.43	-21.11	-19.30

^aO_I line is not clearly seen.

correcting for instrumental broadening by using quadrature formula) show variations ranging from 300 to 600 km s⁻¹. Such large widths are quite common in T Tauri stars (e.g. Hamann & Persson 1992). There appears to be a positive correlation between the H α and the Ca II EWs, indicating that the two emitting regions are responsive to each other (Hamann & Persson 1992). We do not see this correlation in O I λ 7773 line (in absorption) which occurs in rather cooler regions in comparison.

The strong H α is in emission, with a P Cygni absorption feature seen in the early phase of outburst (2004 February 22 and 23; see Fig. 8), first reported by Reipurth & Aspin (2004). Similar P Cygni absorption features are also seen in the Ca II IR triplet in early outburst spectra of V1647 Ori. The pronounced P Cygni profile seen in H α is likely to be formed in a strong wind that has sufficient optical depth to produce the deep blue-shifted absorption trough. The welldefined blue edge of the absorption trough indicates a wind velocity



Figure 9. The time variation of the H α equivalent width. The filled circles denote our measurements with HCT (2004 February 22–2005 September 28). The filled triangles show the data points (2004 February 13–April 17) from Walter et al. (2004). While the data from Walter et al. (2004) are suggestive of an oscillatory or periodic variation, we probably also see such a variation in our data set.

of 420–460 km s⁻¹ [Fig. 8; see also Maheswar & Bhatt (2004)]. This implies significant mass loss in a powerful wind. We note a significant decrease in the depth and the extent of this strong absorption component over the period of our observations (Fig. 8), as also reported by Aspin & Reipurth (2005). This suggests weakening of the powerful wind.

Fig. 9 shows the time variation of the H α EW over a period of about 19 months, which includes data from Walter et al. (2004). The data from Walter et al. (2004) indicate an oscillatory or periodic variation of the H α EW. We also probably see such a variation in our data set. Power spectrum analysis indicates the period of oscillations to be ~40 d. This variation may probably be caused by temporal variations in the wind/accretion flow or by the rotational period of V1647 Ori. However, poor time sampling prevents us from making definitive conclusions about the periodic variation in H α EW.

From Table 6, we note that the strength of the H α emission line has increased, whereas the OI λ 7773 line in absorption appears quite strong in the early phase of the outburst (2004 February– November) and has become weaker (in fact it is not seen clearly on 2005 September 28). Compared to the early outburst stages, Ca II IR triplet lines also show significant strengthening. Hamann & Persson (1992) have suggested that the OI λ 7773 absorptions are entirely due to warm gas in the envelope, and the presence of this line suggests either a rotating disc or a highly turbulent envelope. If it were an outburst, then turbulence in the envelope gas is expected initially which in turn gives stronger OI λ 7773 line. But as the time progresses, the turbulence may die down and the line becomes weaker. Hence, variability in the OI λ 7773 line indeed indicates an outburst activity in the source.

3.4 Possible EXor event?

Two classes of erupting low-mass YSOs have emerged: FUors, which display outbursts of ~4 mag and more that last several decades and EXors, which show smaller outbursts ($\Delta m \sim 2-3$ mag) that last from a few months to a few years and may occur repeatedly (Herbig 1977; Bell et al. 1995; Hartmann 1998). V1647 Ori had brightened by about 3 mag in *JHK*, relative to the 1998 2MASS measurements and is probably returning to its pre-outburst state. It is therefore possible that we had witnessed an EX Orionis kind of behaviour, since the source has about the correct amplitude (Reipurth & Aspin 2004; Ojha et al. 2005). The time elapsed since the outburst of V1647 Ori is more than 2 yr now. During this period, our optical data show a significant variation in the source brightness (see Table 2); the latest optical images show that V1647 Ori has faded

by more than 3 mag since February 2004. We also note a significant weekly and monthly variations in the brightness of V1647 Ori over the period of our observations (2004 February–2005 November/December; see Tables 2 and 3). Similar kind of variations are also seen from the light curves of Walter et al. (2004). This seems to suggest that we are probably witnessing a sporadic or episodic outburst (e.g. Herbig et al. 2001), which may indicate an EXor kind of eruption of V1647 Ori. However, further photometric monitoring of the object is required to settle the class of eruptive variables to which V1647 Ori belongs.

V1647 Ori has now started a very rapid fading. The object may indeed be returning to its pre-outburst characteristics, though in the NIR it is still 1.3–1.4 mag brighter than the 2MASS (preoutburst) brightness. At both optical and NIR, the object was bluer and brighter near the outburst peak brightness and is now redder and fainter. Such a behaviour would be expected if during the outburst a disc, hotter than the stellar photosphere, dominated the luminosity signifying high accretion rate. With the decline in the accretion, the luminosity is now dominated by the star (cool late type) and a less pronounced disc still emits some excess IR. This also shows that the outburst is essentially caused by accretion enhancement.

4 CONCLUSIONS

In this paper, we have presented long-term, post-outburst photometric and spectroscopic monitoring of V1647 Ori. Following conclusions are drawn from our study.

(i) The optical data show that V1647 Ori has faded by more than 3 mag since February 2004. We also see significant variations in the brightness of V1647 Ori within a period of one week ($\Delta m \sim 0.3$ mag) and one month ($\Delta m \sim 0.8$ mag). This shows the eruptive behaviour of McNeil's nebula.

(ii) McNeil's nebula has a cavity structure with possibly two rims as seen in NIR wavelengths. This kind of morphology might have been produced by the eruption that has cleared out a part of the envelope surrounding the young star. This scenario is supported by the J - H and H - K colours (near the outburst peak brightness) of V1647 Ori which indicate that the circumstellar matter of $A_V \sim$ 8 mag was probably cleared in the eruption.

(iii) An optical/IR comparison of McNeil's nebula shows that the optical nebula is more widely and predominantly extended to the north, while the IR nebula is relatively confined, but definitely extended, to the south, too. This colour gradient of the nebula is most likely due to the presence of a large-scale disc-like structure (or envelope) around the young star.

(iv) The optical spectra show H α line strongly in emission and display a pronounced P Cygni profile, with an absorption trough reaching a velocity of ~460 km s⁻¹ during the early phase of outburst. The depth and the extent of the strong absorption component have decreased significantly over the period of our observations, suggesting that the powerful wind is weakening. Variation in H α line strength on the time-scale of ~1 d is also seen.

(v) The presence of strong emission lines, relatively fast fading rate and the low accretion rate $(10^{-5} M_{\odot} \text{ yr}^{-1}; \text{ Muzerolle et al.} 2005)$ of V1647 Ori suggest probably the evidence for a new EXor event.

ACKNOWLEDGMENTS

The observations reported in this paper were obtained using the 2-m HCT at IAO Hanle, the high-altitude station of the Indian Institute

of Astrophysics, Bangalore. We thank the staff at IAO and at the remote control station at CREST, Hosakote, for assistance during the observations. It is a pleasure to thank the staff at Mount Abu IR observatory for their support during the observations. We especially thank the referee, Dr William Sherry, for the critical reading of our manuscript.

REFERENCES

- Ábrahám P., Kóspál Á., Csizmadia Sz., Moór A., Kun M., Stringfellow G., 2004, A&A, 419, L39
- Andrews S. M., Rothberg B., Simon T., 2004, ApJ, 610, L45
- Anthony-Twarog B. J., 1982, AJ, 87, 1213
- Aspin C., Reipurth B., 2005, IAU Circ., 8600
- Bell K. R., Lin D. N. C., Hartmann L. W., Kenyon S. J., 1995, ApJ, 444, 376
- Bessell M. S., Brett J. M., 1988, PASP, 100, 1134
- Briceño C. et al., 2004, ApJ, 606, L123
- Eisloffel J., Mundt R., 1997, AJ, 114, 280
- Ferland G. J., Persson S. E., 1989, ApJ, 347, 656
- Grosso N., Kastner J. H., Ozawa H., Richmond M., Simon T., Weintraub D. A., Hamaguchi K., Frank A., 2005, A&A, 438, 159
- Hamann F., Persson S. E., 1992, ApJS, 82, 247
- Hartmann L., 1998, in Accretion Processes in Star Formation. Cambridge Univ. Press, Cambridge
- Henden A., 2004, http://spiff.rit.edu/classes/phys440/lectures/newstar/mcneil.dat
- Herbig G. H., 1977, ApJ, 217, 693
- Herbig G. H., Aspin C., Gilmore A. C., Imhoff C. L., Jones A. F., 2001, PASP, 113, 1547
- Hunt L. K., Mannucci F., Testi L., Migliorini S., Stanga R. M., Baffa C., Lisi F., Vanzi L., 1998, AJ, 115, 2594

- Kastner J. H. et al., 2004, Nat, 430, 429
- Kóspál Á., Ábrahám P., Acosta-Pulido J., Csizmadia S., Eredics M., Kun M., Racz M., 2005, Inf. Bull. Var. Stars, 5661, 1
- Kun M. et al., 2004, A&A, submitted (astro-ph/0408432)
- Landolt A. U., 1992, AJ, 104, 340
- Lang K. R., 1978, Astrophysical Formulae. Springer Verlag, Berlin, p. 141
- Maheswar G., Bhatt H. C., 2004, IAU Circ., 8295
- McGehee P. M., Smith J. A., Henden A. A., Richmond M. W., Knapp G. R., Finkbeiner D. P., Ivezić Ž., Brinkmann J., 2004, ApJ, 616, 1058 McNeil J. W., 2004, IAU Circ., 8284
- Meyer M. R., Calvet N., Hillenbrand L. A., 1997, AJ, 114, 288
- Muzerolle J., Megeath S. T., Flaherty K. M., Gordon K. D., Rieke G. H., Young E. T., Lada C. J., 2005, ApJ, 620, L107
- Ojha D. K., Ghosh S. K., Verma R. P., Chakraborti A., Anandarao B. G., 2002, Bull. Astron. Soc. India, 30, 827
- Ojha D. K., Kusakabe N., Tamura M., 2004, IAU Circ., 8306
- Ojha D. K., 2005, PASJ, 57, 203
- Reipurth B., Aspin C., 2004, ApJ, 606, L119
- Rieke G. H., Lebofsky M. J., 1985, ApJ, 288, 618
- Samus N. N., 2004, IAU Circ., 8354
- Semkov E. H., 2004, Inf. Bull. Var. Stars, 5578, 1
- Stetson P. B., 1987, PASP, 99, 191
- Tamura M., Gatley I., Waller W., Werner M. W., 1991, ApJ, 374, L25
- Vacca W. D., Cushing M. C., Simon T., 2004, ApJ, 609, L29
- Vig S., Ghosh S. K., Ojha D. K., Kulkarni V. K., 2004, IAU Circ., 8354
- Vig S., Ghosh S. K., Kulkarni V. K., Ojha D. K., 2006, A&A, 446, 1021
- Walter F. M., Stringfellow G. S., Sherry W. H., Pollatou A. F., 2004, AJ, 128, 1872

This paper has been typeset from a TFX/LATFX file prepared by the author.