

THE ROTATIONAL PERIOD OF 1406 KOMPPA

Russell I. Durkee
Shed of Science Observatory
Minneapolis, MN, 55410 USA
shedofscience@earthlink.net

Daniel A. Klinglesmith III
Curtis Warren
Etsorn Campus Observatory
New Mexico Tech
Socorro, NM, 87801 USA

John W. Briggs
HUT Observatory
Eagle, CO, 81631 USA

(Received: 7 October)

Our observations of main-belt asteroid 1406 Komppa indicate a synodic period of $P = 3.508 \pm 0.002$ h, $A = 0.16 \pm 0.05$ mag.

Observations of 1406 Komppa were taken over 7 nights in 2011 August and September. The Shed of Science used a 0.35-m Schmidt Cassegrain (SCT) with an SBIG ST-10XE CCD camera working at f/8.5, resulting in a scale of 0.94 arcsec/pixel. Exposures were made through a Celestron UHC LPR filter. The Etsorn Campus Observatory of New Mexico Institute of Mining and Technology has two identical 0.35-m SCT operating at f/11 with SBIG STL-1001E CCD cameras resulting in an image scale of 1.25 arcsec/pixel. HUT Observatory has a 16-inch f/8 Ritchey-Chretien reflector by DFM Engineering. For these observations we used an Apogee Alta model U47 CCD with a Bessell R filter. The exposures were binned 2x2 for an effective image scale of 1.65 arcsec/pixel.

All images were dark and flat field corrected. Images were measured using *MPO Canopus* (Bdw Publishing) with a differential photometry technique. The *MPO Canopus* Comp Star Selector was used to link sessions. The data were light-time corrected. Period analysis was also done with *MPO Canopus*, incorporating the Fourier analysis algorithm developed by Harris (Harris *et al.* 1989).

Behrend (2011) reported a period near 7.0 h, or twice the period we are reporting here. Our current results are in close agreement with earlier work on this object by Polishhook (2009).


Acknowledgements

Partial funding for work at the Shed of Science is provided by Gene Shoemaker NEO Grants from the Planetary Society. Work at the HUT Observatory is supported by the Mittelman Foundation.

References

- Behrend, R. (2011). "Courbes de rotation d'astéroïdes et de comètes, CdR." <http://obswww.unige.ch/~behrend/page4cou.html#001406>
- Harris, A.W., Young, J.W., Bowell, E., Martin, L.J., Millis, R.L., Poutanen, M., Scaltriti, F., Zappala, V., Schober, H.J., Debehogne, H., and Zeigler, K.W. (1989). "Photoelectric Observations of Asteroids 3, 24, 60, 261, and 863." *Icarus* **77**, 171-186.

Polishhook, D. (2009). "Lightcurves and Spin-Periods from the Wise Observatory: 2008 August – 2009 March." *Minor Planet Bull.* **36**, 104-107.



ROTATIONAL PERIOD DETERMINATION FOR 1820 LOHMANN

Luis E. Martinez
Lenomiya Observatory (H13)
515 E. Barrus Place
Casa Grande, AZ, USA 85122
lenomiyaobs@cox.net

(Received: 8 October)

Observations of main-belt minor planet 1820 Lohmann were undertaken by Lenomiya Observatory during 2011 August-September. Data analysis found a period of 14.048 ± 0.001 h with amplitude 0.48 ± 0.01 mag.

1820 Lohmann is a main-belt asteroid, discovered in 1949 by K. Reinmuth in Heidelberg and named in honor of Werber Lohmann (1911-1983), an astronomer at Heidelberg (Schmadel, 2003). The asteroid was selected from Collaborative Asteroid Lightcurve Link (CALL) site's Lightcurve Targets list (Warner, 2011). A search of the Asteroid Lightcurve Database and other sources does not reveal any previously reported lightcurve results for 1820 Lohmann.

Observations of the asteroid were made at the Lenomiya Observatory with a Celestron CPC-1100 0.28-m Schmidt-Cassegrain (SCT) working at f/6.3 using a focal reducer. Images were guided and unfiltered using a Santa Barbara Instruments Group ST8XME CCD camera operating at -12° C. The images were binned 2x2, which resulted in an effective array of 765x510 18-micron pixels and scale of 1.92 arcseconds/pixel. The 1,442 images were exposed in the range of 60 s to 100 s, depending on atmospheric conditions, in order to maximize SNR. They were calibrated using *CCDSOFT* version 5.00.205 and measured using *MPO Canopus* version 10.4.0.4 (Warner, 2011).

Analysis of the data disclosed a bimodal lightcurve with a period of 14.048 ± 0.001 h with an amplitude of 0.48 ± 0.01 mag.

Individual sessions spanned the complex section of the curve ruling out any trimodal or monomodal fit. Due to weather problems, only about 85% phase coverage could be obtained. However, I believe this period to be secure because the only shape that can produce a lightcurve with an amplitude as large as 0.40 magnitudes is an elongated body that features two maxima and minima per cycle that are nearly symmetrically placed.

Acknowledgments

The author wishes to express gratitude to Frederick Pilcher for his support and helpful suggestions on the *MPO Canopus* software and period determination.

References


Schmadel, L.D. (2003). Dictionary of minor planets. (5th ed., p. 146). New York: Springer.

Warner, B.D. (2011). "Potential Lightcurve Targets 2011 July 1 – September 30."

http://www.minorplanet.info/CALL/targets_2011_Q3.htm

Warner, B.D. (2011). MPO Software, Canopus version 10.4.0.4 Bdw Publishing.

<http://minorplanetobserver.com>



ASTEROID LIGHTCURVE ANALYSIS AT THE OAKLEY SOUTHERN SKY OBSERVATORY: 2011 APRIL–MAY

Richard Ditteon

Rose-Hulman Institute of Technology
5500 Wabash Ave., CM 171, Terre Haute, IN 47803
ditteon@rose-hulman.edu

Lauren Horn

Dupo High School, East Charondelet, Illinois

Amy Kamperman

Loveland High School, Loveland, Ohio

Bradley Vorjohan

Maryville High School, Maryville, Tennessee

Elaine Kirkpatrick

Rose-Hulman Institute of Technology

(Received: 15 October)

Photometric data for 19 asteroids were collected over 21 nights during 2011 April and May at the Oakley Southern Sky Observatory. The asteroids observed included: 518 Halawe, 828 Lindemannia, 999 Zachia, 1305 Pongola, 1359 Prieska, 1858 Lobachevskij, 2008 Konstitutsiya, 2141 Simferopol, 2271 Kiso, 2691 Sersic, 2776 Baikal, 2841 Puijo, 2931 Mayakovsky, 3031 Houston, 3044 Saltykov, 3181 Ahnert, 4362 Carlisle, and 5092 Manara.

Nineteen asteroids were observed from the Oakley Southern Sky Observatory near Coonabarabran, New South Wales, Australia on the nights of 2011 April 22-30, May 1, May 3-7, May 9-10, and May 25-28. From these images, we were able to find lightcurves for eight asteroids. Of those eight, six were previously unrecorded results. Our result for one asteroid was reasonably close to the already published period, but the other was significantly different.

The selection of asteroids was based on their sky position approximately one hour after sunset. Asteroids without previously published lightcurves were given higher priority than asteroids with known periods. Asteroids with uncertain periods were also selected with the hope that we would be able to improve earlier results. We used an RC Optical, 20-inch Ritchey-Chretien optical tube assembly mounted on a Paramount ME. The camera was a Santa Barbara Instrument Group STL-1001E with a clear filter. The image scale was 1.2 arcseconds per pixel at f/8.3. Calibration of the images was done using master twilight flats, darks, and bias frames. All calibration frames were created using *CCDSofr*. *MPO Canopus* was used to measure the processed images and complete the period analysis of the asteroids' lightcurves.

As far as we are aware, these are the first reported lightcurve and period determinations for asteroids 518 Halawe, 2008 Konstitutsiya, 2141 Simferopol, 2931 Mayakovsky, 3031 Houston, and 3248 Farinella. Our data for 828 Lindemannia, 999 Zachia, 1359 Prieska, 2271 Kiso, 2691 Sersic, 2776 Baikal, 2841 Puijo, 3044 Saltykov, 3181 Ahnert, 4362 Carlisle, and 5092 Manara were too noisy to determine their periods. Supporting lightcurve plots are included. Results from all of the asteroids are listed in the table below. Additional comments have been included as needed.

999 Zachia. We were unable to confirm the period of 22.77 ± 0.03 h found by Warner (2000).

1305 Pongola. Our results are within experimental uncertainty of the period of 8.03 ± 0.10 h found by Binzel (1987).

1858 Lobachevskij. Our current results are not compatible with the period of 7.00 ± 0.01 h reported by Ditteon (2002). However, the data from 2002 were reexamined and found to fit a period of 5.435 ± 0.009 h, which is close to our new value. In addition to the plot of the current data, a plot of the reexamined data is also included.

2841 Puijo. We were unable to confirm the period of 3.545 ± 0.005 h found by Warner (2004).