

References

Harris, A.W., Young, J.W., Scaltriti, F., Zappala, V. (1984). "Lightcurves and phase relations of the asteroids 82 Alkmene and 444 Gyptis." *Icarus* **57**, 251-258.

Harris, A.W., Young, J.W., Bowell, E., Martin, L.J., Millis, R.L., Poutanen, M., Scaltriti, F., Zappala, V., Schober, H.J., Debehogne, H., Zeigler, K.W. (1989). "Photoelectric Observations of Asteroids 3, 24, 60, 261, and 863." *Icarus* **77**, 171-186.

Pravec, P., Wolf, M., Sarounova, L. (1998). "Lightcurves of 26 Near-Earth Asteroids." *Icarus* **136**, 124-153.

Warner, B.D., Harris, A.W., Pravec, P. (2009). "The Asteroid Lightcurve Database." *Icarus* **202**, 134-146. Updated 2016 Dec. <http://www.minorplanet.info/lightcurvedatabase.html>

Warner, B.D. (2015). "Near-Earth Asteroid Lightcurve Analysis at CS3-Palmer Divide Station: 2014 June-October." *Minor Planet Bull.* **42**, 41-53.

Waszczak, A., Chang, C.-K., Ofek, E.O., Laher, R., Masci, F., Levitan, D., Surace, J., Cheng, Y.-C., Ip, W.-H., Kinoshita, D., Helou, G., Prince, T.A., Kulkarni, S. (2015). "Asteroid Light Curves from the Palomar Transient Factory Survey: Rotation Periods and Phase Functions from Sparse Photometry." *Astron. J.* **150**, A75.

Zappala, V., Cellini, A., Barucci, A.M., Fulchignoni, M., Lupishko, D.E. (1990). "An analysis of the amplitude-phase relationship among asteroids." *Astron. Astrophys.* **231**, 548-560.

CCD LIGHTCURVES FOR MAIN-BELT ASTEROIDS 423 DIOTIMA AND 925 ALPHONSINA


Kevin B. Alton
UnderOak Observatory
70 Summit Ave.
Cedar Knolls, NJ 07927
kbulton@optonline.net

(Received: 2017 Mar 13)

Fourier analysis of a new CCD-derived lightcurves found synodic periods for 423 Diotima (4.775 ± 0.001 h) and 925 Alphonsina (7.879 ± 0.001 h).

The instrument used at UnderOak Observatory (UO) for this investigation was a 0.28-m SCT equipped with an SBIG ST-8XME thermoelectrically-cooled CCD camera. Image calibration/registration procedures employed at UO have been published elsewhere (Alton, 2013). Data reduction with *MPO Canopus* (Warner, 2015) used at least four non-varying comparison stars in the same FOV to generate lightcurves by differential aperture photometry. Data were light-time corrected but not reduced to standard magnitudes. Fourier analysis (Harris et al., 1989) yielded a period solution from the folded datasets and then independently verified with *Peranso* (Vannmunster 2006) using ANOVA (Schwarzenberg-Czerny, 1996). Phased lightcurve data are available upon request (mail@underoakobservatory.com).

423 Diotima. This is a relatively large ($D = 209 \pm 5$ km) but somewhat dark ($p_V = 0.0515$) taxonomic type C main-belt asteroid that was discovered in 1896 by A. Charlois. Schober (1983) first proposed a synodic period of 8 ± 0.33 h and the possible existence of a satellite. Both of these eventually proved to be inaccurate.



Further refinements to the now accepted synodic period (4.775 h) were reported by Zappala et al. (1985), Dotto et al. (1995), Dymock (2005), Fauerbach and Bennett (2005), and Chiorny et al. (2007). A shape model for 423 Diotima was published by Āurech et al (2007). At UO, images (clear filter; 60 s) were taken over four nights from 2016 Feb 28 thru Mar 3.


Fourier analysis of the 619 lightcurve data points produced a best fit at 4.775 ± 0.001 h. The maximum peak-to-peak amplitude of 0.16 mag observed during the 2016 apparition was consistent with those reported in the Asteroid Lightcurve Database (LCDB; Warner et al., 2009).

925 Alphonsina was discovered in 1920 by J. Solà. It is a moderately sized (57.5 ± 0.4 km) type S asteroid with a somewhat inclined orbit ($i = 21.1^\circ$). Harris and Young (1989) published the first lightcurve with a period solution of 7.880 ± 0.001 h. Additional studies by Hanuš and Āurech (2010) and Hanuš et al (2011) established a *sidereal* period of 7.87754 ± 0.00005 h. This

Number	Name	20yy mm/dd	Pts	Phase	L _{PAB}	B _{PAB}	Period(h)	P.E.	Amp	A.E.
423	Diotima	16/02/28-03/03	619	15.4,16.0	108	9	4.775	0.001	0.16	0.01
925	Alphonsia	17/01/02-02/17	1139	21.8	54	19	7.879	0.001	0.57	0.02

Table I. Observing circumstances and results. Pts is the number of data points. The phase angle is given for the first and last date. L_{PAB} and B_{PAB} are the approximate phase angle bisector longitude and latitude at mid-date range (see Harris et al., 1984).

asteroid has also been shape modeled using a combination of lightcurve inversion and occultation silhouettes (Durech et al, 2011). At UO, images (I_c filter; 60 s) were taken over seven nights from 2017 Jan 2 thru Feb 17.



Fourier analysis of the 1139 lightcurve data points produced a best fit at 7.879 ± 0.001 h. The maximum peak-to-peak amplitude of 0.57 mag observed during this most recent apparition was significantly higher than those of 0.11-0.31 mag reported in the Asteroid Lightcurve Database (Warner et al., 2009).

Acknowledgements

Many thanks to the SAO/NASA Astrophysics Data System, the JPL Small-Body Database Browser and the Asteroid Lightcurve Database (LCDB; Warner et al., 2009), all of which were essential to preparing this report.

References

- Alton, K.B. (2013). "Simultaneous CCD Photometry of Two Eclipsing Binary Stars in Pegasus – Part 1: KW Pegasi." *JAAVSO* **41**, 97-113.
- Chiorny, V.G., Shevchenko, V.G., Krugly, Yu. N., Velichko, F.P., and Gaftonyuk, N.M. (2007). "Photometry of asteroids: Lightcurves of 24 asteroids obtained in 1993-2005." *Planetary and Space Science* **55**, 986-997.
- Dotto, E., De Angelis, G., Di Martino, M., Barucci, M.A., Fulchignoni, M., De Sanctis, G. and Burchi, R. (1995). "Pole Orientation and Shape of 12 Asteroids." *Icarus* **117**, 313-327.
- Đurech, J., Kaasalainen, M., Marciniak, A., Allen, W.H., Behrend, R., Bembrick, C., Bennett, T., Bernasconi, L., Berthier, J., Bolt, G., Boroumand, S., Crespo da Silva, L., Crippa, R., Crow, M., Durkee, R. and 27 colleagues (2007). "Physical models of ten asteroids from an observers' collaboration network." *Astron. Astrophys.* **465**, 331-337.
- Đurech, J., Kaasalainen, M., Herald, D., Dunham, D., Timerson, B., Hanuš, J., Frappa, E., Talbot, J., Hayamizu, T., Warner, B. D., Pilcher, F., Galád, A. (2011). "Combining asteroid models derived by lightcurve inversion with asteroidal occultation silhouettes." *Icarus* **214**, 652-670.
- Dymock, R. (2005). "Lightcurve of 423 Diotima." *Minor Planet Bulletin* **32**, No.3, 52.
- Fauerbach, M., Bennett, T. (2005). "Photometric lightcurve observations of 125 Liberatrix, 218 Bianca, 423 Diotima, 702 Alauda, 1963 Bezovec and (5849) 1990 HF1." *Minor Planet Bulletin* **32**, 80-81.
- Harris, A.W., Young, J.W., Scaltriti, F., Zappala, V. (1984). "Lightcurves and phase relations of the asteroids 82 Alkmene and 444 Gytis." *Icarus* **57**, 251-258.
- Harris, A.W., Young, J.W. (1989). "Asteroid lightcurve observations from 1979-1981." *Icarus*, **81**, 365-374.
- Harris, A.W., Young, J.W., Bowell, E., Martin, L.J., Millis, R. L., Poutanen, M., Scaltriti, F., Zappala, V., Schober, H.J., Debehogne, H., Zeigler, K. (1989). "Photoelectric Observations of Asteroids 3, 24, 60, 261, and 863." *Icarus* **77**, 171-186.
- Hanuš, J., Ďurech, J. (2010). "New asteroid models based on combined dense and sparse photometry." *Bull. Am. Astron. Soc.* **42**, 1035.
- Hanuš, J., Ďurech, J., Brožl, M., Warner, B.D., Pilcher, F., Stephens, R., Oey, J., Bernasconi, L., Casulli, S., Behrend, R., Polishook, D., Henych, T., Lehký, M., Yoshida, F., Ito, T. (2011). "A study of asteroid pole-latitude distribution based on an extended set of shape models derived by the lightcurve inversion method." *Astron. and Astrophys.* **530**, A134.
- Schober, H.J. (1983). "The large C-type asteroid 423 Diotima: rotation period, lightcurve and implications for a possible satellite." *Astron. Astrophys.* **127**, 301-303.
- Schwarzenberg-Czerny, A. (1996). "Fast and Statistically Optimal Period Search in Uneven Sampled Observations." *Astrophys J.* **460**, L107.
- Vannmunster, T. (2006). *Peranso v2.5*, Period Analysis Software. CBA Belgium Observatory.
- Warner, B.D., Harris, A.W. and Pravec, P. (2009). "The asteroid lightcurve database." *Icarus* **202**, 134-146.
- Warner, B.D. (2015). MPO Software, *MPO Canopus v10.7.1.3*. Bdw Publishing, Eaton, CO.
- Zappalà, V., Di Martino, M., Hanslmeier, A., Schober, H.J. (1985). "New cases of ambiguity among large asteroids' spin rates." *Astron. Astrophys.* **147**, 35-38.